

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57194 A2

(51) International Patent Classification⁷: C12N 9/00

(21) International Application Number: PCT/US01/03471

(22) International Filing Date: 2 February 2001 (02.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/179,982	3 February 2000 (03.02.2000)	US
60/183,542	18 February 2000 (18.02.2000)	US
60/213,124	22 June 2000 (22.06.2000)	US
60/220,970	26 July 2000 (26.07.2000)	US
09/657,986	8 September 2000 (08.09.2000)	US
60/234,840	22 September 2000 (22.09.2000)	US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/179,982 (CIP)
Filed on	3 February 2000 (03.02.2000)
US	60/183,542 (CIP)
Filed on	18 February 2000 (18.02.2000)
US	60/213,124 (CIP)
Filed on	22 June 2000 (22.06.2000)
US	60/220,970 (CIP)
Filed on	26 July 2000 (26.07.2000)
US	09/657,986 (CIP)
Filed on	8 September 2000 (08.09.2000)
US	60/234,840 (CIP)
Filed on	22 September 2000 (22.09.2000)

(71) Applicant (for all designated States except US): CORVAS INTERNATIONAL, INC. [US/US]; 3030 Science Park Road, San Diego, CA 92121 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MADISON, Edwin, L. [US/US]; 501 Stratford Court, Del Mar, CA 92014 (US). ONG, Edgar, O. [CA/US]; 8328 Regents Road #3C, San Diego, CA 92122 (US). YEH, Jiunn-cheru [—/US]; 3128-E Via Alicante Drive, La Jolla, CA 92037 (US).

(74) Agents: SEIDMAN, Stephanie, L. et al.; Heller Ehrman White & McAuliffe LLP, Suite 700, 4250 Executive Square, La Jolla, CA 92037 (US).

(81) Designated States (national): AE, AG, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE PROTEASES, THE ENCODED PROTEINS AND METHODS BASED THEREON

(57) Abstract: Provided herein are polypeptides that include the protease domain of a type II transmembrane serine protease (MTSP) as a single chain. Methods using the polypeptides to identify compounds that modulate the protease activity of an MTSP are provided. Also provided are MTSPs designated MTSP3 and MTSP4 and a form of an MTSP designated MTSP6.

WO 01/57194 A2



-1-

**NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE
PROTEASES, THE ENCODED PROTEINS AND METHODS BASED THEREON
RELATED APPLICATIONS**

- Benefit of priority is claimed to U.S. provisional application Serial No. 60/179,982, to Edwin L. Madison and Edgar O. Ong, filed February 3, 2000, entitled "NUCLEOTIDE AND PROTEIN SEQUENCES OF A TRANSMEMBRANE SERINE PROTEASE AND METHODS BASED THEREOF"; to U.S. provisional application Serial No. 60/183,542, to Edwin L. Madison and Edgar O. Ong, filed February 18, 2000, entitled "NUCLEOTIDE AND PROTEIN SEQUENCES OF A TRANSMEMBRANE SERINE PROTEASE AND METHODS BASED THEREOF"; to U.S. provisional application Serial No. 60/213,124, to Edwin L. Madison and Edgar O. Ong, filed June 22, 2000, entitled "NUCLEOTIDE AND PROTEIN SEQUENCES OF A TRANSMEMBRANE SERINE PROTEASE AND METHODS BASED THEREOF"; to U.S. provisional application Serial No. 60/220,970, to Edwin L. Madison and Edgar O. Ong, filed July 26, 2000, entitled "NUCLEOTIDE AND PROTEIN SEQUENCES OF A TRANSMEMBRANE SERINE PROTEASE AND METHODS BASED THEREOF"; and to U.S. provisional application Serial No. 60/234,840 to Edwin L. Madison, Edgar O. Ong and Jiunn-Chern Yeh, filed September 22, 2000, entitled "NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE PROTEASES, THE ENCODED PROTEINS AND METHODS BASED THEREON" is claimed herein. Benefit of priority is also claimed to U.S. application Serial No. 09/657,968, to Edwin L. Madison, Joseph Edward Semple, Gary Samuel Coombs, John Eugene Reiner, Edgar O. Ong, Gian Luca Araldi, filed September 8, 2000, entitled "INHIBITORS OF SERINE PROTEASE ACTIVITY OF MATRIPTASE OR MTSP1". This application is a continuation-in-part of U.S. application Serial No. 09/657,986.

- This application is related to U.S. provisional application Serial No. 60/166,391 to Edwin L. Madison and Edgar O. Ong, filed November 18, 1999 entitled "NUCLEOTIDE AND PROTEIN SEQUENCES OF PROTEASE DOMAINS OF ENDOTHELIASE AND METHODS BASED THEREON". This a application is also

RECTIFIED SHEET (RULE 91)

-2-

related to International PCT application No. PCT/US00/31803, filed November 17, 2000.

Where permitted, the above-noted provisional applications, patent application and International PCT application are incorporated by reference in their entirety. All patents, applications, published applications and other publications and sequences from GenBank and other data bases referred to herein are incorporated by reference in their entirety.

FIELD OF INVENTION

Nucleic acid molecules that encode proteases and portions thereof, particularly protease domains are provided. Also provided are prognostic, diagnostic and therapeutic methods using the proteases and domains thereof and the encoding nucleic acid molecules.

BACKGROUND OF THE INVENTION AND OBJECTS THEREOF

Cancer a leading cause of death in the United States, developing in one in three Americans; one of every four Americans dies of cancer. Cancer is characterized by an increase in the number of abnormal neoplastic cells, which proliferate to form a tumor mass, the invasion of adjacent tissues by these neoplastic tumor cells, and the generation of malignant cells that metastasize via the blood or lymphatic system to regional lymph nodes and to distant sites.

Among the hallmarks of cancer is a breakdown in the communication among tumor cells and their environment. Normal cells do not divide in the absence of stimulatory signals, cease dividing in the presence of inhibitory signals. Growth-stimulatory and growth-inhibitory signals, are routinely exchanged between cells within a tissue. In a cancerous, or neoplastic, state, a cell acquires the ability to "override" these signals and to proliferate under conditions in which normal cells do not grow.

In order to proliferate tumor cells acquire a number of distinct aberrant traits reflecting genetic alterations. The genomes of certain well-studied tumors carry several different independently altered genes, including activated oncogenes and inactivated tumor suppressor genes. Each of these genetic changes appears to be responsible for imparting some of the traits that, in the aggregate, represent the full neoplastic phenotype.

-3-

A variety of biochemical factors have been associated with different phases of metastasis. Cell surface receptors for collagen, glycoproteins such as laminin, and proteoglycans, facilitate tumor cell attachment, an important step in invasion and metastases. Attachment triggers the release of degradative enzymes which facilitate the penetration of tumor cells through tissue barriers. Once the tumor cells have entered the target tissue, specific growth factors are required for further proliferation. Tumor invasion (or progression) involves a complex series of events, in which tumor cells detach from the primary tumor, break down the normal tissue surrounding it, and migrate into a blood or lymphatic vessel to be carried to a distant site. The breaking down of normal tissue barriers is accomplished by the elaboration of specific enzymes that degrade the proteins of the extracellular matrix that make up basement membranes and stromal components of tissues.

A class of extracellular matrix degrading enzymes have been implicated in tumor invasion. Among these are the matrix metalloproteinases (MMP). For example, the production of the matrix metalloproteinase stromelysin is associated with malignant tumors with metastatic potential (see, *e.g.*, McDonnell *et al.* (1990) *Smnrs. in Cancer Biology* 1:107-115; McDonnell *et al.* (1990) *Cancer and Metastasis Reviews* 9:309-319).

The capacity of cancer cells to metastasize and invade tissue is facilitated by degradation of the basement membrane. Several proteinase enzymes, including the MMPs, have been reported to facilitate the process of invasion of tumor cells. MMPs are reported to enhance degradation of the basement membrane, which thereby permits tumorous cells to invade tissues. For example, two major metalloproteinases having molecular weights of about 70 kDa and 92 kDa appear to enhance ability of tumor cells to metastasize.

Type II Transmembrane Serine Proteases (TTSPs)

In addition to the MMPs, serine proteases have been implicated in neoplastic disease progression. Most serine proteases, which are either secreted enzymes or are sequestered in cytoplasmic storage organelles, have roles in blood coagulation, wound healing, digestion, immune responses and tumor invasion and metastasis. A class cell surface proteins designated type II

-4-

transmembrane serine proteases, which are membrane-anchored proteins with N-terminal extracellular domains, has been identified. As cell surface proteins, they are positioned to play a role in intracellular signal transduction and in mediating cell surface proteolytic events.

- 5 Cell surface proteolysis is a mechanism for the generation of biologically active proteins that mediate a variety of cellular functions. These membrane-anchored proteins, include a disintegrin-like and metalloproteinase (ADAM) and membrane-type matrix metalloproteinase (MT-MMP). In mammals, at least 17 members of the family are known, including seven in humans (see, Hooper *et al.*
- 10 (2001) *J. Biol. Chem.* 276:857-860). These include: corin (accession nos. AF133845 and AB013874; see, Yan *et al.* (1999) *J. Biol. Chem.* 274:14926-14938; Tomia *et al.* (1998) *J. Biochem.* 124:784-789; Uan *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:8525-8529); enterpeptidase (also designated enterokinase; accession no. U09860 for the human protein; see, Kitamoto *et al.*
- 15 (1995) *Biochem.* 27: 4562-4568; Yahagi *et al.* (1996) *Biochem. Biophys. Res. Commun.* 219:806-812; Kitamoto *et al.* (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91:7588-7592; Matsushima *et al.* (1994) *J. Biol. Chem.* 269:19976-19982;); human airway trypsin-like protease (HAT; accession no. AB002134; see Yamaoka *et al.* *J. Biol. Chem.* 273:11894-11901); MTSP1 and matriptase (also
- 20 called TADG-15; see SEQ ID Nos. 1 and 2; accession nos. AF133086/AF118224, AF04280022; Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-1161; Lin *et al.* (1999) *J. Biol. Chem.* 274:18231-18236; Takeuchi *et al.* (2000) *J. Biol. Chem.* 275:26333-26342; and Kim *et al.* (1999) *Immunogenetics* 49:420-429); hepsin (see, accession nos. M18930, AF030065,
- 25 X70900; Leytus *et al.* (1988) *Biochem.* 27: 11895-11901; Vu *et al.* (1997) *J. Biol. Chem.* 272:31315-31320; and Farley *et al.* (1993) *Biochem. Biophys. Acta* 1173:350-352; and see, U.S. Patent No. 5,972,616); TMPRS2 (see, Accession Nos. U75329 and AF113596; Paoloni-Giacobino *et al.* (1997) *Genomics* 44:309-320; and Jacquinet *et al.* (2000) *FEBS Lett.* 468: 93-100); and TMPRSS4 (see,
- 30 Accession No. NM 016425; Wallrapp *et al.* (2000) *Cancer* 60:2602-2606).

Serine proteases, including transmembrane serine proteases, have been implicated in processes involved in neoplastic development and progression.

-5-

While the precise role of these proteases has not been elaborated, serine proteases and inhibitors thereof are involved in the control of many intra- and extracellular physiological processes, including degradative actions in cancer cell invasion, metastatic spread, and neovascularization of tumors, that are involved
5 in tumor progression. It is believed that proteases are involved in the degradation of extracellular matrix (ECM) and contribute to tissue remodeling, and are necessary for cancer invasion and metastasis. The activity and/or expression of some proteases have been shown to correlate with tumor progression and development.

10 For example, a membrane-type serine protease MTSP1 (also called matriptase; see SEQ ID Nos. 1 and 2 from U.S. Patent No. 5,972,616; and GenBank Accession No. AF118224; (1999) *J. Biol. Chem.* 274:18231-18236; U.S. Patent No. 5,792,616; see, also Takeuchi (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-1161) that is expressed in epithelial cancer and normal tissue
15 (Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. USA*, 96(20):11054-61) has been identified. Matriptase was originally identified in human breast cancer cells as a major gelatinase (see, U.S. Patent No. 5,482,848), a type of matrix metalloprotease (MMP). It has been proposed that it plays a role in the metastasis of breast cancer. Its primary cleavage specificity is Arg-Lys residues.
20 Matriptase also is expressed in a variety of epithelial tissues with high levels of activity and/or expression in the human gastrointestinal tract and the prostate.

Prostate-specific antigen (PSA), a kallikrein-like serine protease, degrades extracellular matrix glycoproteins fibronectin and laminin, and, has been postulated to facilitate invasion by prostate cancer cells (Webber *et al.* (1995)
25 *Clin. Cancer Res.*, 1(10):1089-94). Blocking PSA proteolytic activity with PSA-specific monoclonal antibodies results in a dose-dependent decrease *in vitro* in the invasion of the reconstituted basement membrane Matrigel by LNCaP human prostate carcinoma cells which secrete high levels of PSA.

Hepsin, a cell surface serine protease identified in hepatoma cells, is
30 overexpressed in ovarian cancer (Tanimoto *et al.* (1997) *Cancer Res.*, 57(14):2884-7). The hepsin transcript appears to be abundant in carcinoma tissue and is almost never expressed in normal adult tissue, including normal

-6-

ovary. It has been suggested that hepsin is frequently overexpressed in ovarian tumors and therefore may be a candidate protease in the invasive process and growth capacity of ovarian tumor cells.

A serine protease-like gene, designated normal epithelial cell-specific 1 (NES1) (Liu et al., *Cancer Res.*, 56(14):3371-9 (1996)) has been identified. Although expression of the NES1 mRNA is observed in all normal and immortalized nontumorigenic epithelial cell lines, the majority of human breast cancer cell lines show a drastic reduction or a complete lack of its expression. The structural similarity of NES1 to polypeptides known to regulate growth factor activity and a negative correlation of NES1 expression with breast oncogenesis suggest a direct or indirect role for this protease-like gene product in the suppression of tumorigenesis.

Hence transmembrane serine proteases appear to be involved in the etiology and pathogenesis of tumors. There is a need to further elucidate their role in these processes and to identify additional transmembrane proteases. Therefore, it is an object herein to provide transmembrane serine protease (MTSP) proteins and nucleic acids encoding such MTSP proteases that are involved in the regulation of or participate in tumorigenesis and/or carcinogenesis. It is also an object herein to provide prognostic, diagnostic, therapeutic screening methods using the such proteases and the nucleic acids encoding such proteases.

SUMMARY OF THE INVENTION

Provided herein are isolated protease domains of the Transmembrane Serine Protease family, particularly the Type II Transmembrane Serine Protease (TTSP) family (also referred to herein as MTSPs), and more particularly TTSP family members whose functional activity differs in tumor cells from non-tumor cells in the same tissue. For example, the MTSPs include those that are activated and/or expressed in tumor cells at different levels, typically higher, from non-tumor cells; and those from cells in which substrates therefor differ in tumor cells from non-tumor cells or otherwise alter the specificity of the MTSP.

The MTSP family as intended herein does not include any membrane anchored or spanning proteases that are expressed on endothelial cells.

-7-

Included among the MTSPs are several heretofore unidentified MTSP family members, designated herein as MTSP3 and MTSP4 and a new form of a protein designated herein as MTSP6. In addition to the protease domains of each of MTSP3 and MTSP4, the full-length proteins, including those that results from
5 splice variants, zymogens and activated forms, and uses thereof, are also provided.

The protease domains as provided herein are single-chain polypeptides, with an N-terminus (such as IV, VV, IL and II) generated at the cleavage site (generally having the consensus sequence R↓VVGG, R↓IVGG, R↓IVNG,
10 R↓ILGG, R↓VGLL, R↓ILGG or a variation thereof; an N-terminus R↓V or R↓I, where the arrow represents the cleavage point) when the zymogen is activated. To identify the protease domain an RI should be identified, and then the following amino acids compared to the above noted motif.

The protease domains generated herein, however, do not result from
15 activation, which produces a two chain activated product, but rather are single chain polypeptides with the N-terminus include the consensus sequence ↓VVGG, ↓IVGG, ↓VGLL, ↓ILGG or ↓IVNG or other such motif at the N-terminus. As shown herein, such polypeptides, although not the result of activation and not double-chain forms, exhibit proteolytic (catalytic) activity.
20 These protease domain polypeptides are used in assays to screen for agents that modulate the activity of the MTSP. Such assays are also provided herein. In exemplary assays, the affects of test compounds in the ability of a protease domains to proteolytically cleave a known substrate, typically a fluorescently, chromogenically or otherwise detectably labeled substrate, are assessed.
25 Agents, generally compounds, particularly small molecules, that modulate the activity of the protease domain are candidate compounds for modulating the activity of the MTSP. The protease domains can also be used to produce single-chain protease-specific antibodies. The protease domains provided herein include, but are not limited to, the single chain region having an N-terminus at
30 the cleavage site for activation of the zymogen, through the C-terminus, or C-terminal truncated portions thereof that exhibit proteolytic activity as a single-chain polypeptide in *in vitro* proteolysis assays, of any MTSP family member,

-8-

preferably from a mammal, including and most preferably human, that, for example, is expressed in tumor cells at different levels from non-tumor cells, and that is not expressed on an endothelial cell. These include, but are not limited to : MTSP1 (or matriptase), MTSP3, MTSP4 and MTSP6. Other MTSP protease domains of interest herein, particularly for use in *in vitro* drug screening proteolytic assays, include, but are not limited to: corin (accession nos. AF133845 and AB013874; see, Yan *et al.* (1999) *J. Biol. Chem.* 274:14926-14938; Tomia *et al.* (1998) *J. Biochem.* 124:784-789; Uan *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97:8525-8529; SEQ ID Nos. 61 and 62 for the human protein); enterpeptidase (also designated enterokinase; accession no. U09860 for the human protein; see, Kitamoto *et al.* (1995) *Biochem.* 27: 4562-4568; Yahagi *et al.* (1996) *Biochem. Biophys. Res. Commun.* 219:806-812; Kitamoto *et al.* (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91:7588-7592; Matsushima *et al.* (1994) *J. Biol. Chem.* 269:19976-19982; see SEQ ID Nos. 63 and 64 for the human protein); human airway trypsin-like protease (HAT; accession no. AB002134; see Yamaoka *et al.* *J. Biol. Chem.* 273:11894-11901; SEQ ID Nos. 65 and 66 for the human protein); hepsin (see, accession nos. M18930, AF030065, X70900; Leytus *et al.* (1988) *Biochem.* 27: 11895-11901; Vu *et al.* (1997) *J. Biol. Chem.* 272:31315-31320; and Farley *et al.* (1993) *Biochem. Biophys. Acta* 1173:350-352; SEQ ID Nos. 67 and 68 for the human protein); TMPRS2 (see, Accession Nos. U75329 and AF113596; Paoloni-Giacobino *et al.* (1997) *Genomics* 44:309-320; and Jacquinet *et al.* (2000) *FEBS Lett.* 468: 93-100; SEQ ID Nos. 69 and 70 for the human protein) TMPRSS4 (see, Accession No. NM 016425; Wallrapp *et al.* (2000) *Cancer* 60:2602-2606; SEQ ID Nos. 71 and 72 for the human protein); and TADG-12 (also designated MTSP6, see SEQ ID Nos. 11 and 12; see International PCT application No. WO 00/52044, which claims priority to U.S. application Serial No. 09/261,416).

Also provided are muteins of the single chain protease domains and MTSPs, particularly muteins in which the Cys residue in the protease domain that is free (*i.e.*, does not form disulfide linkages with any other Cys residue in the protein) is substituted with another amino acid substitution, preferably with a conservative amino acid substitution or a substitution that does not eliminate the

RECTIFIED SHEET (RULE 91)

-9-

activity, and muteins in which a glycosylation site(s) is eliminated. Muteins in which other conservative amino acid substitutions in which catalytic activity is retained are also contemplated (see, *e.g.*, Table 1, for exemplary amino acid substitutions). See, also, Figure 4, which identifies the free Cys residues in

5 MTSP3, MTSP4 and MTSP6.

Hence, provided herein are members of a family of transmembrane serine protease (MTSP) proteins, and functional domains, especially protease (or catalytic) domains thereof, muteins and other derivatives and analogs thereof. Also provided herein are nucleic acids encoding the MTSPs.

10 Exemplary MTSPs (see, *e.g.*, SEQ ID No. 1-12, 49 and 50) are provided herein, as are the single chain protease domains thereof as follows: SEQ ID Nos. 1, 2, 49 and 50 set forth amino acid and nucleic acid sequences of MTSP1 and the protease domain thereof; SEQ ID No. 3 sets forth the MTSP3 nucleic acid sequence and SEQ ID No. 4 the encoded MTSP3 amino acids; SEQ ID No. 5
15 MTSP4 a nucleic acid sequence of the protease domain and SEQ ID No. 6 the encoded MTSP4 amino acid protease domain; SEQ ID No. 7 MTSP4-L a nucleic acid sequence and SEQ ID No. 8 the encoded MTSP4-L amino acid sequence; SEQ ID No. 9 an MTSP4-S encoding nucleic acid sequence and SEQ ID No. 10 the encoded MTSP4-S amino acid sequence; and SEQ ID No. 11 an MTSP6
20 encoding nucleic acid sequence and SEQ ID No. 12 the encoded MTSP6 amino acid sequence. The single chain protease domains of each are delineated below.

Nucleic acid molecules that encode a single-chain protease domain or catalytically active portion thereof are provided. Also provided are nucleic acid molecules that hybridize to such *MTSP* encoding nucleic acid along their full
25 length and encode the protease domain or portion thereof are provided. Hybridization is preferably effected under conditions of at least low, generally at least moderate, and often high stringency.

Additionally provided herein are antibodies that specifically bind to the MTSPs, cells, combinations, kits and articles of manufacture that contain the
30 nucleic acid encoding the MTSP and/or the MTSP. Further provided herein are prognostic, diagnostic, therapeutic screening methods using MTSPs and the nucleic acids encoding MTSP. Also provided are transgenic non-human animals

-10-

bearing inactivated genes encoding the MTSP and bearing the genes encoding the MTSP under non-native promotor control. Such animals are useful in animal models of tumor initiation, growth and/or progression models.

Provided herein are members of a family of membrane serine proteases (MTSP) that are expressed in certain tumor or cancer cells such lung, prostate, colon and breast cancers. In particular, it is shown herein, that MTSPs, particularly, MTSP3, MTSP4 and MTSP6 are expressed in lung carcinoma, breast carcinoma, colon adenocarcinoma and/or ovarian carcinomas as well as in certain normal cells and tissues (see *e.g.*, EXAMPLES for tissue-specific expression profiles of each protein exemplified herein). The MTSPs that are of particular interest herein, are those that are expressed in tumor cells, for example, those that appear to be expressed at different levels in tumor cells from normal cells, or whose functional activity is different in tumor cells from normal cells, such as by an alteration in a substrate therefor, or a cofactor. Hence the MTSP provided herein can serve as diagnostic markers for certain tumors. The level of activated MTSP3, MTSP4 and MTSP6 can be diagnostic of prostate cancer. In addition, MTSP4 is expressed and/or activated in lymphomas, leukemias, lung cancer, breast, prostate and colon cancers. MTSP6 is activated and/or expressed in breast, lung, prostate, colon and ovarian cancers. Furthermore, compounds that modulate the activity of these MTSPs, as assessed by the assays provided herein, particularly the *in vitro* proteolytic assays that use the single chain protease domains, are potential therapeutic candidates for treatment of various malignancies and neoplastic disease.

Also provided herein are methods of modulating the activity of the MTSPs and screening for compounds that modulate, including inhibit, antagonize, agonize or otherwise alter the activity of the MTSPs. Of particular interest is the extracellular domain of these MTSPs that includes the proteolytic (catalytic) portion of the protein.

MTSP proteins, including, but not limited to, MTSP3, MTSP4, and MTSP6, including splice variants thereof, and nucleic acids encoding MTSPs, and domains, derivatives and analogs thereof are provided herein. Single chain protease domains, in the N-terminal is that which would be generated by

RECTIFIED SHEET (RULE 91)

-11-

activation of the zymogen, from any MTSP, particularly those that are not expressed in endothelial cells and that are expressed in tumor cells are also provided.

Antibodies that specifically bind to the MTSP, particularly the single chain
5 protease domain, and any and all forms of MTSP3 and MTSP4, and cells, combinations, kits and articles of manufacture containing the MTSP proteins, domains thereof, or encoding nucleic acids are also provided herein. Transgenic non-human animals bearing inactivated genes encoding the MTSP and bearing the genes encoding the MTSP under a non-native promotor control are
10 additionally provided herein. Also provided are nucleic acid molecules encoding each of the MTSPs and domains thereof.

Also provided are plasmids containing any of the nucleic acid molecules provided herein. Cells containing the plasmids are also provided. Such cells include, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells,
15 insect cells and animal cells.

Also provided is a method of producing a MTSP by growing the above-described cells under conditions whereby the MTSP is expressed by the cells, and recovering the expressed MTSP protein. Methods for isolating nucleic acid encoding other MTSPs are also provided.

20 Also provided are cells, preferably eukaryotic cells, such as mammalian cells and yeast cells, in which the MTSP protein, preferably MTSP3 and MTSP4, is expressed in the surface of the cells. Such cells are used in drug screening assays to identify compounds that modulate the activity of the MTSP protein. These assays including *in vitro* binding assays, and transcription based assays in
25 which signal transduction mediated by the MTSP is assessed.

Further provided herein are prognostic, diagnostic and therapeutic screening methods using the MTSP and the nucleic acids encoding MTSP. In particular, the prognostic, diagnostic and therapeutic screening methods are used for preventing, treating, or for finding agents useful in preventing or
30 treating, tumors or cancers such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.

-12-

Also provided are methods for screening for compounds that modulate the activity of any MTSP. The compounds are identified by contacting them with the MTSP and a substrate for the MTSP. A change in the amount of substrate cleaved in the presence of the compounds compared to that in the
5 absence of the compound indicates that the compound modulates the activity of the MTSP. Such compounds are selected for further analyses or for use to modulate the activity of the MTSP, such as inhibitors or agonists. The compounds can also be identified by contacting the substrates with a cell that expresses the MTSP or the extracellular domain or proteolytically active portion
10 thereof. For assays in which the extracellular domain or a proteolytically active portion thereof is employed, the MTSP is any MTSP that is expressed on cells, other than endothelial cells, including, but not limited to MTSP1, MTSP3, MTSP4 and MTSP6.

Also provided herein are modulators of the activity of the MTSP,
15 especially the modulators obtained according to the screening methods provide herein. Such modulators may have use in treating cancerous conditions, and other neoplastic conditions.

Pharmaceutical composition containing the protease domains of an MTSP protein, and the MTSP proteins, MTSP3, MTSP4 and MTSP6 are provided herein
20 in a pharmaceutically acceptable carrier or excipient are provided herein.

Also provided are articles of manufacture that contain the MTSP proteins and protease domains of MTSPs in single chain form. The articles contain a) packaging material; b) the polypeptide (or encoding nucleic acid), particularly the single chain protease domain thereof; and c) a label indicating that the article is
25 for using ins assays for identifying modulators of the activities of an MTSP protein is provided herein.

Conjugates containing a) a MTSP protease domain in single chain from; and b) a targeting agent linked to the MTSP directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the conjugate; ii)
30 attachment of the conjugate to a surface; iii) detection of the conjugate; or iv) targeted delivery to a selected tissue or cell, is provided herein. The conjugate

-13-

can contain a plurality of agents linked thereto. The conjugate can be a chemical conjugate; and it can be a fusion protein.

In yet another embodiment, the targeting agent is a protein or peptide fragment. The protein or peptide fragment can include a protein binding
5 sequence, a nucleic acid binding sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence.

Method of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP, particularly an MTSP3, MTSP4 or MTSP 6, in a subject is provided. The method can be practiced by measuring the level of the
10 DNA, RNA, protein or functional activity of the MTSP. An increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP, relative to the level of the DNA, RNA, protein or functional activity found in an analogous sample not having the disease or disorder (or other suitable control) is
15 indicative of the presence of the disease or disorder in the subject or other relative any other suitable control.

Combinations are provided herein. The combination can include: a) an inhibitor of the activity of an MTSP; and b) an anti-cancer treatment or agent. The MTSP inhibitor and the anti-cancer agent can be formulated in a single
20 pharmaceutical composition or each is formulated in a separate pharmaceutical composition. The MTSP inhibitor can be an antibody or a fragment or binding portion thereof against the MTSP, such as an antibody that specifically binds to the protease domain, an inhibitor of the MTSP production, or an inhibitor of the MTSP membrane-localization or an inhibitor of MTSP activation. Other MTSP
25 inhibitors include, but are not limited to, an antisense nucleic acid encoding the MTSP, particularly a portion of the protease domain; a nucleic acid encoding at least a portion of a gene encoding the MTSP with a heterologous nucleotide sequence inserted therein such that the heterologous sequence inactivates the biological activity encoded MTSP or the gene encoding it. The portion of the
30 gene encoding the MTSP preferably flanks the heterologous sequence to promote homologous recombination with a genomic gene encoding the MTSP.

Also, provided are methods for treating or preventing a tumor or cancer in a mammal by administering to a mammal an effective amount of an inhibitor of

-14-

an MTSP3, MTSP4 or MTSP6, whereby the tumor or cancer is treated or prevented. The MTSP inhibitor used in the treatment or for prophylaxis is administered with a pharmaceutically acceptable carrier or excipient. The mammal treated can be a human. The treatment or prevention method can additionally include administering an anti-cancer treatment or agent simultaneously with or subsequently or before administration of the MTSP inhibitor.

Also provided is a recombinant non-human animal in which an endogenous gene of an MTSP has been deleted or inactivated by homologous recombination or insertional mutagenesis of the animal or an ancestor thereof. A recombinant non-human animal is provided herein, where the gene of an MTSP is under control of a promoter that is not the native promoter of the gene or that is not the native promoter of the gene in the non-human animal or where the nucleic acid encoding the MTSP is heterologous to the non-human animal and the promoter is the native or a non-native promoter.

Also provided are methods of treatments of tumors by administering a prodrug that is activated by an MTSP that is expressed or active in tumor cells, particularly those in which its functional activity in tumor cells is greater than in none-tumor cells. The prodrug is administered and, upon administration, active MTSP expressed on cells cleaves the prodrug and releases active drug in the vicinity of these cells. The active anti-cancer drug accumulates in the vicinity of the tumor. This is particularly useful in instances in which an MTSP is expressed or active in greater quantity, higher level or predominantly in tumor cells compared to other cells.

25 BRIEF DESCRIPTION OF DRAWINGS

Figure 1 illustrates the domain organization of the MTSP3;

Figure 2 illustrates the domain organization of the MTSP4 splice variants and domains thereof; MTSP4-L includes a transmembrane domain, a CUB domain, a low density lipoprotein receptor (LDLR) domains, and a serine protease catalytic domain; MTSP4-S lacking the portion between amino acids 136-279.

Figure 3 depicts the domain organization of MTSP6.

-15-

Figure 4 provides an alignment of the C-terminal portions of MTSP3, the two splice variant-encoded forms of MTSP4, and MTSP6, that encompasses the protease domains thereof; the figure shows the cleavage sites, which form the N-terminus of the protease domain of each protein; a potential glycosylation site is noted and the free Cys residues in the protease domain of each are noted (*). Muteins of each protein may be prepared by replacing the residues in the glycosylation site, particularly the N residue, and the free Cys residues, with preferably conservative amino acid residues. Such muteins are also provided herein.

10 DETAILED DESCRIPTION OF THE INVENTION

A. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, applications, published applications and other publications and sequences from GenBank and other data bases referred to herein are incorporated by reference in their entirety.

As used herein, the abbreviations for any protective groups, amino acids and other compounds, are, unless indicated otherwise, in accord with their common usage, recognized abbreviations, or the IUPAC-IUB Commission on Biochemical Nomenclature (see, (1972) Biochem. 11:942-944).

As used herein, serine protease refers to a diverse family of proteases wherein a serine residue is involved in the hydrolysis of proteins or peptides. The serine residue can be part of the catalytic triad mechanism, which includes a serine, a histidine and an aspartic acid in the catalysis, or be part of the hydroxyl/ ϵ -amine or hydroxyl/ α -amine catalytic dyad mechanism, which involves a serine and a lysine in the catalysis.

As used herein, "transmembrane serine protease (MTSP)" refers to a family of transmembrane serine proteases that share common structural features as described herein (see, also Hooper *et al.* (2001) *J. Biol. Chem.* 276:857-860). Thus, reference, for example, to "MTSP" encompasses all proteins encoded by the MTSP gene family, including but are not limited to: MTSP1, MTSP3, MTSP4 and MTSP6, or an equivalent molecule obtained from any other source or that

-16-

has been prepared synthetically or that exhibits the same activity. Other MTSPs include, but are not limited to, corin, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4. Sequences of encoding nucleic molecules and the encoded amino acid sequences of exemplary MTSPs and/or domains thereof are set forth in SEQ ID Nos. 1-12, 49, 50 and 61-72. The term also encompass MTSPs with conservative amino acid substitutions that do not substantially alter activity of each member, and also encompasses splice variants thereof. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Of particular interest are MTSPs of mammalian, including human, origin. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224).

As used herein, a "protease domain of an MTSP" refers to the protease domain of MTSP that is located within the extracellular domain of a MTSP and exhibits serine proteolytic activity. It includes at least the smallest fragment thereof that acts catalytically as a single chain form. Hence it is at least the minimal portion of the extracellular domain that exhibits proteolytic activity as assessed by standard assays *in vitro* assays. Those of skill in this art recognize that such protease domain is the portion of the protease that is structurally equivalent to the trypsin or chymotrypsin fold.

Exemplary MTSP proteins, with the protease domains indicated, are illustrated in Figures 1-3. Smaller portions thereof that retain protease activity are contemplated. The protease domains vary in size and constitution, including insertions and deletions in surface loops. They retain conserved structure, including at least one of the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease domains of proteases. Thus, for purposes herein, the protease domain is a portion of a MTSP, as defined herein, and is homologous to a domain of other MTSPs, such as corin, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2,

-17-

and TMPRSS4, which have been previously identified; it was not recognized, however, that an isolated single chain form of the protease domain could function proteolytically in *in vitro* assays. As with the larger class of enzymes of the chymotrypsin (S1) fold (see, *e.g.*, Internet accessible MEROPS data base),

5 the MTSPs protease domains share a high degree of amino acid sequence identity. The His, Asp and Ser residues necessary for activity are present in conserved motifs. The activation site, which results in the N-terminus of second chain in the two chain forms is has a conserved motif and readily can be identified (see, *e.g.*, amino acids 801-806, SEQ ID No. 62, amino acids 406-

10 410, SEQ ID No. 64; amino acids 186-190, SEQ ID No. 66; amino acids 161-166, SEQ ID No. 68; amino acids 255-259, SEQ ID No. 70; amino acids 190-194, SEQ ID No. 72).

As used herein, the catalytically active domain of an MTSP refers to the protease domain. Reference to the protease domain of an MTSP refers includes

15 the single and double-chain forms of any of these proteins. The zymogen form of each protein is single chain form, which can be converted to the active two chain form by cleavage. The protease domain may also be converted to a two chain form. By active form is meant a form active *in vivo*.

Significantly, it is shown herein, that, at least *in vitro*, the single chain

20 forms of the MTSPs and the catalytic domains or proteolytically active portions thereof (typically C-terminal truncations) thereof exhibit protease activity. Hence provided herein are isolated single chain forms of the protease domains of MTSPs and their use in *in vitro* drug screening assays for identification of agents that modulate the activity thereof.

25 As used herein an MTSP3, whenever referenced herein, includes at least one or all of or any combination of:

a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 3;

a polypeptide encoded by a sequence of nucleotides that

30 hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in SEQ ID No. 3;

RECTIFIED SHEET (RULE 91)

-18-

a polypeptide that comprises the sequence of amino acids set forth as amino acids 205-437 of SEQ ID No. 4;

a polypeptide that comprises a sequence of amino acids having at least about 85% or 90% sequence identity with the sequence of amino acids set forth in SEQ ID No. 4; and/or

a splice variant of the MTSP3 set forth in SEQ ID Nos. 3 and 4.

The MTSP3 may be from any animal, particularly a mammal, and includes but are not limited to, humans, rodents, fowl, ruminants and other animals. The full length zymogen or double chain activated form is contemplated or any domain thereof, including the protease domain, which can be a double chain activated form, or a single chain form.

As used herein an MTSP4, whenever referenced herein, includes at least one or all of or any combination of:

a polypeptide encoded by the sequence of nucleotides set forth in any of SEQ ID No. 5, 7 or 9;

a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in any of SEQ ID Nos. 5, 7 or 9;

a polypeptide that comprises the sequence of amino acids set forth in any of SEQ ID Nos. 6, 8 or 10;

a polypeptide that comprises a sequence of amino acids having at least about 85% or 90% or 95% sequence identity with the sequence of amino acids set forth in SEQ ID No. 6, 8 or 10; and/or

a splice variant of the MTSP4s set forth in SEQ ID Nos. 7-10.

The MTSP4 may be from any animal, particularly a mammal, and includes but are not limited to, humans, rodents, fowl, ruminants and other animals. The full length zymogen or double chain activated form is contemplated or any domain thereof, including the protease domain, which can be a double chain activated form, or a single chain form.

As used herein an MTSP6, whenever referenced herein, includes at least one or all of or any combination of:

RECTIFIED SHEET (RULE 91)

-19-

a polypeptide encoded by the sequence of nucleotides set forth in any of SEQ ID No. 11;

a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of low, moderate or high stringency to the sequence of nucleotides set forth in any of SEQ ID Nos. 11;

a polypeptide that comprises the sequence of amino acids set forth in any of SEQ ID Nos. 12;

a polypeptide that comprises a sequence of amino acids having at least about 90% or 95% or 98% sequence identity with the sequence of amino acids set forth in SEQ ID No. 12; and/or

a splice variant of the MTSP4s set forth in SEQ ID No. 12.

The MTSP6 may be from any animal, particularly a mammal, and includes but are not limited to, humans, rodents, fowl, ruminants and other animals. The full length zymogen or double chain activated form is contemplated or any domain thereof, including the protease domain, which can be a double chain activated form, or a single chain form. Of particular interest herein is the MTSP6 of SEQ ID No. 12.

As used herein, a human protein is one encoded by DNA present in the genome of a human, including all allelic variants and conservative variations as long as they are not variants found in other mammals.

As used herein, a "nucleic acid encoding a protease domain or catalytically active portion of a MTSP" shall be construed as referring to a nucleic acid encoding only the recited single chain protease domain or active portion thereof, and not the other contiguous portions of the MTSP as a continuous sequence.

As used herein, a CUB domain is a motif that mediates protein-protein interactions in complement components C1r/C1s and has also been identified in various proteins involved in developmental processes.

As used herein, catalytic activity refers to the activity of the MTSP as a serine proteases. Function of the MTSP refers to its function in tumor biology, including promotion of or involvement in tumorigenesis, metastasis or carcinogenesis, and also roles in signal transduction.

-20-

As used herein, a "propeptide" or "pro sequence" is sequence of amino acids positioned at the amino terminus of a mature biologically active polypeptide. When so-positioned, the resulting polypeptide is called a zymogen. Zymogens, generally, are biologically inactive and can be converted to mature

5 active polypeptides by catalytic or autocatalytic cleavage of the propeptide from the zymogen. A zymogen is an enzymatically inactive protein that is converted to a proteolytic enzyme by the action of an activator. Cleavage may be effected autocatalytically.

As used herein, "disease or disorder" refers to a pathological condition in

10 an organism resulting from, *e.g.*, infection or genetic defect, and characterized by identifiable symptoms.

As used herein, neoplasm (neoplasia) refers to abnormal new growth, and thus means the same as *tumor*, which may be benign or malignant. Unlike *hyperplasia*, neoplastic proliferation persists even in the absence of the original

15 stimulus.

As used herein, neoplastic disease refers to any disorder involving cancer, including tumor development, growth, metastasis and progression.

As used herein, cancer refers to a general term for diseases caused by any type of malignant tumor.

As used herein, malignant, as applies to tumors, refers to primary tumors

20 that have the capacity of *metastasis* with loss of *growth control* and *positional control*.

As used herein, an anti-cancer agent (used interchangeable with "anti-tumor or anti-neoplastic agent") refers to any agents used in the anti-cancer

25 treatment. These include any agents, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with neoplastic disease, tumor and cancer, and can be used in methods, combinations and compositions provided herein. Non-limiting

30 examples of anti-neoplastic agents include anti-angiogenic agents, alkylating agents, antimetabolite, certain natural products, platinum coordination complexes, anthracenediones, substituted ureas, methylhydrazine derivatives,

-21-

adrenocortical suppressants, certain hormones, antagonists and anti-cancer polysaccharides.

As used herein, a splice variant refers to a variant produced by differential processing of a primary transcript of genomic DNA that results in more than one type of mRNA. Splice variants of MTSPs are provided herein.

As used herein, angiogenesis is intended to broadly encompass the totality of processes directly or indirectly involved in the establishment and maintenance of new vasculature (neovascularization), including, but not limited to, neovascularization associated with tumors.

As used herein, anti-angiogenic treatment or agent refers to any therapeutic regimen and compound, when used alone or in combination with other treatment or compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with undesired and/or uncontrolled angiogenesis. Thus, for purposes herein an anti-angiogenic agent refers to an agent that inhibits the establishment or maintenance of vasculature. Such agents include, but are not limited to, anti-tumor agents, and agents for treatments of other disorders associated with undesirable angiogenesis, such as diabetic retinopathies, restenosis, hyperproliferative disorders and others.

As used herein, non-anti-angiogenic anti-tumor agents refer to anti-tumor agents that do not act primarily by inhibiting angiogenesis.

As used herein, pro-angiogenic agents are agents that promote the establishment or maintenance of the vasculature. Such agents include agents for treating cardiovascular disorders, including heart attacks and strokes.

As used herein, undesired and/or uncontrolled angiogenesis refers to pathological angiogenesis wherein the influence of angiogenesis stimulators outweighs the influence of angiogenesis inhibitors. As used herein, deficient angiogenesis refers to pathological angiogenesis associated with disorders where there is a defect in normal angiogenesis resulting in aberrant angiogenesis or an absence or substantial reduction in angiogenesis.

As used herein, endotheliase refers to a mammalian protein, including humans, that has a transmembrane domain and is expressed on the surface of

-22-

endothelial cells and includes a protease domain, particularly an extracellular protease domain, and is preferably a serine protease. Thus, reference, for example, to endotheliase encompasses all proteins encoded by the endotheliase gene family, or an equivalent molecule obtained from any other source or that
5 has been prepared synthetically or that exhibits the same activity. The endotheliase gene family are transmembrane proteases expressed in endothelial cells. Endotheliases are excluded from the MTSPs contemplated herein.

As used herein, the protease domain of an endotheliase refers to the polypeptide portion of the endotheliase that is the extracellular portion that
10 exhibits protease activity. The protease domain is a polypeptide that includes at least the minimum number of amino acids, generally more than 50 or 100, required for protease activity. Protease activity may be assessed empirically, such as by testing the polypeptide for its ability to act as a protease. Assays, such as the assays described in the EXAMPLES, employing a known substrate in
15 place of the test compounds may be used. Furthermore, since proteases, particularly serine proteases, have characteristic structures and sequences or motifs, the protease domain may be readily identified by such structure and sequence or motif.

As used herein, the protease domain of an MTSP protein refers to the
20 protease domain of an MTSP that is located within or is the extracellular domain of an MTSP and exhibits serine proteolytic activity. Hence it is at least the minimal portion of the extracellular domain that exhibits proteolytic activity as assessed by standard assays *in vitro*. It refers, herein, to a single chain form heretofore thought to be inactive.

25 Exemplary protease domains include at least a sufficient portion of sequences of amino acids set forth as amino acids 615-855 in SEQ ID No. 2 (encoded by nucleotides 1865-2587 in SEQ ID No. 1; see also SEQ ID Nos. 49 and 50) from MTSP1, amino acids 205-437 of SEQ ID NO. 4 from MTSP3, SEQ ID No. 6, which sets forth the protease domain of MTSP4, and amino acids 217-443 of
30 SEQ ID No. 11 from MTSP6. Also contemplated are nucleic acid molecules that encode polypeptide that has proteolytic activity in an *in vitro* proteolysis assay and that have at least 80%, 85%, 90% or 95% sequence identity with the full

RECTIFIED SHEET (RULE 91)

-23-

length of a protease domain of an MTSP protein, or that hybridize along their full length to a nucleic acids that encode a protease domain, particularly under conditions of moderate, generally high, stringency.

For each of these protease domains, residues at the N-terminus can be critical for activity, since it has been shown that an Asp in the N-terminus of such proteases is essential for formation of the catalytically active conformation upon activation cleavage of the zymogen form of the protease. It is shown herein that the protease domain of the single chain form of the protease is catalytically active. Hence the protease domain will require the N-terminal amino acids; the c-terminus portion may be truncated. The amount that can be removed can be determined empirically by testing the protein for protease activity in an *in vitro* assays that assesses catalytic cleavage.

Hence smaller portions of the protease domains, particularly the single chain domains, thereof that retain protease activity are contemplated. Such smaller versions will generally be C-terminal truncated versions of the protease domains. The protease domains vary in size and constitution, including insertions and deletions in surface loops. Such domains exhibit conserved structure, including at least one structural feature, such as the active site triad, primary specificity pocket, oxyanion hole and/or other features of serine protease domains of proteases. Thus, for purposes herein, the protease domain is a single chain portion of an MTSP, as defined herein, but is homologous in its structural features and retention of sequence of similarity or homology the protease domain of chymotrypsin or trypsin. Most significantly, the polypeptide will exhibit proteolytic activity as a single chain.

As used herein, by homologous means about greater than 25% nucleic acid sequence identity, preferably 25% 40%, 60%, 80%, 90% or 95%. The terms "homology" and "identity" are often used interchangeably. In general, sequences are aligned so that the highest order match is obtained (see, *e.g.*: *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994;

-24-

Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073).

By sequence identity, the number of conserved amino acids are
5 determined by standard alignment algorithms programs, and are used with
default gap penalties established by each supplier. Substantially homologous
nucleic acid molecules would hybridize typically at moderate stringency or at
high stringency all along the length of the nucleic acid of interest. Also
contemplated are nucleic acid molecules that contain degenerate codons in place
10 of codons in the hybridizing nucleic acid molecule.

Whether any two nucleic acid molecules have nucleotide sequences that
are at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% "identical" can be
determined using known computer algorithms such as the "FAST A" program,
using for example, the default parameters as in Pearson *et al.* (1988) *Proc. Natl.*
15 *Acad. Sci. USA* 85:2444 (other programs include the GCG program package
(Devereux, J., *et al.*, *Nucleic Acids Research* 12(1):387 (1984)), BLASTP,
BLASTN, FASTA (Atschul, S.F., *et al.*, *J Molec Biol* 215:403 (1990); Guide to
Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and
Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). For example, the BLAST
20 function of the National Center for Biotechnology Information database may be
used to determine identity. Other commercially or publicly available programs
include, DNASTar "MegAlign" program (Madison, WI) and the University of
Wisconsin Genetics Computer Group (UWG) "Gap" program (Madison WI)).
Percent homology or identity of proteins and/or nucleic acid molecules may be
25 determined, for example, by comparing sequence information using a GAP
computer program (*e.g.*, Needleman *et al.* (1970) *J. Mol. Biol.* 48:443, as
revised by Smith and Waterman ((1981) *Adv. Appl. Math.* 2:482). Briefly, the
GAP program defines similarity as the number of aligned symbols (*i.e.*,
nucleotides or amino acids) which are similar, divided by the total number of
30 symbols in the shorter of the two sequences. Default parameters for the GAP
program may include: (1) a unary comparison matrix (containing a value of 1 for
identities and 0 for non-identities) and the weighted comparison matrix of

-25-

Gribskov *et al.* (1986) *Nucl. Acids Res.* 14:6745, as described by Schwartz and Dayhoff, eds., *ATLAS OF PROTEIN SEQUENCE AND STRUCTURE*, National Biomedical Research Foundation, pp. 353-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no
5 penalty for end gaps.

Therefore, as used herein, the term "identity" represents a comparison between a test and a reference polypeptide or polynucleotide. For example, a test polypeptide may be defined as any polypeptide that is 90% or more identical to a reference polypeptide. As used herein, the term at least "90%
10 identical to" refers to percent identities from 90 to 99.99 relative to the reference polypeptides. Identity at a level of 90% or more is indicative of the fact that, assuming for exemplification purposes a test and reference polynucleotide length of 100 amino acids are compared. No more than 10% (i.e., 10 out of 100) amino acids in the test polypeptide differs from that of the
15 reference polypeptides. Similar comparisons may be made between a test and reference polynucleotides. Such differences may be represented as point mutations randomly distributed over the entire length of an amino acid sequence or they may be clustered in one or more locations of varying length up to the maximum allowable, e.g. 10/100 amino acid difference (approximately 90%
20 identity). Differences are defined as nucleic acid or amino acid substitutions, or deletions. At level of homologies or identities above about 85-90%, the result should be independent of the program and gap parameters set; such high levels of identity readily can be assess, often without relying on software.

As used herein, primer refers to an oligonucleotide containing two or
25 more deoxyribonucleotides or ribonucleotides, preferably more than three, from which synthesis of a primer extension product can be initiated. Experimental conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization and extension, such as DNA polymerase, and a suitable buffer, temperature and pH.

30 As used herein, animals include any animal, such as, but are not limited to, goats, cows, deer, sheep, rodents, pigs and humans. Non-human animals, exclude humans as the contemplated animal. The MTSPs provided herein are

-26-

from any source, animal, plant, prokaryotic and fungal. Preferred MTSPs are of animal origin, preferably mammalian origin.

As used herein, genetic therapy involves the transfer of heterologous DNA to the certain cells, target cells, of a mammal, particularly a human, with a
5 disorder or conditions for which such therapy is sought. The DNA is introduced into the selected target cells in a manner such that the heterologous DNA is expressed and a therapeutic product encoded thereby is produced.
Alternatively, the heterologous DNA may in some manner mediate expression of DNA that encodes the therapeutic product, or it may encode a product, such as
10 a peptide or RNA that in some manner mediates, directly or indirectly, expression of a therapeutic product. Genetic therapy may also be used to deliver nucleic acid encoding a gene product that replaces a defective gene or supplements a gene product produced by the mammal or the cell in which it is introduced. The introduced nucleic acid may encode a therapeutic compound, such as a growth
15 factor inhibitor thereof, or a tumor necrosis factor or inhibitor thereof, such as a receptor therefor, that is not normally produced in the mammalian host or that is not produced in therapeutically effective amounts or at a therapeutically useful time. The heterologous DNA encoding the therapeutic product may be modified prior to introduction into the cells of the afflicted host in order to enhance or
20 otherwise alter the product or expression thereof. Genetic therapy may also involve delivery of an inhibitor or repressor or other modulator of gene expression.

As used herein, heterologous DNA is DNA that encodes RNA and proteins that are not normally produced *in vivo* by the cell in which it is expressed or that
25 mediates or encodes mediators that alter expression of endogenous DNA by affecting transcription, translation, or other regulatable biochemical processes. Heterologous DNA may also be referred to as foreign DNA. Any DNA that one of skill in the art would recognize or consider as heterologous or foreign to the cell in which is expressed is herein encompassed by heterologous DNA.
30 Examples of heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins, such as a protein that confers drug resistance, DNA that encodes therapeutically effective substances, such as anti-cancer agents,

-27-

enzymes and hormones, and DNA that encodes other types of proteins, such as antibodies. Antibodies that are encoded by heterologous DNA may be secreted or expressed on the surface of the cell in which the heterologous DNA has been introduced.

5 Hence, herein heterologous DNA or foreign DNA, includes a DNA molecule not present in the exact orientation and position as the counterpart DNA molecule found in the genome. It may also refer to a DNA molecule from another organism or species (*i.e.*, exogenous).

10 As used herein, a therapeutically effective product is a product that is encoded by heterologous nucleic acid, typically DNA, that, upon introduction of the nucleic acid into a host, a product is expressed that ameliorates or eliminates the symptoms, manifestations of an inherited or acquired disease or that cures the disease.

15 As used herein, recitation that a polypeptide consists essentially of the protease domain means that the only MTSP portion of the polypeptide is a protease domain or a catalytically active portion thereof. The polypeptide may optionally, and generally will, include additional non-MTSP-derived sequences of amino acids.

20 As used herein, cancer or tumor treatment or agent refers to any therapeutic regimen and/or compound that, when used alone or in combination with other treatments or compounds, can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with deficient angiogenesis.

25 As used herein, domain refers to a portion of a molecule, *e.g.*, proteins or nucleic acids, that is structurally and/or functionally distinct from other portions of the molecule.

30 As used herein, protease refers to an enzyme catalyzing hydrolysis of proteins or peptides. For purposes herein, the protease domain is a single chain form of an MTSP protein. For MTSP3 and MTSP4 the protease domain also includes two chain forms.

-28-

As used herein, catalytic activity refers to the activity of the MTSP as a protease as assessed in *in vitro* proteolytic assays that detect proteolysis of a selected substrate.

As used herein, nucleic acids include DNA, RNA and analogs thereof,
5 including protein nucleic acids (PNA) and mixture thereof. Nucleic acids can be single or double stranded. When referring to probes or primers, optionally labeled, with a detectable label, such as a fluorescent or radiolabel, single-stranded molecules are contemplated. Such molecules are typically of a length such that they are statistically unique and of a low copy number (typically less
10 than 5, preferably less than 3) for probing or priming a library. Generally a probe or primer contains at least 14, 16 or 30 contiguous sequence complementary to or identical to a gene of interest. Probes and primers can be 10, 20, 30, 50, 100 or more nucleic acids long.

As used herein, nucleic acid encoding a fragment or portion of an MTSP
15 refers to a nucleic acid encoding only the recited fragment or portion of MTSP, and not the other contiguous portions of the MTSP.

As used herein, heterologous or foreign DNA and RNA are used interchangeably and refer to DNA or RNA that does not occur naturally as part of the genome in which it is present or which is found in a location or locations in
20 the genome that differ from that in which it occurs in nature. Heterologous nucleic acid is generally not endogenous to the cell into which it is introduced, but has been obtained from another cell or prepared synthetically. Generally, although not necessarily, such nucleic acid encodes RNA and proteins that are not normally produced by the cell in which it is expressed. Any DNA or RNA
25 that one of skill in the art would recognize or consider as heterologous or foreign to the cell in which it is expressed is herein encompassed by heterologous DNA. Heterologous DNA and RNA may also encode RNA or proteins that mediate or alter expression of endogenous DNA by affecting transcription, translation, or other regulatable biochemical processes.

30 As used herein, operative linkage of heterologous DNA to regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences refers to the relationship

RECTIFIED SHEET (RULE 91)

-29-

between such DNA and such sequences of nucleotides. For example, operative linkage of heterologous DNA to a promoter refers to the physical relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes,
5 binds to and transcribes the DNA in reading frame.

As used herein, a sequence complementary to at least a portion of an RNA, with reference to antisense oligonucleotides, means a sequence having sufficient complementarity to be able to hybridize with the RNA, preferably under moderate or high stringency conditions, forming a stable duplex; in the case of
10 double-stranded MTSP antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize depends on the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with a MTSP encoding RNA it can contain and still form
15 a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

For purposes herein, conservative amino acid substitutions may be made in any of MTSPs and protease domains thereof provided that the resulting
20 protein exhibits protease activity. Conservative amino acid substitutions, such as those set forth in Table 1, are those that do not eliminate proteolytic activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single
25 amino acid substitutions in non-essential regions of a polypeptide do not substantially alter biological activity (see, e.g., Watson et al. *Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224). Also included within the definition, is the catalytically active fragment of an MTSP, particularly a single chain protease portion. Conservative amino acid
30 substitutions are made, for example, in accordance with those set forth in TABLE 1 as follows:

-30-

TABLE 1

	Original residue	Conservative substitution
	Ala (A)	Gly; Ser, Abu
	Arg (R)	Lys, orn
5	Asn (N)	Gln; His
	Cys (C)	Ser
	Gln (Q)	Asn
	Glu (E)	Asp
	Gly (G)	Ala; Pro
10	His (H)	Asn; Gln
	Ile (I)	Leu; Val; Met; Nle; Nva
	Leu (L)	Ile; Val; Met; Nle; Nv
	Lys (K)	Arg; Gln; Glu
	Met (M)	Leu; Tyr; Ile; NLe Val
15	Ornithine	Lys; Arg
	Phe (F)	Met; Leu; Tyr
	Ser (S)	Thr
	Thr (T)	Ser
	Trp (W)	Tyr
20	Tyr (Y)	Trp; Phe
	Val (V)	Ile; Leu; Met; Nle; Nv

Other substitutions are also permissible and may be determined empirically or in accord with known conservative substitutions.

As used herein, Abu is 2-aminobutyric acid; Orn is ornithine.

25 As used herein, the amino acids, which occur in the various amino acid sequences appearing herein, are identified according to their well-known, three-letter or one-letter abbreviations. The nucleotides, which occur in the various DNA fragments, are designated with the standard single-letter designations used routinely in the art.

30 As used herein, a splice variant refers to a variant produced by differential processing of a primary transcript of genomic DNA that results in more than one type of mRNA.

As used herein, a probe or primer based on a nucleotide sequence disclosed herein, includes at least 10, 14, preferably at least 16 or 30 or 100
35 contiguous sequence of nucleotides of SEQ ID Nos. 1, 3, 5, 7, 9 or 11.

As used herein, amelioration of the symptoms of a particular disorder by administration of a particular pharmaceutical composition refers to any lessening,

-31-

whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

As used herein, antisense polynucleotides refer to synthetic sequences of nucleotide bases complementary to mRNA or the sense strand of double stranded DNA. Admixture of sense and antisense polynucleotides under appropriate conditions leads to the binding of the two molecules, or hybridization. When these polynucleotides bind to (hybridize with) mRNA, inhibition of protein synthesis (translation) occurs. When these polynucleotides bind to double stranded DNA, inhibition of RNA synthesis (transcription) occurs. The resulting inhibition of translation and/or transcription leads to an inhibition of the synthesis of the protein encoded by the sense strand. Antisense nucleic acid molecule typically contain a sufficient number of nucleotides to specifically bind to a target nucleic acid, generally at least 5 contiguous nucleotides, often at least 14 or 16 or 30 contiguous nucleotides or modified nucleotides complementary to the coding portion of a nucleic acid molecule that encodes a gene of interest, for example, nucleic acid encoding a single chain protease domain of an MTSP.

As used herein, an array refers to a collection of elements, such as antibodies, containing three or more members. An addressable array is one in which the members of the array are identifiable, typically by position on a solid phase support. Hence, in general the members of the array will be immobilized to discrete identifiable loci on the surface of a solid phase.

As used herein, antibody refers to an immunoglobulin, whether natural or partially or wholly synthetically produced, including any derivative thereof that retains the specific binding ability the antibody. Hence antibody includes any protein having a binding domain that is homologous or substantially homologous to an immunoglobulin binding domain. Antibodies include members of any immunoglobulin claims, including IgG, IgM, IgA, IgD and IgE.

As used herein, antibody fragment refers to any derivative of an antibody that is less than full length, retaining at least a portion of the full-length antibody's specific binding ability. Examples of antibody fragments include, but are not limited to, Fab, Fab', F(ab)₂, single-chain Fvs (scFV), FV, dsFV diabody

-32-

and Fd fragments. The fragment can include multiple chains linked together, such as by disulfide bridges. An antibody fragment generally contains at least about 50 amino acids and typically at least 200 amino acids.

As used herein, an Fv antibody fragment is composed of one variable
5 heavy domain (V_H) and one variable light domain linked by noncovalent interactions.

As used herein, a dsFV refers to an Fv with an engineered intermolecular disulfide bond, which stabilizes the V_H - V_L pair.

As used herein, an $F(ab)_2$ fragment is an antibody fragment that results
10 from digestion of an immunoglobulin with pepsin at pH 4.0-4.5; it may be recombinantly produced.

As used herein, Fab fragments is an antibody fragment that results from digestion of an immunoglobulin with papain; it may be recombinantly produced.

As used herein, scFVs refer to antibody fragments that contain a variable
15 light chain (V_L) and variable heavy chain (V_H) covalently connected by a polypeptide linker in any order. The linker is of a length such that the two variable domains are bridged without substantial interference. Preferred linkers are $(Gly-Ser)_n$ residues with some Glu or Lys residues dispersed throughout to increase solubility.

20 As used herein, humanized antibodies refer to antibodies that are modified to include human sequences of amino acids so that administration to a human will not provoke an immune response. Methods for preparation of such antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an
25 antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify such regions.

As used herein, diabodies are dimeric scFV; diabodies typically have shorter peptide linkers than scFVs, and they preferentially dimerize.

30 As used herein, humanized antibodies refer to antibodies that are modified to include human sequences of amino acids so that administration to a human will not provoke an immune response. Methods for preparation of such

-33-

antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify
5 such regions.

As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA.

As used herein the term assessing is intended to include
10 quantitative and qualitative determination in the sense of obtaining an absolute value for the activity of an MTSP, or a domain thereof, present in the sample, and also of obtaining an index, ratio, percentage, visual or other value indicative of the level of the activity. Assessment may be direct or indirect and the chemical species actually detected need not of course be the proteolysis
15 product itself but may for example be a derivative thereof or some further substance.

As used herein, biological activity refers to the in vivo activities of a compound or physiological responses that result upon in vivo administration of a compound, composition or other mixture. Biological activity, thus, encompasses
20 therapeutic effects and pharmaceutical activity of such compounds, compositions and mixtures. Biological activities may be observed in in vitro systems designed to test or use such activities. Thus, for purposes herein the biological activity of a luciferase is its oxygenase activity whereby, upon oxidation of a substrate, light is produced.

25 As used herein, a combination refers to any association between two or among more items.

As used herein, a composition refers to any mixture. It may be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

-34-

As used herein, a conjugate refers to the compounds provided herein that include one or more MTSPs, particularly single chain protease domains thereof, and one or more targeting agents. These conjugates include those produced by recombinant means as fusion proteins, those produced by chemical means, such as by chemical coupling, through, for example, coupling to sulfhydryl groups, and those produced by any other method whereby at least one MTSP, or a domain thereof, is linked, directly or indirectly via linker(s) to a targeting agent.

As used herein, a targeting agent, is any moiety, such as a protein or effective portion thereof, that provides specific binding of the conjugate to a cell surface receptor, which, preferably, internalizes the conjugate or MTSP portion thereof. A targeting agent may also be one that promotes or facilitates, for example, affinity isolation or purification of the conjugate; attachment of the conjugate to a surface; or detection of the conjugate or complexes containing the conjugate.

As used herein, an antibody conjugate refers to a conjugate in which the targeting agent is an antibody.

As used herein, humanized antibodies refer to antibodies that are modified to include human sequences of amino acids so that administration to a human will not provoke an immune response. Methods for preparation of such antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify such regions.

As used herein, derivative or analog of a molecule refers to a portion derived from or a modified version of the molecule.

As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

As used herein, an effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such amount may

-35-

be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Repeated administration may be required to achieve the desired amelioration of symptoms.

As used herein equivalent, when referring to two sequences of nucleic acids means that the two sequences in question encode the same sequence of amino acids or equivalent proteins. When equivalent is used in referring to two proteins or peptides, it means that the two proteins or peptides have substantially the same amino acid sequence with only conservative amino acid substitutions (see, *e.g.*, Table 1, above) that do not substantially alter the activity or function of the protein or peptide. When equivalent refers to a property, the property does not need to be present to the same extent [*e.g.*, two peptides can exhibit different rates of the same type of enzymatic activity], but the activities are preferably substantially the same. Complementary, when referring to two nucleotide sequences, means that the two sequences of nucleotides are capable of hybridizing, preferably with less than 25%, more preferably with less than 15%, even more preferably with less than 5%, most preferably with no mismatches between opposed nucleotides. Preferably the two molecules will hybridize under conditions of high stringency.

As used herein, an agent that modulates the activity of a protein or expression of a gene or nucleic acid either decreases or increases or otherwise alters the activity of the protein or, in some manner up- or down-regulates or otherwise alters expression of the nucleic acid in a cell.

As used herein, inhibitor of an the activity of an MTSP encompasses any substances that prohibit or decrease production, post-translational modification(s), maturation, or membrane localization of the MTSP or any substances that interfere with or decrease the proteolytic efficacy of thereof, particular of a single chain form *in vitro*.

As used herein, a method for treating or preventing neoplastic disease means that any of the symptoms, such as the tumor, metastasis thereof, the vascularization of the tumors or other parameters by which the disease is

-36-

characterized are reduced, ameliorated, prevented, placed in a state of remission, or maintained in a state of remission. It also means that the hallmarks of neoplastic disease and metastasis may be eliminated, reduced or prevented by the treatment. Non-limiting examples of the hallmarks include uncontrolled

- 5 degradation of the basement membrane and proximal extracellular matrix, migration, division, and organization of the endothelial cells into new functioning capillaries, and the persistence of such functioning capillaries.

- As used herein, operatively linked or operationally associated refers to the functional relationship of DNA with regulatory and effector sequences of
- 10 nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences. For example, operative linkage of DNA to a promoter refers to the physical and functional relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and
- 15 transcribes the DNA. In order to optimize expression and/or *in vitro* transcription, it may be necessary to remove, add or alter 5' untranslated portions of the clones to eliminate extra, potential inappropriate alternative translation initiation (*i.e.*, start) codons or other sequences that may interfere with or reduce expression, either at the level of transcription or translation.
- 20 Alternatively, consensus ribosome binding sites (see, e.g., Kozak *J. Biol. Chem.* 266:19867-19870 (1991)) can be inserted immediately 5' of the start codon and may enhance expression. The desirability of (or need for) such modification may be empirically determined.

- As used herein, pharmaceutically acceptable salts, esters or other
- 25 derivatives of the conjugates include any salts, esters or derivatives that may be readily prepared by those of skill in this art using known methods for such derivatization and that produce compounds that may be administered to animals or humans without substantial toxic effects and that either are pharmaceutically active or are prodrugs.

- 30 As used herein, a prodrug is a compound that, upon in vivo administration, is metabolized or otherwise converted to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a

-37-

prodrug, the pharmaceutically active compound is modified such that the active compound will be regenerated by metabolic processes. The prodrug may be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes and drug metabolism in vivo, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, e.g., Nogrady (1985) Medicinal Chemistry A Biochemical Approach, Oxford University Press, New York, pages 388-392).

- 10 As used herein, a drug identified by the screening methods provided herein refers to any compound that is a candidate for use as a therapeutic or as lead compound for designed a therapeutic. Such compounds can be small molecules, including small organic molecules, peptides, peptide mimetics, antisense molecules, antibodies, fragments of antibodies, recombinant antibodies and other such compound which can serve as drug candidate or lead compound.

15 As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA.

- 20 As used herein, a promoter region or promoter element refers to a segment of DNA or RNA that controls transcription of the DNA or RNA to which it is operatively linked. The promoter region includes specific sequences that are sufficient for RNA polymerase recognition, binding and transcription initiation. This portion of the promoter region is referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding and transcription initiation activity of RNA polymerase. These sequences may be *cis* acting or may be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, may be constitutive or regulated. Exemplary promoters contemplated for use in prokaryotes include the bacteriophage T7 and T3 promoters.

- 30 As used herein, a receptor refers to a molecule that has an affinity for a given ligand. Receptors may be naturally-occurring or synthetic molecules. Receptors may also be referred to in the art as anti-ligands. As used herein, the

-38-

- receptor and anti-ligand are interchangeable. Receptors can be used in their unaltered state or as aggregates with other species. Receptors may be attached, covalently or noncovalently, or in physical contact with, to a binding member, either directly or indirectly via a specific binding substance or linker. Examples
- 5 of receptors, include, but are not limited to: antibodies, cell membrane receptors surface receptors and internalizing receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants [such as on viruses, cells, or other materials], drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins, sugars, polysaccharides, cells, cellular membranes, and organelles.
- 10 Examples of receptors and applications using such receptors, include but are not restricted to:
- a) enzymes: specific transport proteins or enzymes essential to survival of microorganisms, which could serve as targets for antibiotic [ligand] selection;
 - b) antibodies: identification of a ligand-binding site on the antibody

15 molecule that combines with the epitope of an antigen of interest may be investigated; determination of a sequence that mimics an antigenic epitope may lead to the development of vaccines of which the immunogen is based on one or more of such sequences or lead to the development of related diagnostic agents or compounds useful in therapeutic treatments such as for auto-immune diseases

 - 20 c) nucleic acids: identification of ligand, such as protein or RNA, binding sites;
 - d) catalytic polypeptides: polymers, preferably polypeptides, that are capable of promoting a chemical reaction involving the conversion of one or more reactants to one or more products; such polypeptides generally include a

25 binding site specific for at least one reactant or reaction intermediate and an active functionality proximate to the binding site, in which the functionality is capable of chemically modifying the bound reactant [see, e.g., U.S. Patent No. 5,215,899];

 - e) hormone receptors: determination of the ligands that bind with high

30 affinity to a receptor is useful in the development of hormone replacement therapies; for example, identification of ligands that bind to such receptors may lead to the development of drugs to control blood pressure; and

-39-

f) opiate receptors: determination of ligands that bind to the opiate receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

As used herein, sample refers to anything which may contain an analyte
5 for which an analyte assay is desired. The sample may be a biological sample, such as a biological fluid or a biological tissue. Examples of biological fluids include urine, blood, plasma, serum, saliva, semen, stool, sputum, cerebral spinal fluid, tears, mucus, amniotic fluid or the like. Biological tissues are aggregate of cells, usually of a particular kind together with their intercellular substance that
10 form one of the structural materials of a human, animal, plant, bacterial, fungal or viral structure, including connective, epithelium, muscle and nerve tissue. Examples of biological tissues also include organs, tumors, lymph nodes, arteries and individual cell(s).

As used herein: stringency of hybridization in determining percentage
15 mismatch is as follows:

- 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C
- 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C
- 3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C

Those of skill in this art know that the washing step selects for stable
20 hybrids and also know the ingredients of SSPE (see, e.g., Sambrook, E.F. Fritsch, T. Maniatis, in: Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory Press (1989), vol. 3, p. B.13, see, also, numerous catalogs that describe commonly used laboratory solutions). SSPE is pH 7.4 phosphate-buffered 0.18 NaCl. Further, those of skill in the art recognize that the stability
25 of hybrids is determined by T_m , which is a function of the sodium ion concentration and temperature ($T_m = 81.5^\circ \text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G} + \text{C}) - 600/l$), so that the only parameters in the wash conditions critical to hybrid stability are sodium ion concentration in the SSPE (or SSC) and temperature.

It is understood that equivalent stringencies may be achieved using
30 alternative buffers, salts and temperatures. By way of example and not limitation, procedures using conditions of low stringency are as follows (see also Shilo and Weinberg, *Proc. Natl. Acad. Sci. USA*, 78:6789-6792 (1981)): Filters

RECTIFIED SHEET (RULE 91)

-40-

containing DNA are pretreated for 6 hours at 40°C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 µg/ml denatured salmon sperm DNA (10X SSC is 1.5 M sodium chloride, and 0.15 M sodium citrate, adjusted to a pH of 7).

- 5 Hybridizations are carried out in the same solution with the following modifications: 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 µg/ml salmon sperm DNA, 10% (wt/vol) dextran sulfate, and 5-20 X 10⁸ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 40°C, and then washed for 1.5 hours at 55°C in a solution containing 2X SSC, 25 mM
- 10 Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS. The wash solution is replaced with fresh solution and incubated an additional 1.5 hours at 60°C. Filters are blotted dry and exposed for autoradiography. If necessary, filters are washed for a third time at 65-68°C and reexposed to film. Other conditions of low stringency which may be used are well known in the art (*e.g.*, as employed for
- 15 cross-species hybridizations).

- By way of example and not way of limitation, procedures using conditions of moderate stringency is provided. For example, but not limited to, procedures using such conditions of moderate stringency are as follows: Filters containing DNA are pretreated for 6 hours at 55°C in a solution containing 6X
- 20 SSC, 5X Denhart's solution, 0.5% SDS and 100 µg/ml denatured salmon sperm DNA. Hybridizations are carried out in the same solution and 5-20 X 10⁸ cpm ³²P-labeled probe is used. Filters are incubated in hybridization mixture for 18-20 hours at 55°C, and then washed twice for 30 minutes at 60°C in a solution containing 1X SSC and 0.1% SDS. Filters are blotted dry and exposed for
- 25 autoradiography. Other conditions of moderate stringency which may be used are well-known in the art. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.1% SDS.

- By way of example and not way of limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is
- 30 carried out for 8 hours to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 µg/ml denatured salmon sperm DNA. Filters are hybridized for 48 hours

-41-

at 65°C in prehybridization mixture containing 100 µg/ml denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Washing of filters is done at 37°C for 1 hour in a solution containing 2X SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA. This is followed by a wash in 0.1X SSC at 50°C for 45

5 minutes before autoradiography. Other conditions of high stringency which may be used are well known in the art.

The term substantially identical or homologous or similar varies with the context as understood by those skilled in the relevant art and generally means at least 70%, preferably means at least 80%, more preferably at least 90%, and
10 most preferably at least 95% identity.

As used herein, substantially identical to a product means sufficiently similar so that the property of interest is sufficiently unchanged so that the substantially identical product can be used in place of the product.

As used herein, substantially pure means sufficiently homogeneous to
15 appear free of readily detectable impurities as determined by standard methods of analysis, such as thin layer chromatography (TLC), gel electrophoresis and high performance liquid chromatography (HPLC), used by those of skill in the art to assess such purity, or sufficiently pure such that further purification would not detectably alter the physical and chemical properties, such as enzymatic and
20 biological activities, of the substance. Methods for purification of the compounds to produce substantially chemically pure compounds are known to those of skill in the art. A substantially chemically pure compound may, however, be a mixture of stereoisomers or isomers. In such instances, further purification might increase the specific activity of the compound.

25 As used herein, target cell refers to a cell that expresses an MTSP *in vivo*.

As used herein, test substance refers to a chemically defined compound (*e.g.*, organic molecules, inorganic molecules, organic/inorganic molecules, proteins, peptides, nucleic acids, oligonucleotides, lipids, polysaccharides, saccharides, or hybrids among these molecules such as glycoproteins, etc.) or
30 mixtures of compounds (*e.g.*, a library of test compounds, natural extracts or culture supernatants, etc.) whose effect on an MTSP, particularly a single chain

-42-

form that includes the protease domain or a sufficient portion thereof for activity, as determined by *in vitro* method, such as the assays provided herein.

As used herein, the terms a therapeutic agent, therapeutic regimen, radioprotectant, chemotherapeutic mean conventional drugs and drug therapies, including vaccines, which are known to those skilled in the art. Radiotherapeutic agents are well known in the art.

As used herein, treatment means any manner in which the symptoms of a conditions, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

10 As used herein, vector (or plasmid) refers to discrete elements that are used to introduce heterologous DNA into cells for either expression or replication thereof. The vectors typically remain episomal, but may be designed to effect integration of a gene or portion thereof into a chromosome of the genome. Also contemplated are vectors that are artificial chromosomes, such as yeast artificial
15 chromosomes and mammalian artificial chromosomes. Selection and use of such vehicles are well known to those of skill in the art. An expression vector includes vectors capable of expressing DNA that is operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting expression of such DNA fragments. Thus, an expression vector refers to a
20 recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well known to those of skill in the art and include those that are replicable in eukaryotic cells and/or prokaryotic cells and those that remain episomal or those
25 which integrate into the host cell genome.

As used herein, protein binding sequence refers to a protein or peptide sequence that is capable of specific binding to other protein or peptide sequences generally, to a set of protein or peptide sequences or to a particular protein or peptide sequence.

30 As used herein, epitope tag refers to a short stretch of amino acid residues corresponding to an epitope to facilitate subsequent biochemical and immunological analysis of the epitope tagged protein or peptide. Epitope tagging

-43-

is achieved by appending the sequence of the epitope tag to the protein-encoding sequence in an appropriate expression vector. Epitope tagged proteins can be affinity purified using highly specific antibodies raised against the tags.

As used herein, metal binding sequence refers to a protein or peptide
5 sequence that is capable of specific binding to metal ions generally, to a set of metal ions or to a particular metal ion.

As used herein, a composition refers to a any mixture. It may be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

10 As used herein, a combination refers to any association between two or among more items.

As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

15 As used herein, a cellular extract refers to a preparation or fraction which is made from a lysed or disrupted cell.

As used herein, an agent is said to be randomly selected when the agent is chosen randomly without considering the specific sequences involved in the association of a protein alone or with its associated substrates, binding partners,
20 etc. An example of randomly selected agents is the use a chemical library or a peptide combinatorial library, or a growth broth of an organism.

As used herein, an agent is the to be rationally selected or designed when the agent is chosen on a non-random basis which takes into account the sequence of the target site and/or its conformation in connection with the
25 agent's action. As described in the Examples, there are proposed binding sites for serine protease and (catalytic) sites in the protein having SEQ ID NO:3 or SEQ ID NO:4. Agents can be rationally selected or rationally designed by utilizing the peptide sequences that make up these sites. For example, a rationally selected peptide agent can be a peptide whose amino acid sequence is
30 identical to the ATP or calmodulin binding sites or domains.

For clarity of disclosure, and not by way of limitation, the detailed description is divided into the subsections that follow.

-44-

**B. MTSP PROTEINS, MUTEINS, DERIVATIVES AND ANALOGS THEREOF
MTSPs**

The MTSPs are a family of transmembrane serine proteases that are found in mammals and also other species that share a number of common structural features including: a proteolytic extracellular C-terminal domain; a transmembrane domain, with a hydrophobic domain near the N-terminus; a short cytoplasmic domain; and a variable length stem region containing modular domains. The proteolytic domains share sequence homology including conserved his, asp, and ser residues necessary for catalytic activity that are present in conserved motifs. The MTSPs are synthesized as zymogens, and activated to double chain forms by cleavage. It is shown herein that the single chain proteolytic domain can function *in vitro* and, hence is useful in *in vitro* assays for identifying agents that modulate the activity of members of this family. Also provided are family members designated MTSP3, MTSP4 and an MTSP6 variant.

The MTSP family is a target for therapeutic intervention and also some, may serve as diagnostic markers for tumor development, growth and/or progression. As discussed, the members of this family are involved in proteolytic processes that are implicated in tumor development, growth and/or progression. This implication is based upon their functions as proteolytic enzymes in processes related to ECM degradative pathways. In addition, their levels of expression or level of activation or their apparent activity resulting from substrate levels or alterations in substrates and levels thereof differs in tumor cells and non-tumor cells in the same tissue. Hence, protocols and treatments that alter their activity, such as their proteolytic activities and roles in signal transduction, and/or their expression, such as by contacting them with a compound that modulates their activity and/or expression, could impact tumor development, growth and/or progression. Also, in some instances, the level of activation and/or expression may be altered in tumors, such as lung carcinoma, colon adenocarcinoma and ovarian carcinoma.

The MTSP may serve as a diagnostic marker for tumors. It is shown herein, that MTSP3 and MTSP4 and the MTSP6 variant provided herein are

RECTIFIED SHEET (RULE 91)

-45-

expressed and/or activated in certain tumors; hence their activation or expression may serve as a diagnostic marker for tumor development, growth and/or progression. In other instances the MTSP protein can exhibit altered activity by virtue of a change in activity or expression of a co-factor therefor or a substrate therefor. In addition, in some instances, these MTSPs and/or variants thereof may be shed from cell surfaces. Detection of the shed MTSPs, particularly the extracellular domains, in body fluids, such as serum, blood, saliva, cerebral spinal fluid, synovial fluid and interstitial fluids, urine, sweat and other such fluids and secretions, may serve as a diagnostic tumor marker. In particular, detection of higher levels of such shed polypeptides in a subject compared to a subject known not to have any neoplastic disease or compared to earlier samples from the same subject, can be indicative of neoplastic disease in the subject.

Provided herein are isolated substantially pure single polypeptides that contain the protease domain of an MTSP as a single chain. The MTSPs contemplated herein are not expressed on endothelial cells, and, preferably, are expressed on tumor cells, typically at a level that differs from the level in which they are expressed in the non-tumor cell of the same type. Hence, for example, if the MTSP is expressed in an ovarian tumor cell, to be of interest herein with respect to ovarian cancer, it is expressed at the same level in non-tumor ovarian cells. MTSP protease domains include the single chain protease domains of MTSP1, MTSP3, MTSP4, MTSP6 and other such proteases, including, but are not limited to, corin, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4.

Provided are the protease domains or proteins that include a portion of an MTSP that is the protease domain of any MTSP, particularly an MTSP1, MTSP3, MTSP4 and MTSP6. The protein can also include other non-MTSP sequences of amino acids, but will include the protease domain or a sufficient portion thereof to exhibit catalytic activity in any *in vitro* assay that assess such protease activity, such as any provided herein.

Also provided herein are nucleic acid molecules that encode MTSP proteins and the encoded proteins. In particular, nucleic acid molecules

RECTIFIED SHEET (RULE 91)

-46-

encoding MTSP-3 and MTSP-4 from animals, including splice variants thereof are provided. The encoded proteins are also provided. Also provided are functional domains thereof.

In specific aspects, the MTSP protease domains, portions thereof, and
5 muteins thereof are from or based on animal MTSPs, including, but are not limited to, rodent, such as mouse and rat; fowl, such as chicken; ruminants, such as goats, cows, deer, sheep; ovine, such as pigs; and humans.

In particular, MTSP derivatives can be made by altering their sequences by substitutions, additions or deletions that provide for functionally equivalent
10 molecules. Due to the degeneracy of nucleotide coding sequences, other nucleic sequences which encode substantially the same amino acid sequence as a *MTSP* gene can be used. These include but are not limited to nucleotide sequences comprising all or portions of *MTSP* genes that are altered by the substitution of different codons that encode the amino acid residue within the sequence, thus
15 producing a silent change. Likewise, the MTSP derivatives include, but are not limited to, those containing, as a primary amino acid sequence, all or part of the amino acid sequence of MTSP, including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a silent change. For example, one or more amino acid residues
20 within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine,
25 proline, phenylalanine, tryptophan and methionine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid (see, *e.g.*, Table 1).

30 In a preferred embodiment, the substantially purified MTSP protease is encoded by a nucleic acid that hybridizes to the a nucleic acid molecule containing the protease domain encoded by the nucleotide sequence set forth in

-47-

any of SEQ. ID Nos. 1, 3, 5, 7, 9 or 11 under at least moderate, generally high, stringency conditions, such that the protease domain encoding nucleic acid thereof hybridizes along its full length. In preferred embodiments the substantially purified MTSP protease is a single chain polypeptide that includes

5 substantially the sequence of amino acids set forth in any SEQ ID Nos. 2, 4, 6, 8, 10 and 12 that encodes the protease domain. Specific sequences for the following human MTSPs and domains thereof are provided as follows: SEQ ID No. 3 MTSP3 nucleic acid sequence; SEQ ID No. 4 MTSP3 amino acid sequence; SEQ ID No. 5 MTSP4 nucleic acid encoding the protease

10 domain; SEQ ID No. 6 MTSP4 amino acid sequence of the protease domain; SEQ ID No. 7 MTSP4-L nucleic acid sequence; SEQ ID No. 8 MTSP4-L amino acid sequence; SEQ ID No. 9 MTSP4-S nucleic acid sequence; SEQ ID No. 10 MTSP4-S amino acid sequence; SEQ ID No. 11 MTSP6 nucleic acid sequence; SEQ ID No. 12 MTSP6 amino acid sequence. SEQ ID No. 1 sets forth the nucleic

15 acid sequence of the long form of MTSP1; SEQ ID No. 2 the encoded amino acid sequence; SEQ ID No. 49 sets forth the sequence of a protease domain of an MTSP1, and SEQ ID No. 50 the sequence of the encoded single chain protease domain thereof. Figures 1-3 depict the structural organization of the MTSP3, MTSP4 and MTSP6, respectively.

20 In particular, exemplary protease domains include at least a sufficient portion of sequences of amino acids set forth as amino acids 615-855 in SEQ ID No. 2 (encoded by nucleotides 1865-2587 in SEQ ID No. 1; see also SEQ ID Nos. 49 and 50) from MTSP1 (matriptase), amino acids 205-437 of SEQ ID NO. 4 from MTSP3, SEQ ID No. 6, which sets forth the protease domain of MTSP4, and amino acids 217-443 of SEQ ID No. 11 from MTSP6. Also

25 contemplated are nucleic acid molecules that encode a single chain MTSP protease that have proteolytic activity in an *in vitro* proteolysis assay and that have at least 60%, 70%, 80%, 85%, 90% or 95% sequence identity with the full length of a protease domain of an MTSP protein, or that hybridize along their

30 full length to a nucleic acids that encode a protease domain, particularly under conditions of moderate, generally high, stringency. As above, the encoded polypeptides contain the protease as a single chain.

-48-

The isolated nucleic acids may include of at least 8 nucleotides of an *MTSP* sequence. In other embodiments, the nucleic acids may contain least 25 (continuous) nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, or 200 nucleotides of a *MTSP* sequence, or a full-length *MTSP* coding sequence. In
5 another embodiment, the nucleic acids are smaller than 35, 200 or 500 nucleotides in length. Nucleic acids can be single or double stranded. Nucleic acids that hybridizes to or complementary to the foregoing sequences, in particular the inverse complement to nucleic acids that hybridizes to the foregoing sequences (*i.e.*, the inverse complement of a nucleic acid strand has
10 the complementary sequence running in reverse orientation to the strand so that the inverse complement would hybridize without mismatches to the nucleic acid strand; thus, for example, where the coding strand is that hybridizes to a nucleic acid with no mismatches between the coding strand and the that hybridizes strand, then the inverse complement of the that hybridizes strand is identical to
15 the coding strand) are also provided. In specific aspects, nucleic acids are provided that include a sequence complementary to (specifically are the inverse complement of) at least 10, 25, 50, 100, or 200 nucleotides or the entire coding region of an *MTSP* encoding nucleic acid, particularly the protease domain thereof. For *MTSP3* and *MTSP4* the full-length protein or domain or active
20 fragment thereof.

For each of the nucleic acid molecules, the nucleic acid can be DNA or RNA or PNA or other nucleic acid analogs or may include non-natural nucleotide bases.

Also provided are isolated nucleic acid molecules that include a sequence
25 of nucleotides complementary to the nucleotide sequence encoding an *MTSP*.

Probes and primers derived from the nucleic acid molecules are provided, Such probes and primers contain at least 8, 14, 16, 30, 100 or more contiguous nucleotides with identity to contiguous nucleotides of an *MTSP*, including, but are not limited to, *MTSP1*, *MTSP3*, *MTSP4* and *MTSP6*. The probes and primers
30 are optionally labelled with a detectable label, such as a radiolabel or a fluorescent tag, or can be mass differentiated for detection by mass spectrometry or other means.

Plasmids and vectors containing the nucleic acid molecules are also provided. Cells containing the vectors, including cells that express the encoded proteins are provided. The cell can be a bacterial cell, a yeast cell, a fungal cell, a plant cell, an insect cell or an animal cell. Methods for producing an MTSP or
5 single chain form of the protease domain thereof by, for example, growing the cell under conditions whereby the encoded MTSP is expressed by the cell, and recovering the expressed protein, are provided herein. As noted, for MTSP3 and MTSP4, the full-length zymogens and activated proteins and activated (two strand) protease and single chain protease domains are provided.

10 Except for the MTSP proteins (MTSP3 and MTSP4) heretofore unidentified and provided herein, the isolated polypeptides contain the MTSP protease domain or a catalytically active portion thereof and, generally, do not contain additional MTSP. Hence isolated, substantially pure proteases, protease domains or catalytically active portion thereof in single chain form of MTSPs are
15 provided. The protease domains may be included in a longer protein, but such longer protein is not the MTSP zymogen.

Thus, MTSP3 and MTSP4 proteins are provided. For these proteins, the domains, fragments, derivatives or analogs that are functionally active, *i.e.*, capable of exhibiting one or more functional activities associated with the MTSP
20 protein, *e.g.*, serine protease activity, immunogenicity and antigenicity, are provided. As discussed above, the protease domains thereof are also provided. For MTSP3 and MTSP4, the zymogens and activated forms, and also, the single chain and double chain, activated protease domains are provided.

Also provided are nucleic acid molecules that hybridize to the above-
25 noted sequences of nucleotides encoding MTSP3 and MTSP4 (SEQ ID Nos. 3, 5, 7 and 9) at least at low stringency, more preferably at moderate stringency, and most preferably at high stringency, and that encode the protease domain and/or the full length protein or other domains of an MTSP family member, such as MTSP3, MTSP4, MTSP6 or a splice variant or allelic variant thereof, or MTSP6
30 or a splice variant or allelic variant thereof. Preferably the molecules hybridize under such conditions along their full length for at least one domain and encode at least one domain, such as the protease or extracellular domain, of the

-50-

polypeptide. In particular, such nucleic acid molecules include any isolated nucleic fragment that encodes at least one domain of a membrane serine protease, that (1) contains a sequence of nucleotides that encodes the protease or a domain thereof, and (2) is selected from among:

- 5 (a) a sequence of nucleotides that encodes the protease or a domain thereof includes a sequence of nucleotides set forth above;
- (b) a sequence of nucleotides that encodes such portion or the full length protease and hybridizes under conditions of high stringency, preferably to nucleic acid that is complementary to a mRNA
10 transcript present in a mammalian cell that encodes such protein or fragment thereof;
- (c) a sequence of nucleotides that encodes a transmembrane protease or domain thereof that includes a sequence of amino acids encoded by such portion or the full length open reading frame; and
15 (d) a sequence of nucleotides that encodes the transmembrane protease that includes a sequence of amino acids encoded by a sequence of nucleotides that encodes such subunit and hybridizes under conditions of high stringency to DNA that is complementary to the mRNA transcript.

20 Exemplary MTSPs

The above discussion provides an overview and some details of the exemplified MTSPs. The following discussion provides additional details (see, also, EXAMPLES).

MTSP1 (matriptase)

- 25 Matriptase is a trypsin-like serine protease with broad spectrum cleavage activity and two potential regulatory modules. It was named "matriptase" because its ability to degrade the extra-cellular matrix and its trypsin-like activity. When isolated from breast cancer cells (or T-47D cell conditioned medium), matriptase has been reported to be primarily in an uncomplexed form.
- 30 Matriptase has been isolated from human milk; when isolated from human milk, matriptase was reported to be in one of two complexed forms, 95 kDa (the predominant form) and 110 kDa; uncomplexed matriptase was not detected.

-51-

(Liu, *et al.*, *J. Biol. Chem.* 274(26):18237-18242 (1999).) It has been proposed that matriptase exists as an uncomplexed protease when in its active state. In breast milk, matriptase has been reported to exist in complex with a fragment of hepatocyte growth factor inhibitor-1 (HAI-1), a Kuntz-type serine protease

5 inhibitor having activity against trypsin-like serine proteases.

Ecotin and Ecotin M84R/M85R are macromolecular inhibitors of serine proteases of the chymotrypsin fold and inhibit ductal branching, morphogenesis and differentiation of the explanted ductal prostate. PC-3 is a cell line derived from prostate cancer epithelial cells. Ecotin and M84R/M85R ecotin were found

10 to decrease tumor size and metastasis in PC-3 implanted nude mice.

Matriptase has been isolated and its encoding nucleic acids cloned from T-47D human breast cancer cell-conditioned medium (Lin *et al.* (1999) *J. Biol. Chem.* 274:18231-18236). Upon analysis of the cDNA, it was determined that the full length protease has 683 amino acids and contains three main structural

15 regions: a serine protease domain near the carboxyl-terminal region, four tandem low-density lipoprotein receptor domains, and two tandem complement subcomponents C1r and C1s.

Studies to identify additional serine proteases made by cancer cells were done using PC-3 cells. A serine protease termed "MT-SP1", reported to be a

20 transmembrane protease was cloned (Takeuchi *et al.* (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96:11054-11061). It was subsequently found the originally identified matriptase sequence is included in the translated sequence of the cDNA that encodes MT-SP1. The matriptase cDNA was reported to be a partial MT-SP1 cDNA and to lack 516 of the coding nucleotides (Takeuchi, *et al.*, *J.*

25 *Biol. Chem* 275:26333-26342 (2000).) Since the reported matriptase encoding cDNA sequence encoded a possible initiating methionine, it was proposed that alternative splicing could yield a protein lacking the N-terminal region of MTSP1.

Matriptase and MT-SP1 demonstrate trypsin-like protease activity and are

30 Type II transmembrane proteins with a common extracellular protease domain. Studies of substrate specificity of MT-SP1 reveal that protease-activated receptor 2 (PAR2) and single-chain urokinase-type plasminogen activator (sc-

-52-

uPA) are macromolecular substrates of MT-SP1. PAR2 is functions in inflammation, cytoprotection and/or cell adhesion, while sc-uPa is functions in tumor cell invasion and metastasis.

An exemplary nucleotide sequences encoding a human MTSP1 is set forth in SEQ ID Nos 1 and 2 (see also SEQ ID Nos. 49 and 50 for the protease domain thereof). As previously noted SEQ ID No. 1 sets for an MTSP1-encoding nucleic acid sequence. This sequence is the longer version and includes the protease domain, which is common to both variants. Nucleic acids encoding the MTSP that hybridizes to the nucleotide sequence set forth in SEQ ID No. 1 can be obtained by any method known in the art, *e.g.*, by PCR amplification using synthetic primers that hybridizes to the 3' and 5' ends of the sequence and/or by cloning from a cDNA or genomic library using a PCR amplification product or an oligonucleotide specific for the gene sequence (*e.g.*, as described in Section C herein). Homologs (*e.g.*, nucleic acids of the above-listed genes of species other than human) or other related sequences (*e.g.*, paralogs) and muteins can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular sequence provided as a probe using methods well known in the art for nucleic acid hybridization and cloning.

Isolated single chain protease domains of MTSP1 proteins from animals are provided herein. As shown herein, the single chain protease domain is catalytically active and can be used in a variety of drug screening assays, particularly *in vitro* proteolytic assays. Exemplary MTSP protease domains are set forth as the amino acids (615-855 of SEQ ID No. 2) encoded by nucleotides 1865-2587 of SEQ ID No. 1 (see, also, SEQ ID Nos. 49 and 50). The MTSP1 single chain protease domain is catalytically active

Muteins of the MTSP1 proteins are provided. In the activated double chain molecule, residue 731 forms a disulfide bond with the Cys at residue 604. In the single chain form, the residue at 731 in the protease domain is free. Muteins in which Cys residues, particularly the free Cys residue (amino acid 731 in SEQ ID No. 2) in the single chain protease domain are provided. Other muteins in which conservative amino acids replacements are effected and that retain proteolytic activity as a single chain are also provided. Such changes may

-53-

be systematically introduced and tested for activity in *in vitro* assays, such as those provided herein.

MTSP3

In a specific embodiment, a nucleic acid that encodes a MTSP,
5 designated MTSP3 is provided. In particular, the nucleic acid includes an open reading frame within the following sequence of nucleotides set forth in SEQ ID No. 3. In particular the protein is encoded by the open reading frame that begins at nucleotide 261 and ends at 1574.

Also provided are nucleic acid molecules that hybridize under conditions
10 of at least low stringency, preferably moderate stringency, more preferably high stringency to the following sequence of nucleic acids (SEQ ID No. 3), particularly to the open reading frame encompassed by nucleotides that encode a single protease domain thereof, or any domain of MTSP3

Also included are substantially purified MTSP3 zymogen, activated double
15 chains, single chain protease domains and double chain protease domains. These are encoded by a nucleic acid that includes sequence encoding a protease domain that exhibits proteolytic activity and that hybridizes to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID No. 3, typically under moderate, generally under high stringency, conditions and most preferably along
20 the full length of the protease domain. Splice variants are also contemplated herein.

In a preferred embodiment, the isolated nucleic acid fragment hybridizes to the nucleic acid having the nucleotide sequence set forth in SEQ ID No: 3 under high stringency conditions, and preferably comprises the sequence of
25 nucleotides set forth in any of SEQ ID Nos. 3 or comprises a portion thereof that encodes a transmembrane domain and may additionally include a LDLR domain, a scavenger-receptor cysteine rich (SRCR) domain and a serine protease catalytic domain or any other identified domain (see FIGURES) or comprises nucleic acid molecule that encodes the protein encoded by SEQ ID NO. 4.

30 The isolated nucleic acid fragment is DNA, including genomic or cDNA, or is RNA, or can include other components, such as protein nucleic acid. The isolated nucleic acid may include additional components, such as heterologous or

-54-

native promoters, and other transcriptional and translational regulatory sequences, these genes may be linked to other genes, such as reporter genes or other indicator genes or genes that encode indicators.

Also provided is an isolated nucleic acid molecule that includes the
5 sequence of molecules that is complementary to the nucleotide sequence encoding the MTSP or the portion thereof.

Also provided are fragments thereof that can be used as probes or primers and that contain at least about 10 nucleotides, more preferably 14 nucleotides, more preferably at least about 16 nucleotides, most preferably at
10 least about 30 nucleotides.

Hence provided herein are polypeptides that are encoded by such nucleic acid molecules. Included among those polypeptides are the MTSP3 protease domain or a polypeptide with conservative amino acid changes such that the specificity and protease activity remains substantially unchanged. In particular, a
15 substantially purified mammalian MTSP protein is provided that has a transmembrane domain and may additionally include a CUB domain, one or more of an LDLR domain(s), a scavenger-receptor cysteine rich (SRCR) domain and a serine protease catalytic domain is provided.

Also provided is a substantially purified protein comprising a sequence of
20 amino acids that has at least 60%, more preferably at least about 90%, most preferably at least about 95%, identity to the MTSP3, wherein the percentage identity is determined using standard algorithms and gap penalties that maximize the percentage identity. The human MTSP3 protein is most preferred, although other mammalian MTSP3 proteins are contemplated.

25 Muteins of MTSP3, particularly those in which Cys residues, such as the Cys310 in the single chain protease domain, is replaced with another amino acid that does not eliminate the activity, are provided.

MTSP4

Among the proteins provided herein is MTSP4. MTSP4 is highly
30 expressed in the liver, and is expressed in substantially lower levels in other tissues (see, EXAMPLES). It is also expressed in non-liver-derived tumors (see EXAMPLES), including Burkitt's lymphoma, colorectal adenocarcinoma

-55-

(SW480), lung carcinoma (A549), and in leukemic cells, indicating a role in one or more of tumor progression, tumor invasion, tumor growth and tumor metastases.

Also provided are nucleic acid molecules that hybridize under conditions
5 of at least low stringency, preferably moderate stringency, more preferably high stringency to the sequence of nucleic acids set forth in SEQ ID Nos. 5, 7 or 9), particularly to the open reading frame encompassed by nucleotides that encode a single protease domain thereof, or any domain of an MTSP4.

Also included are substantially purified MTSP4 zymogens, activated
10 double chains, single chain protease domains and double chain protease domains. These are encoded by a nucleic acid that includes sequence encoding a protease domain that exhibits proteolytic activity and that hybridizes to a nucleic acid molecule having a nucleotide sequence set forth in SEQ ID Nos. 5, 7 and 9, typically under moderate, generally under high stringency, conditions and
15 most preferably along the full length of the protease domain.

In a preferred embodiment, the isolated nucleic acid fragment hybridizes to the nucleic acid having the nucleotide sequence set forth in SEQ ID No: 5, 7 or 9 under high stringency conditions, and preferably comprises the sequence of nucleotides set forth in any of SEQ ID Nos. 5, 7 or 9 comprises a portion thereof
20 that encodes a transmembrane domain and may additionally include a LDLR domain, a scavenger-receptor cysteine rich (SRCR) domain and a serine protease catalytic domain or any other identified domain (see FIGURES) or comprises nucleic acid molecule that encodes the protein encoded by SEQ ID NO. 6, 9 or 10..

25 The isolated nucleic acid fragment is DNA, including genomic or cDNA, or is RNA, or can include other components, such as protein nucleic acid. The isolated nucleic acid may include additional components, such as heterologous or native promoters, and other transcriptional and translational regulatory sequences, these genes may be linked to other genes, such as reporter genes or
30 other indicator genes or genes that encode indicators.

-56-

Also provided is an isolated nucleic acid molecule that includes the sequence of molecules that is complementary to the nucleotide sequence encoding and MTSP4 or the portion thereof.

Also provided are fragments thereof that can be used as probes or
5 primers and that contain at least about 10 nucleotides, more preferably 14 nucleotides, more preferably at least about 16 nucleotides, most preferably at least about 30 nucleotides.

In particular nucleic acid molecules encoding two forms of MTSP4 are
provide. The encoded proteins are multi-domain, type II membrane-type serine
10 proteases and include a transmembrane domain at the N terminus followed by a CUB domain, 3 LDLR domains and a trypsin-like serine protease domain at the C terminus. The difference between the two forms, which are splice variants, is the absence in MTSP4-S of a 432-bp nucleotide sequence between the transmembrane and the CUB domains (see FIGURE 2; see, also SEQ ID Nos. 5-
15 10).

Also provided is a nucleic acid that encodes the extracellular protease domain of an MTSP4 is provided. Both forms of MTSP4 exemplified herein include a protease domain in common (see SEQ ID Nos. 5 and 6).

In particular, the extracellular protease domain of the MTSP4 proteins is
20 encoded by the open reading frame that begins at nucleotide 1 and ends at 708 (TGA) (SEQ ID No. 5. This open reading frame encodes a portion of the MSTP4 protein and includes the protease domain. Full length MSTP4 proteins (SEQ ID Nos. 7 and 9) include the above domain. The extracellular protease domain, as a single chain, and also an activated double chain, exhibit protease activity. The
25 disulfide bonds that form that two chain form of MTSP forms are likely between Cys415 and Cys535 for MTSP4-S, and between Cys559 and Cys679 for MTSP4-L.

For use of the single chain protease domain thereof, it is of interest to replace the free Cys (i.e. Cys535 (Cys679)) in the protease domain with another
30 amino acid, such as any amino acid that does not alter the function (such change is likely to be any amino acid). Thus, muteins of MTSP4, particularly

-57-

those in which Cys residues, such as the Cys535 and Cys679 in the single chain protease domains of MTSP4-S and MTSP4-L, respectively, are provided.

MTSP6

Nucleic acid and the encoded MTSP6 protein of an exemplary MTSP6 are also provided. The respective sequences are set forth in SEQ ID Nos. 11 and 12. The MTSP6 DNA and protein sequences were analyzed using DNA Strider (version 1.2). The ORF encoding the MTSP6 variant provided herein is composed of 1,362 bp, which translate into a 453-amino acid protein. MTSP6 is a multi-domain, type-II membrane-type serine protease containing a transmembrane domain (amino acids 48-68) at the N-terminus followed by a LDLRa domain (LDL receptor domain class a) (amino acids 72-108), a SR domain (Scavenger receptor Cys-rich domain)(amino acids 109-205), and a trypsin-like serine protease domain (amino acids 216-443) (see FIGURE 3). Muteins of MTSP6, particularly those in which Cys residues, such as the Cys324 in the single chain protease domain of MTSP6 are provided.

International PCT application No. WO 00/52044 describes MTSPs that resemble the MTSP6 provided herein. The polypeptide provided therein differs at single amino acid positions, such as 90 in SEQ ID No. 12 (Ala is replaced with a Thr), and significantly from the instant MTSP6 in that ten amino acids (amino acid nos. 46-55 in SEQ ID No. 12) are replaced with the eleven amino acids: phe glu val phe ser gln ser ser ser leu gly (SEQ ID No. 59) resulting in a protein that is one 454 amino acids long.

There are a few other amino acid sequence differences and a number of nucleic acid sequence differences. Significantly, there are substantial differences in the protease domain at amino acids 368-394 (368 ICNHRDVYGGIISPSMLCAGYLTGGVD---- 394; SEQ ID No. 12) are replaced at position 369-396 with amino acids:

369 DLQPQ—GRVRWHHLPLHALRGLPDGWRWN 396, where the differences from 368-394 (Seq ID No. 12) are indicated.

In addition, a second C-terminus truncated variant with an altered protease domain is identified in the PCT application. The variant is the same as the 454 variant through amino acid 261 thereof (corresponding to 160 of SEQ

-58-

ID No. 12 herein), followed by 33 amino acids (see SEQ ID No. 60 herein) that differ by virtue of a frame shift.

C. Tumor specificity and tissue expression profiles

Each MTSP has a characteristic tissue expression profile; the MTSPs in particular, although not exclusively expressed or activated in tumors, exhibit characteristic tumor tissue expression or activation profiles. In some instances, MTSPs may have different activity in a tumor cell from a non-tumor cell by virtue of a change in a substrate or cofactor thereof or other factor that would alter the apparent functional activity of the MTSP. Hence each can serve as a diagnostic marker for particular tumors, by virtue of a level of activity and/or expression or function in a subject (i.e. a mammal, particularly a human) with neoplastic disease, compared to a subject or subjects that do not have the neoplastic disease. In addition, detection of activity (and/or expression) in a particular tissue can be indicative of neoplastic disease. Shed MTSPs in body fluids can be indicative of neoplastic disease. Also, by virtue of the activity and/or expression profiles of each, they can serve as therapeutic targets, such as by administration of modulators of the activity thereof, or, as by administration of a prodrug specifically activated by one of the MTSPs.

Tissue expression profiles

20

MTSP3

The MTSP3 transcript was detected in lung carcinoma (LX-1), colon adenocarcinoma (CX-1), colon adenocarcinoma (GI-112) and ovarian carcinoma (GI-102). No apparent signal was detected in another form of lung carcinoma (GI-117), breast carcinoma (GI-101), pancreatic adenocarcinoma (GI-103) and prostatic adenocarcinoma (PC3). MTSP1 is expressed in breast cancers.

MTSP4

The MTSP4 transcript, a DNA fragment encoding part of the LDL receptor domain and the protease domain was used to probe an RNA blot composed of 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH). As in the northern analysis of gel blot, a very strong signal was observed in the liver. Signals in other tissues were

RECTIFIED SHEET (RULE 91)

-59-

observed in (decreasing signal level): fetal liver > heart = kidney = adrenal gland = testis = fetal heart and kidney = skeletal muscle = bladder = placenta > brain = spinal cord = colon = stomach = spleen = lymph node = bone marrow = trachea = uterus = pancreas = salivary gland = mammary gland = lung. MTSP4 is also expressed less abundantly in several tumor cell lines including HeLa S3 = leukemia K-562 = Burkitt's lymphomas (Raji and Daudi) = colorectal adenocarcinoma (SW480) > lung carcinoma (A549) = leukemia MOLT-4 = leukemia HL-60. PCR of the MTSP4 transcript from cDNA libraries made from several human primary tumors xenografted in nude mice (human tumor multiple tissue cDNA panel, catalog number K1522-1, CLONTECH) was performed using MTSP4-specific primers. The MTSP4 transcript was detected in breast carcinoma (GI-101), lung carcinoma (LX-1), colon adenocarcinoma (GI-112) and pancreatic adenocarcinoma (GI-103). No apparent signal was detected in another form of lung carcinoma (GI-117), colon adenocarcinoma (CX-1), ovarian carcinoma (GI-102) and prostatic adenocarcinoma (PC3). The MTSP4 transcript was also detected in LNCaP and PC-3 prostate cancer cell lines as well as in HT-1080 human fibrosarcoma cell line.

Gene expression profile of MTSP6 in normal and tumor tissues

To obtain information regarding the gene expression profile of the MTSP6 transcript, a 495 bp DNA fragment obtained from PCR reaction with primers Ch17-NSP-3 and NSP-4AS was used to probe an RNA blot composed of 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH). The strongest signal was observed in duodenum. Signal in other tissues were observed in (decreased signal level): Stomach > trachea = mammary gland = thyroid gland = salivary gland = pituitary gland = pancreas > kidney > lung > jejunum = ileum = ileocecum = appendix = fetal kidney > fetal lung. Very weak signals can also be detected in several other tissues.

MTSP6 is also expressed in several tumor cell lines including HeLa S3 > colorectal adenocarcinoma (SW480) > leukemia MOLT-4 > leukemia K-562. PCR analysis of the MTSP6 transcript from cDNA libraries made from several

-60-

human primary tumors xenografted in nude mice (human tumor multiple tissue cDNA panel, catalog number K1522-1, CLONTECH) was performed using MTSP6-specific primers (Ch17-NSP-3 and Ch17-NSP2AS). The MTSP6 transcript was strongly detected in lung carcinoma (LX-1), moderately detected
5 in pancreatic adenocarcinoma (GI-103), weakly detected in ovarian carcinoma (GI-102); and very weakly detected in colon adenocarcinoma (GI-112 and CX-1), breast carcinoma (GI-101), lung carcinoma (GI-117) and prostatic adenocarcinoma (PC3). The MTSP6 transcript was also detected in breast cancer cell line MDA-MB-231, prostate cancer cell line PC-3, but not in HT-1080
10 human fibrosarcoma cell line. MTSP6 is also expressed in mammary gland carcinoma cDNA (Clontech). MTSP6 is also over expressed in ovarian tumor cells.

D. Identification and isolation of MTSP protein genes

The MTSP proteins, or domains thereof, can be obtained by methods well
15 known in the art for protein purification and recombinant protein expression. Any method known to those of skill in the art for identification of nucleic acids that encode desired genes may be used. Any method available in the art can be used to obtain a full length (*i.e.*, encompassing the entire coding region) cDNA or genomic DNA clone encoding an MTSP protein. In particular, the polymerase
20 chain reaction (PCR) can be used to amplify a sequence identified as being differentially expressed in normal and tumor cells or tissues, *e.g.*, nucleic acids encoding an MTSP protein (SEQ. NOs: 1-12), in a genomic or cDNA library. Oligonucleotide primers that hybridize to sequences at the 3' and 5' termini of the identified sequences can be used as primers to amplify by PCR sequences
25 from a nucleic acid sample (RNA or DNA), preferably a cDNA library, from an appropriate source (*e.g.*, tumor or cancer tissue).

PCR can be carried out, *e.g.*, by use of a Perkin-Elmer Cetus thermal
cycler and Taq polymerase (Gene Amp[®]). The DNA being amplified can include mRNA or cDNA or genomic DNA from any eukaryotic species. One can choose
30 to synthesize several different degenerate primers, for use in the PCR reactions. It is also possible to vary the stringency of hybridization conditions used in priming the PCR reactions, to amplify nucleic acid homologs (*e.g.*, to obtain

-61-

- MTSP protein* sequences from species other than humans or to obtain human sequences with homology to MTSP protein) by allowing for greater or lesser degrees of nucleotide sequence similarity between the known nucleotide sequence and the nucleic acid homolog being isolated. For cross species
- 5 hybridization, low stringency conditions are preferred. For same species hybridization, moderately stringent conditions are preferred. After successful amplification of the nucleic acid containing all or a portion of the identified MTSP protein sequence or of a nucleic acid encoding all or a portion of an MTSP protein homolog, that segment may be molecularly cloned and sequenced, and
- 10 used as a probe to isolate a complete cDNA or genomic clone. This, in turn, will permit the determination of the gene's complete nucleotide sequence, the analysis of its expression, and the production of its protein product for functional analysis. Once the nucleotide sequence is determined, an open reading frame encoding the MTSP protein gene protein product can be determined by any
- 15 method well known in the art for determining open reading frames, for example, using publicly available computer programs for nucleotide sequence analysis. Once an open reading frame is defined, it is routine to determine the amino acid sequence of the protein encoded by the open reading frame. In this way, the nucleotide sequences of the entire MTSP protein genes as well as the amino acid
- 20 sequences of MTSP protein proteins and analogs may be identified.

- Any eukaryotic cell potentially can serve as the nucleic acid source for the molecular cloning of the MTSP protein gene. The nucleic acids can be isolated from vertebrate, mammalian, human, porcine, bovine, feline, avian, equine, canine, as well as additional primate sources, insects, plants, etc. The
- 25 DNA may be obtained by standard procedures known in the art from cloned DNA (e.g., a DNA "library"), by chemical synthesis, by cDNA cloning, or by the cloning of genomic DNA, or fragments thereof, purified from the desired cell (see, for example, Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New
- 30 York; Glover, D.M. (ed.), 1985, *DNA Cloning: A Practical Approach*, MRL Press, Ltd., Oxford, U.K. Vol. I, II). Clones derived from genomic DNA may contain regulatory and intron DNA regions in addition to coding regions; clones derived

-62-

from cDNA will contain only exon sequences. Whatever the source, the gene should be molecularly cloned into a suitable vector for propagation of the gene.

In the molecular cloning of the gene from genomic DNA, DNA fragments are generated, some of which will encode the desired gene. The DNA may be
5 cleaved at specific sites using various restriction enzymes. Alternatively, one may use DNase in the presence of manganese to fragment the DNA, or the DNA can be physically sheared, for example, by sonication. The linear DNA fragments can then be separated according to size by standard techniques, including but not limited to, agarose and polyacrylamide gel electrophoresis and
10 column chromatography.

Once the DNA fragments are generated, identification of the specific DNA fragment containing the desired gene may be accomplished in a number of ways. For example, a portion of the MTSP protein (of any species) gene (*e.g.*, a PCR amplification product obtained as described above or an oligonucleotide
15 having a sequence of a portion of the known nucleotide sequence) or its specific RNA, or a fragment thereof be purified and labeled, and the generated DNA fragments may be screened by nucleic acid hybridization to the labeled probe (Benton and Davis, *Science* 196:180 (1977); Grunstein and Hogness, *Proc. Natl. Acad. Sci. U.S.A.* 72:3961 (1975)). Those DNA fragments with substantial
20 homology to the probe will hybridize. It is also possible to identify the appropriate fragment by restriction enzyme digestion(s) and comparison of fragment sizes with those expected according to a known restriction map if such is available or by DNA sequence analysis and comparison to the known nucleotide sequence of MTSP protein. Further selection can be carried out on
25 the basis of the properties of the gene. Alternatively, the presence of the gene may be detected by assays based on the physical, chemical, or immunological properties of its expressed product. For example, cDNA clones, or DNA clones which hybrid-select the proper mRNA, can be selected which produce a protein that, *e.g.*, has similar or identical electrophoretic migration, isoelectric focusing
30 behavior, proteolytic digestion maps, antigenic properties, serine protease activity. If an anti-MTSP protein antibody is available, the protein may be identified by binding of labeled antibody to the putatively MTSP protein

-63-

synthesizing clones, in an ELISA (enzyme-linked immunosorbent assay)-type procedure.

Alternatives to isolating the MTSP protein genomic DNA include, but are not limited to, chemically synthesizing the gene sequence from a known
5 sequence or making cDNA to the mRNA that encodes the MTSP protein. For example, RNA for cDNA cloning of the MTSP protein gene can be isolated from cells expressing the protein. The identified and isolated nucleic acids can then be inserted into an appropriate cloning vector. A large number of vector-host systems known in the art may be used. Possible vectors include, but are not
10 limited to, plasmids or modified viruses, but the vector system must be compatible with the host cell used. Such vectors include, but are not limited to, bacteriophages such as lambda derivatives, or plasmids such as pBR322 or pUC plasmid derivatives or the Bluescript vector (Stratagene, La Jolla, CA). The insertion into a cloning vector can, for example, be accomplished by ligating the
15 DNA fragment into a cloning vector which has complementary cohesive termini. If the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules may be enzymatically modified. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers may comprise
20 specific chemically synthesized oligonucleotides encoding restriction endonuclease recognition sequences. In an alternative method, the cleaved vector and MTSP protein gene may be modified by homopolymeric tailing. Recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, etc., so that many copies of the gene
25 sequence are generated.

In an alternative method, the desired gene may be identified and isolated after insertion into a suitable cloning vector in a "shot gun" approach. Enrichment for the desired gene, for example, by size fractionization, can be done before insertion into the cloning vector.

30 In specific embodiments, transformation of host cells with recombinant DNA molecules that incorporate the isolated MTSP protein gene, cDNA, or synthesized DNA sequence enables generation of multiple copies of the gene.

-64-

Thus, the gene may be obtained in large quantities by growing transformants, isolating the recombinant DNA molecules from the transformants and, when necessary, retrieving the inserted gene from the isolated recombinant DNA.

E. Vectors, plasmids and cells that contain nucleic acids encoding an MTSP protein or protease domain thereof and expression of MTSP proteins

Vectors and cells

For recombinant expression of one or more of the MTSP proteins, the nucleic acid containing all or a portion of the nucleotide sequence encoding the MTSP protein can be inserted into an appropriate expression vector, *i.e.*, a vector that contains the necessary elements for the transcription and translation of the inserted protein coding sequence. The necessary transcriptional and translational signals can also be supplied by the native promoter for MTSP genes, and/or their flanking regions.

Also provided are vectors that contain nucleic acid encoding the MTSPs. Cells containing the vectors are also provided. The cells include eukaryotic and prokaryotic cells, and the vectors are any suitable for use therein.

Prokaryotic and eukaryotic cells, including endothelial cells, containing the vectors are provided. Such cells include bacterial cells, yeast cells, fungal cells, plant cells, insect cells and animal cells. The cells are used to produce an MTSP protein or protease domain thereof by growing the above-described cells under conditions whereby the encoded MTSP protein or protease domain of the MTSP protein is expressed by the cell, and recovering the expressed protease domain protein. For purposes herein, the protease domain is preferably secreted into the medium.

In one embodiment, the vectors include a sequence of nucleotides that encodes a polypeptide that has protease activity and contains all or a portion of only the protease domain, or multiple copies thereof, of an MTSP protein are provided. Also provided are vectors that comprise a sequence of nucleotides that encodes the protease domain and additional portions of an MTSP protein up to and including a full length MTSP protein, as well as multiple copies thereof, are also provided. The vectors may selected for expression of the MTSP protein or protease domain thereof in the cell or such that the MTSP protein is

-65-

expressed as a transmembrane protein. Alternatively, the vectors may include signals necessary for secretion of encoded proteins. When the protease domain is expressed the nucleic acid is preferably linked to a secretion signal, such as the *Saccharomyces cerevisiae* α mating factor signal sequence or a portion

5 thereof.

A variety of host-vector systems may be used to express the protein coding sequence. These include but are not limited to mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing
10 yeast vectors; or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system used, any one of a number of suitable transcription and translation elements may be used.

Any methods known to those of skill in the art for the insertion of DNA
15 fragments into a vector may be used to construct expression vectors containing a chimeric gene containing appropriate transcriptional/translational control signals and protein coding sequences. These methods may include *in vitro* recombinant DNA and synthetic techniques and *in vivo* recombinants (genetic recombination). Expression of nucleic acid sequences encoding MTSP protein, or domains,
20 derivatives, fragments or homologs thereof, may be regulated by a second nucleic acid sequence so that the genes or fragments thereof are expressed in a host transformed with the recombinant DNA molecule(s). For example, expression of the proteins may be controlled by any promoter/enhancer known in the art. In a specific embodiment, the promoter is not native to the genes for
25 MTSP protein. Promoters which may be used include but are not limited to the SV40 early promoter (Bernoist and Chambon, *Nature* 290:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980)), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. USA* 78:1441-1445 (1981)), the
30 regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982)); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Kamaroff et al., *Proc. Natl. Acad. Sci. USA* 75:3727-3731 (1978)) or the

RECTIFIED SHEET (RULE 91)

-66-

tac promoter (DeBoer et al., *Proc. Natl. Acad. Sci. USA* 80:21-25 (1983)); see also "Useful Proteins from Recombinant Bacteria": in *Scientific American* 242:79-94 (1980)); plant expression vectors containing the nopaline synthetase promoter (Herrar-Estrella et al., *Nature* 303:209-213 (1984)) or the cauliflower mosaic virus 35S RNA promoter (Gardner et al., *Nucleic Acids Res.* 9:2871 (1981)), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., *Nature* 310:115-120 (1984)); promoter elements from yeast and other fungi such as the Gal4 promoter, the alcohol dehydrogenase promoter, the phosphoglycerol kinase promoter, the alkaline phosphatase promoter, and the following animal transcriptional control regions that exhibit tissue specificity and have been used in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., *Cell* 38:639-646 (1984); Ornitz et al., *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409 (1986); MacDonald, *Hepatology* 7:425-515 (1987)); insulin gene control region which is active in pancreatic beta cells (Hanahan et al., *Nature* 315:115-122 (1985)), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., *Cell* 38:647-658 (1984); Adams et al., *Nature* 318:533-538 (1985); Alexander et al., *Mol. Cell Biol.* 7:1436-1444 (1987)), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., *Cell* 45:485-495 (1986)), albumin gene control region which is active in liver (Pinckert et al., *Genes and Devel.* 1:268-276 (1987)), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., *Mol. Cell Biol.* 5:1639-1648 (1985); Hammer et al., *Science* 235:53-58 (1987)), alpha-1 antitrypsin gene control region which is active in liver (Kelsey et al., *Genes and Devel.* 1:161-171 (1987)), beta globin gene control region which is active in myeloid cells (Mogam et al., *Nature* 315:338-340 (1985); Kollias et al., *Cell* 46:89-94 (1986)), myelin basic protein gene control region which is active in oligodendrocyte cells of the brain (Readhead et al., *Cell* 48:703-712 (1987)), myosin light chain-2 gene control region which is active in skeletal muscle (Sani, *Nature* 314:283-286 (1985)), and gonadotrophic releasing hormone gene control region which is active in gonadotrophs of the hypothalamus (Mason et al., *Science* 234:1372-1378 (1986)).

-67-

In a specific embodiment, a vector is used that contains a promoter operably linked to nucleic acids encoding an MTSP protein, or a domain, fragment, derivative or homolog, thereof, one or more origins of replication, and optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene).

5 Expression vectors containing the coding sequences, or portions thereof, of an MTSP protein, is made, for example, by subcloning the coding portions into the EcoRI restriction site of each of the three pGEX vectors (glutathione S-transferase expression vectors (Smith and Johnson, *Gene* 7:31-40 (1988))). This allows for the expression of products in the correct reading frame. Preferred
10 vectors and systems for expression of the protease domains of the MTSP proteins are well known *Pichia* vectors (available, for example, from Invitrogen, San Diego, CA), particularly those designed for secretion of the encoded proteins. One exemplary vector is described in the EXAMPLES.

Plasmids for transformation of *E. coli* cells, include, for example, the pET
15 expression vectors (see, U.S patent 4,952,496; available from NOVAGEN, Madison, WI; see, also literature published by Novagen describing the system). Such plasmids include pET 11a, which contains the T7lac promoter, T7 terminator, the inducible *E. coli* lac operator, and the lac repressor gene; pET 12a-c, which contains the T7 promoter, T7 terminator, and the *E. coli* ompT
20 secretion signal; and pET 15b and pET19b (NOVAGEN, Madison, WI), which contain a His-Tag™ leader sequence for use in purification with a His column and a thrombin cleavage site that permits cleavage following purification over the column; the T7-lac promoter region and the T7 terminator.

The vectors are introduced into host cells, such as *Pichia* cells and
25 bacterial cells, such as *E. coli*, and the proteins expressed therein. Preferred *Pichia* strains, include, for example, GS115. Preferred bacterial hosts contain chromosomal copies of DNA encoding T7 RNA polymerase operably linked to an inducible promoter, such as the lacUV promoter (see, U.S. Patent No. 4,952,496). Such hosts include, but are not limited to, the lysogenic *E. coli*
30 strain BL21(DE3).

-68-

Expression and production of proteins

The MTSP domains, derivatives and analogs be produced by various methods known in the art. For example, once a recombinant cell expressing an MTSP protein, or a domain, fragment or derivative thereof, is identified, the individual gene product can be isolated and analyzed. This is achieved by assays based on the physical and/or functional properties of the protein, including, but not limited to, radioactive labeling of the product followed by analysis by gel electrophoresis, immunoassay, cross-linking to marker-labeled product. The MTSP protein proteins may be isolated and purified by standard methods known in the art (either from natural sources or recombinant host cells expressing the complexes or proteins), including but not restricted to column chromatography (*e.g.*, ion exchange, affinity, gel exclusion, reversed-phase high pressure, fast protein liquid, etc.), differential centrifugation, differential solubility, or by any other standard technique used for the purification of proteins. Functional properties may be evaluated using any suitable assay known in the art.

Alternatively, once an MTSP protein or its domain or derivative is identified, the amino acid sequence of the protein can be deduced from the nucleotide sequence of the gene which encodes it. As a result, the protein or its domain or derivative can be synthesized by standard chemical methods known in the art (*e.g.* see Hunkapiller et al, *Nature* 310:105-111 (1984)).

Manipulations of MTSP protein sequences may be made at the protein level. Also contemplated herein are MTSP protein proteins, domains thereof, derivatives or analogs or fragments thereof, which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc.

-69-

In addition, domains, analogs and derivatives of an MTSP protein can be chemically synthesized. For example, a peptide corresponding to a portion of an MTSP protein, which includes the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer.

- 5 Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP protein sequence. Non-classical amino acids include but are not limited to the D-isomers of the common amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid, ϵ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino
- 10 isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D
- 15 (dextrorotary) or L (levorotary).

- In cases where natural products are suspected of being mutant or are isolated from new species, the amino acid sequence of the MTSP protein isolated from the natural source, as well as those expressed *in vitro*, or from synthesized expression vectors *in vivo* or *in vitro*, can be determined from
- 20 analysis of the DNA sequence, or alternatively, by direct sequencing of the isolated protein. Such analysis may be performed by manual sequencing or through use of an automated amino acid sequenator.

Modifications

- A variety of modification of the MTSP proteins and domains are
- 25 contemplated herein. An MTSP-encoding nucleic acid molecule modified by any of numerous strategies known in the art (Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York). The sequences can be cleaved at appropriate sites with restriction endonuclease(s), followed by further enzymatic modification if
- 30 desired, isolated, and ligated *in vitro*. In the production of the gene encoding a domain, derivative or analog of MTSP, care should be taken to ensure that the modified gene retains the original translational reading frame, uninterrupted by

-70-

translational stop signals, in the gene region where the desired activity is encoded.

Additionally, the MTSP-encoding nucleic acid molecules can be mutated *in vitro* or *in vivo*, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions and/or form new restriction endonuclease sites or destroy pre-existing ones, to facilitate further *in vitro* modification. Also, as described herein muteins with primary sequence alterations, such as replacements of Cys residues and elimination of glycosylation sites are contemplated. Such mutations may be effected by any technique for mutagenesis known in the art, including, but not limited to, chemical mutagenesis and *in vitro* site-directed mutagenesis (Hutchinson et al., *J. Biol. Chem.* 253:6551-6558 (1978)), use of TAB® linkers (Pharmacia). In one embodiment, for example, an MTSP protein or domain thereof is modified to include a fluorescent label. In other specific embodiments, the MTSP protein is modified to have a heterofunctional reagent, such heterofunctional reagents can be used to crosslink the members of the complex.

The MTSP proteins may be isolated and purified by standard methods known in the art (either from natural sources or recombinant host cells expressing the complexes or proteins), including but not restricted to column chromatography (*e.g.*, ion exchange, affinity, gel exclusion, reversed-phase high pressure, fast protein liquid, etc.), differential centrifugation, differential solubility, or by any other standard technique used for the purification of proteins. Functional properties may be evaluated using any suitable assay known in the art.

Alternatively, once a MTSP or its domain or derivative is identified, the amino acid sequence of the protein can be deduced from the nucleotide sequence of the gene which encodes it. As a result, the protein or its domain or derivative can be synthesized by standard chemical methods known in the art (*e.g.*, see Hunkapiller et al, *Nature*, 310:105-111 (1984)).

Manipulations of MTSP sequences may be made at the protein level. MTSP domains, derivatives or analogs or fragments, which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation,

-71-

phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule and other cellular ligand, are contemplated herein. Any of numerous chemical modifications may be carried out by known techniques, including but not limited to specific chemical

5 cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 , acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc.

In addition, domains, analogs and derivatives of a MTSP can be chemically synthesized. For example, a peptide corresponding to a portion of a

10 MTSP, which comprises the desired domain or which mediates the desired activity *in vitro* can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the MTSP sequence. Non-classical amino acids include but are not limited to the D-isomers of the common

15 amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-aminobutyric acid, ϵ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -

20 methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

F. SCREENING METHODS

The single chain protease domains, as shown herein, can be used in a

25 variety of methods to identify compounds that modulate the activity thereof. For MTSPs that exhibit higher activity or expression in tumor cells, compounds that inhibit the proteolytic activity are of particular interest. For any MTSPs that are active at lower levels in tumor cells, compounds or agents that enhance the activity are potentially of interest. In all instances the identified compounds will

30 include agents that are candidate cancer treatments.

Several types of assays are exemplified and described herein. It is understood that the protease domains may be used in other assays. It is shown

-72-

here, however, that the single chain protease domains exhibit catalytic activity. As such they are ideal for *in vitro* screening assays. They may also be used in binding assays.

- The MTSP3, MTSP4 and MTSP6 full length zymogens, activated
- 5 enzymes, single and double chain protease domains are contemplated for use in any screening assay known to those of skill in the art, including those provided herein. Hence the following description, if directed to proteolytic assays is intended to apply to use of a single chain protease domain or a catalytically active portion thereof of any MTSP, including an MTSP3, MTSP4 or an MTSP6.
- 10 Other assays, such as binding assays are provided herein, particularly for use with an MTSP3, MTSP4 or MTSP6, including any variants, such as splice variants thereof. MTSP3 and MTSP4 are of most interest in such assays.

1. Catalytic Assays for identification of agents that modulate the protease activity of an MTSP protein

- 15 Methods for identifying a modulator of the catalytic activity of an MTSP, particularly a single chain protease domain or catalytically active portion thereof, are provided herein. The methods can be practiced by: a) contacting the MTSP, particularly a single-chain domain thereof, with a substrate of the MTSP in the presence of a test substance, and detecting the proteolysis of the substrate,
- 20 whereby the activity of the MTSP is assessed, and comparing the activity to a control. For example, the control can be the activity of the MTSP assessed by contacting an MTSP, particularly a single-chain domain thereof, with a substrate of the MTSP, and detecting the proteolysis of the substrate, whereby the activity of the MTSP is assessed. The results in the presence and absence of
- 25 the test compounds are compared. A difference in the activity indicates that the test substance modulates the activity of the MTSP.

- In one embodiment a plurality of the test substances are screened simultaneously in the above screening method. In another embodiment, the MTSP is isolated from a target cell as a means for then identifying agents that
- 30 are potentially specific for the target cell.

In still another embodiment, The test substance is a therapeutic compound, and whereby a difference of the MTSP activity measured in the

-73-

presence and in the absence of the test substance indicates that the target cell responds to the therapeutic compound.

One method include the steps of (a) contacting the MTSP protein or protease domain thereof with one or a plurality of test compounds under
5 conditions conducive to interaction between the ligand and the compounds; and (b) identifying one or more compounds in the plurality that specifically binds to the ligand.

Another method provided herein includes the steps of a) contacting an MTSP protein or protease domain thereof with a substrate of the MTSP protein,
10 and detecting the proteolysis of the substrate, whereby the activity of the MTSP protein is assessed; b) contacting the MTSP protein with a substrate of the MTSP protein in the presence of a test substance, and detecting the proteolysis of the substrate, whereby the activity of the MTSP protein is assessed; and c)
15 comparing the activity of the MTSP protein assessed in steps a) and b), whereby the activity measured in step a) differs from the activity measured in step b) indicates that the test substance modulates the activity of the MTSP protein.

In another embodiment, a plurality of the test substances are screened simultaneously. In comparing the activity of an MTSP protein in the presence and absence of a test substance to assess whether the test substance is a
20 modulator of the MTSP protein, it is unnecessary to assay the activity in parallel, although such parallel measurement is preferred. It is possible to measure the activity of the MTSP protein at one time point and compare the measured activity to a historical value of the activity of the MTSP protein.

For instance, one can measure the activity of the MTSP protein in the
25 presence of a test substance and compare with historical value of the activity of the MTSP protein measured previously in the absence of the test substance, and *vice versa*. This can be accomplished, for example, by providing the activity of the MTSP protein on an insert or pamphlet provided with a kit for conducting the assay.

30 Methods for selecting substrates for a particular MTSP are described in the EXAMPLES, and particular proteolytic assays are exemplified.

-74-

Combinations and kits containing the combinations optionally including instructions for performing the assays are provided. The combinations include an MTSP protein and a substrate of the MTSP protein to be assayed; and, optionally reagents for detecting proteolysis of the substrate. The substrates, which are typically proteins subject to proteolysis by a particular MTSP protein, can be identified empirically by testing the ability of the MTSP protein to cleave the test substrate. Substrates that are cleaved most effectively (i.e., at the lowest concentrations and/or fastest rate or under desirable conditions), are identified.

- 10 Additionally provided herein is a kit containing the above-described combination. Preferably, the kit further includes instructions for identifying a modulator of the activity of an MTSP protein. Any MTSP protein is contemplated as target for identifying modulators of the activity thereof.

2. Binding assays

- 15 Also provided herein are methods for identification and isolation of agents, particularly compounds that bind to MTSPs. The assays are designed to identify agents that bind to the zymogen form, the single chain isolated protease domain (or a protein, other than an MTSP protein, that contains the protease domain of an MTSP protein), and to the activated form, including the activated form derived from the full length zymogen or from an extended protease domain. The identified compounds are candidates or leads for identification of compounds for treatments of tumors and other disorders and diseases involving aberrant angiogenesis. The MTSP proteins used in the methods include any MTSP protein as defined herein, and preferably use MTSP single chain domain or proteolytically active portion thereof.

- 25 A variety of methods are provided herein. These methods may be performed in solution or in solid phase reactions in which the MTSP protein(s) or protease domain(s) thereof are linked, either directly or indirectly via a linker, to a solid support. Screening assays are described in the Examples, and these assays have been used to identify candidate compounds.

30 For purposes herein, all binding assays described above are provided for MTSP3, MTSP4 and MTSP6. For MTSP1 (including any variant thereof) and

RECTIFIED SHEET (RULE 91)

-75-

thereof) and other such proteases, binding assays that employ the isolated single chain protease domain or a protein containing such domain (other than the MTSP from which the protease is derived) are provided.

- Methods for identifying an agent, such as a compound, that specifically
- 5 binds to an MTSP single chain protease domain or an MTSP, such as an MTSP3, MTSP4 or an MTSP6, are provided herein. The method can be practiced by (a) contacting the MTSP with one or a plurality of test agents under conditions conducive to binding between the MTSP and an agent; and (b) identifying one or more agents within the plurality that specifically binds to the MTSP.
- 10 For example, in practicing such methods the MTSP polypeptide is mixed with a potential binding partner or an extract or fraction of a cell under conditions that allow the association of potential binding partners with the polypeptide. After mixing, peptides, polypeptides, proteins or other molecules that have become associated with an MTSP are separated from the mixture. The binding partner
- 15 that bound to the MTSP can then be removed and further analyzed. To identify and isolate a binding partner, the entire protein, for instance the entire disclosed protein of SEQ ID Nos. 6, 8 10 or 12 can be used. Alternatively, a fragment of the protein can be used.

- A variety of methods can be used to obtain cell extracts. Cells can be
- 20 disrupted using either physical or chemical disruption methods. Examples of physical disruption methods include, but are not limited to, sonication and mechanical shearing. Examples of chemical lysis methods include, but are not limited to, detergent lysis and enzyme lysis. A skilled artisan can readily adapt methods for preparing cellular extracts in order to obtain extracts for use in the
- 25 present methods.

- Once an extract of a cell is prepared, the extract is mixed with the MTSP under conditions in which association of the protein with the binding partner can occur. A variety of conditions can be used, the most preferred being conditions that closely resemble conditions found in the cytoplasm of a human cell.
- 30 Features such as osmolarity, pH, temperature, and the concentration of cellular extract used, can be varied to optimize the association of the protein with the binding partner.

-76-

After mixing under appropriate conditions, the bound complex is separated from the mixture. A variety of techniques can be used to separate the mixture. For example, antibodies specific to an MTSP can be used to immunoprecipitate the binding partner complex. Alternatively, standard chemical separation techniques such as chromatography and density/sediment centrifugation can be used.

After removing the non-associated cellular constituents in the extract, the binding partner can be dissociated from the complex using conventional methods. For example, dissociation can be accomplished by altering the salt concentration or pH of the mixture.

To aid in separating associated binding partner pairs from the mixed extract, the MTSP can be immobilized on a solid support. For example, the protein can be attached to a nitrocellulose matrix or acrylic beads. Attachment of the protein or a fragment thereof to a solid support aids in separating peptide/binding partner pairs from other constituents found in the extract. The identified binding partners can be either a single protein or a complex made up of two or more proteins.

Alternatively, the nucleic acid molecules encoding the single chain proteases can be used in a yeast two-hybrid system. The yeast two-hybrid system has been used to identify other protein partner pairs and can readily be adapted to employ the nucleic acid molecules herein described.

Another *in vitro* binding assay, particularly for an MTSP3, MTSP4 or an MTSP6 uses a mixture of a polypeptide that contains at least the catalytic domain of one of these proteins and one or more candidate binding targets or substrates. After incubating the mixture under appropriate conditions, one determines whether the MTSP or a polypeptide fragment thereof containing the catalytic domain binds with the candidate substrate. For cell-free binding assays, one of the components includes or is coupled to a detectable label. The label may provide for direct detection, such as radioactivity, luminescence, optical or electron density, *etc.*, or indirect detection such as an epitope tag, an enzyme, *etc.* A variety of methods may be employed to detect the label depending on the nature of the label and other assay components. For example,

-77-

the label may be detected bound to the solid substrate or a portion of the bound complex containing the label may be separated from the solid substrate, and the label thereafter detected.

3. Detection of signal transduction

5 The cell surface location of the MTSPs suggests a role for some or all of these proteins in signal transduction. Assays for assessing signal transduction are well known to those of skill in the art, and may be adapted for use with the MTSP protein.

Assays for identifying agents that effect or alter signal transduction
10 mediated by an MTSP, particularly the full length or a sufficient portion to anchor the extracellular domain or a function portion thereof of an MTSP on the surface of a cell are provided. Such assays, include, for example, transcription based assays in which modulation of a transduced signal is assessed by detecting an effect on an expression from a reporter gene (see, *e.g.*, U.S. Patent No.
15 5,436,128).

4. Methods for Identifying Agents that Modulate the Expression a Nucleic Acid Encoding an MTSP, particularly an MTSP3, MTSP4 or MTSP6

Another embodiment provides methods for identifying agents that
20 modulate the expression of a nucleic acid encoding an MTSP, particularly an MTSP3, MTSP4 or MTSP. Such assays use any available means of monitoring for changes in the expression level of the nucleic acids encoding an MTSP, such as MTSP3 or MTSP4.

In one assay format, cell lines that contain reporter gene fusions between
25 the open reading frame of MTSP3, MTSP4 or MTSP6 or a domain thereof, particularly the protease domain and any assayable fusion partner may be prepared. Numerous assayable fusion partners are known and readily available including the firefly luciferase gene and the gene encoding chloramphenicol acetyltransferase (Alam *et al.*, *Anal. Biochem.* 188: 245-54 (1990)). Cell lines
30 containing the reporter gene fusions are then exposed to the agent to be tested under appropriate conditions and time. Differential expression of the reporter gene between samples exposed to the agent and control samples identifies

-78-

agents which modulate the expression of a nucleic acid encoding an MTSP3, MTSP4 or MTSP6.

Additional assay formats may be used to monitor the ability of the agent to modulate the expression of a nucleic acid encoding an MTSP3, MTSP4 or
5 MTSP6. For instance, mRNA expression may be monitored directly by hybridization to the nucleic acids. Cell lines are exposed to the agent to be tested under appropriate conditions and time and total RNA or mRNA is isolated by standard procedures (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR
10 CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press). Probes to detect differences in RNA expression levels between cells exposed to the agent and control cells may be prepared from the nucleic acids. It is preferable, but not necessary, to design probes which hybridize only with target nucleic acids under conditions of high stringency. Only highly
15 complementary nucleic acid hybrids form under conditions of high stringency. Accordingly, the stringency of the assay conditions determines the amount of complementarity which should exist between two nucleic acid strands in order to form a hybrid. Stringency should be chosen to maximize the difference in stability between the probe:target hybrid and potential probe:non-target hybrids.

Probes may be designed from the nucleic acids through methods known
20 in the art. For instance, the G + C content of the probe and the probe length can affect probe binding to its target sequence. Methods to optimize probe specificity are commonly available (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press); and Ausubel *et al.* (1995) CURRENT PROTOCOLS IN
25 MOLECULAR BIOLOGY, Greene Publishing Co., NY).

Hybridization conditions are modified using known methods (see, *e.g.*, Sambrook *et al.* (1989) MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed. Cold Spring Harbor Laboratory Press); and Ausubel *et al.* (1995) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Co., NY),
30 as required for each probe. Hybridization of total cellular RNA or RNA enriched for polyA RNA can be accomplished in any available format. For instance, total cellular RNA or RNA enriched for polyA RNA can be affixed to a solid support,

-79-

and the solid support exposed to at least one probe comprising at least one, or part of one of the nucleic acid molecules under conditions in which the probe will specifically hybridize. Alternatively, nucleic acid fragments comprising at least one, or part of one of the sequences can be affixed to a solid support, such as a porous glass wafer. The glass wafer can then be exposed to total cellular RNA or polyA RNA from a sample under conditions in which the affixed sequences will specifically hybridize. Such glass wafers and hybridization methods are widely available, for example, those disclosed by Beattie (WO 95/11755). By examining for the ability of a given probe to specifically hybridize to an RNA sample from an untreated cell population and from a cell population exposed to the agent, agents which up or down regulate the expression of a nucleic acid encoding the protein having the sequence of SEQ ID NO:3 or SEQ ID NO:4 are identified.

5. Methods for Identifying Agents that Modulate at Least One Activity of an MTPS, such as MTSP3, MTSP4 or MTSP6

Methods for identifying agents that modulate at least one activity of an MTPS, such as an MTSP3, MTSP4 or MTSP6 are provided. Such methods or assays may use any means of monitoring or detecting the desired activity.

In one format, the relative amounts of a protein between a cell population that has been exposed to the agent to be tested compared to an un-exposed control cell population may be assayed (*e.g.*, a prostate cancer cell line, a lung cancer cell line, a colon cancer cell line or a breast cancer cell line). In this format, probes, such as specific antibodies, are used to monitor the differential expression of the protein in the different cell populations. Cell lines or populations are exposed to the agent to be tested under appropriate conditions and time. Cellular lysates may be prepared from the exposed cell line or population and a control, unexposed cell line or population. The cellular lysates are then analyzed with the probe.

For example, N- and C- terminal fragments of the MTSP can be expressed in bacteria and used to search for proteins which bind to these fragments. Fusion proteins, such as His-tag or GST fusion to the N- or C-terminal regions of the MTSP, such as an MTSP3, MTSP4 or an MTSP6, can be prepared for use as

-80-

a substrate. These fusion proteins can be coupled to, for example, Glutathione-Sepharose beads and then probed with cell lysates. Prior to lysis, the cells may be treated with a candidate agent which may modulate an MTSP, such as an MTSP3, MTSP4 or an MTSP6, or proteins that interact with domains thereon.

- 5 Lysate proteins binding to the fusion proteins can be resolved by SDS-PAGE, isolated and identified by protein sequencing or mass spectroscopy, as is known in the art.

Antibody probes are prepared by immunizing suitable mammalian hosts in appropriate immunization protocols using the peptides, polypeptides or proteins
10 if they are of sufficient length (*e.g.*, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40 or more consecutive amino acids the MTSP protein, such as an MTSP3, an MTSP4 or an MTSP6), or if required to enhance immunogenicity, conjugated to suitable carriers. Methods for preparing immunogenic conjugates with carriers, such as bovine serum albumin (BSA), keyhole limpet hemocyanin
15 (KLH), or other carrier proteins are well known in the art. In some circumstances, direct conjugation using, for example, carbodiimide reagents may be effective; in other instances linking reagents such as those supplied by Pierce Chemical Co., Rockford, IL, may be desirable to provide accessibility to the hapten. Hapten peptides can be extended at either the amino or carboxy
20 terminus with a Cys residue or interspersed with cysteine residues, for example, to facilitate linking to a carrier. Administration of the immunogens is conducted generally by injection over a suitable time period and with use of suitable adjuvants, as is generally understood in the art. During the immunization schedule, titers of antibodies are taken to determine adequacy of antibody
25 formation.

Anti-peptide antibodies can be generated using synthetic peptides corresponding to, for example, the carboxy terminal amino acids of the MTSP. Synthetic peptides can be as small as 1-3 amino acids in length, but are preferably at least 4 or more amino acid residues long. The peptides can be
30 coupled to KLH using standard methods and can be immunized into animals, such as rabbits or ungulate. Polyclonal antibodies can then be purified, for example using Actigel beads containing the covalently bound peptide.

-81-

While the polyclonal antisera produced in this way may be satisfactory for some applications, for pharmaceutical compositions, use of monoclonal preparations is preferred. Immortalized cell lines which secrete the desired monoclonal antibodies may be prepared using the standard method of Kohler *et al.*, (*Nature* 256: 495-7 (1975)) or modifications which effect immortalization of lymphocytes or spleen cells, as is generally known. The immortalized cell lines secreting the desired antibodies are screened by immunoassay in which the antigen is the peptide hapten, polypeptide or protein. When the appropriate immortalized cell culture secreting the desired antibody is identified, the cells can be cultured either *in vitro* or by production *in vivo* via ascites fluid. Of particular interest, are monoclonal antibodies that recognize the catalytic domain of an MTSP, such as an MTSP3, MTSP4 or an MTSP6.

Additionally, the zymogen or two-chain forms the MTSP can be used to make monoclonal antibodies which recognize conformation epitopes. For peptide-directed monoclonal antibodies, peptides from the C1r/C1s domain can be used to generate anti-C1r/C1s domain monoclonal antibodies which can thereby block activation of the zymogen to the two-chain form of the MTSP. This domain can similarly be the substrate for other non-antibody compounds which bind to these preferred domains on either the single-chain or double-chain forms of the MTSP3, MTSP4 or MTSP6, and thereby modulate the activity of thereof or prevent its activation.

The desired monoclonal antibodies are then recovered from the culture supernatant or from the ascites supernatant. Fragments of the monoclonals or the polyclonal antisera which contain the immunologically significant portion can be used as antagonists, as well as the intact antibodies. Use of immunologically reactive fragments, such as the Fab, Fab', or F(ab')₂ fragments are often preferable, especially in a therapeutic context, as these fragments are generally less immunogenic than the whole immunoglobulin.

The antibodies or fragments may also be produced. Regions that bind specifically to the desired regions of receptor can also be produced in the context of chimeras with multiple species origin.

RECTIFIED SHEET (RULE 91)

-82-

Agents that are assayed in the above method can be randomly selected or rationally selected or designed.

The agents can be, as examples, peptides, small molecules, and carbohydrates. A skilled artisan can readily recognize that there is no limit as to
5 the structural nature of the agents.

The peptide agents can be prepared using standard solid phase (or solution phase) peptide synthesis methods, as is known in the art. In addition, the DNA encoding these peptides may be synthesized using commercially available oligonucleotide synthesis instrumentation and produced recombinantly
10 using standard recombinant production systems. The production using solid phase peptide synthesis is necessitated if non-gene-encoded amino acids are to be included.

G. Assay formats and selection of test substances

A variety of formats and detection protocols are known for performing
15 screening assays. Any such formats and protocols may be adapted for identifying modulators of MTSP protein activities. The following includes a discussion of exemplary protocols.

1. High throughput screening assays

Although the above-described assay can be conducted where a single
20 MTSP protein is screened, and/or a single test substance is screened for in one assay, the assay is preferably conducted in a high throughput screening mode, *i.e.*, a plurality of the MTSP proteins are screened against and/or a plurality of the test substances are screened for simultaneously (*See generally, High Throughput Screening: The Discovery of Bioactive Substances* (Devlin, Ed.)
25 Marcel Dekker, 1997; Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997); and Silverman et al., *Curr. Opin. Chem. Biol.*, 2(3):397-403 (1998)). For example, the assay can be conducted in a multi-well (*e.g.*, 24-, 48-, 96-, or 384-well), chip or array format.

High-throughput screening (HTS) is the process of testing a large number
30 of diverse chemical structures against disease targets to identify "hits" (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). Current state-of-the-art HTS operations are highly automated and computerized to handle

-83-

sample preparation, assay procedures and the subsequent processing of large volumes of data.

Detection technologies employed in high-throughput screens depend on the type of biochemical pathway being investigated (Sittampalam et al., *Curr. Opin. Chem. Biol.*, 1(3):384-91 (1997)). These methods include, radiochemical methods, such as the scintillation proximity assays (SPA), which can be adapted to a variety of enzyme assays (Lerner et al., *J. Biomol. Screening*, 1:135-143 (1996); Baker et al., *Anal. Biochem.*, 239:20-24 (1996); Baum et al., *Anal. Biochem.*, 237:129-134 (1996); and Sullivan et al., *J. Biomol. Screening*, 2:19-23 (1997)) and protein-protein interaction assays (Braunwalder et al., *J. Biomol. Screening*, 1:23-26 (1996); Sonatore et al., *Anal. Biochem.*, 240:289-297 (1996); and Chen et al., *J. Biol. Chem.*, 271:25308-25315 (1996)), and non-isotopic detection methods, including but are not limited to, colorimetric and luminescence detection methods, resonance energy transfer (RET) methods, time-resolved fluorescence (HTRF) methods, cell-based fluorescence assays, such as fluorescence resonance energy transfer (FRET) procedures (see, e.g., Gonzalez et al., *Biophys. J.*, 69:1272-1280 (1995)), fluorescence polarization or anisotropy methods (see, e.g., Jameson et al., *Methods Enzymol.*, 246:283-300 (1995); Jolley, *J. Biomol. Screening*, 1:33-38 (1996); Lynch et al., *Anal. Biochem.*, 247:77-82 (1997)), fluorescence correlation spectroscopy (FCS) and other such methods.

2. Test Substances

Test compounds, including small molecules and libraries and collections thereof can be screened in the above-described assays and assays described below to identify compounds that modulate the activity an MTSP protein. Rational drug design methodologies that rely on computational chemistry may be used to screen and identify candidate compounds.

The compounds identified by the screening methods include inhibitors, including antagonists, and may be agonists. Compounds for screening are any compounds and collections of compounds available, know or that can be prepared.

-84-

a. Selection of Compounds

Compounds can be selected for their potency and selectivity of inhibition of serine proteases, especially MTSP protein. As described herein, and as generally known, a target serine protease and its substrate are combined under
5 assay conditions permitting reaction of the protease with its substrate. The assay is performed in the absence of test compound, and in the presence of increasing concentrations of the test compound. The concentration of test compound at which 50% of the serine protease activity is inhibited by the test
10 Concentration) value for that compound. Within a series or group of test compounds, those having lower IC_{50} or EC_{50} values are considered more potent inhibitors of the serine protease than those compounds having higher IC_{50} or EC_{50} values. The IC_{50} measurement is often used for more simplistic assays, whereas the EC_{50} is often used for more complicated assays, such as those
15 employing cells.

Preferred compounds according to this aspect have an IC_{50} value of 100 nM or less as measured in an *in vitro* assay for inhibition of MTSP protein activity. Especially preferred compounds have an IC_{50} value of less than 100 nM.

20 The test compounds also are evaluated for selectivity toward a serine protease. As described herein, and as generally known, a test compound is assayed for its potency toward a panel of serine proteases and other enzymes and an IC_{50} value or EC_{50} value is determined for each test compound in each assay system. A compound that demonstrates a low IC_{50} value or EC_{50} value for
25 the target enzyme, *e.g.*, MTSP protein, and a higher IC_{50} value or EC_{50} value for other enzymes within the test panel (*e.g.*, urokinase tissue plasminogen activator, thrombin, Factor Xa), is considered to be selective toward the target enzyme. Generally, a compound is deemed selective if its IC_{50} value or EC_{50} value in the target enzyme assay is at least one order of magnitude less than the
30 next smallest IC_{50} value or EC_{50} value measured in the selectivity panel of enzymes.

-85-

Presently preferred compounds have an IC_{50} value of 100 nM or less as measured in an *in vitro* assay for inhibition of urokinase activity. Especially preferred compounds have an IC_{50} value in the *in vitro* urokinase inhibition assay that is at least one order of magnitude smaller than the IC_{50} value measured in the *in vitro* tPA inhibition assay. Compounds having a selectivity ratio of IC_{50} u-PA assay: IC_{50} MTSP protein assay of greater than 100 are especially preferred.

Compounds are also evaluated for their activity *in vivo*. The type of assay chosen for evaluation of test compounds will depend on the pathological condition to be treated or prevented by use of the compound, as well as the route of administration to be evaluated for the test compound.

For instance, to evaluate the activity of a compound to reduce tumor growth through inhibition of MTSP protein, the procedures described by Jankun et al., *Canc. Res.*, 57:559-563 (1997) to evaluate PAI-1 can be employed. Briefly, the ATCC cell lines DU145 and LnCaP are injected into SCID mice. After tumors are established, the mice are given test compound according to a dosing regime determined from the compound's *in vitro* characteristics. The Jankun et al. compound was administered in water. Tumor volume measurements are taken twice a week for about five weeks. A compound is deemed active if an animal to which the compound was administered exhibited decreased tumor volume, as compared to animals receiving appropriate control compounds.

Another *in vivo* experimental model designed to evaluate the effect of p-aminobenzamidine, a swine protease inhibitor, on reducing tumor volume is described by Billström et al., *Int. J. Cancer*, 61:542-547 (1995).

To evaluate the ability of a compound to reduce the occurrence of, or inhibit, metastasis, the procedures described by Kobayashi et al., *Int. J. Canc.*, 57:727-733d (1994) can be employed. Briefly, a murein xenograft selected for high lung colonization potential is injected into C57B1/6 mice i.v. (experimental metastasis) or s.c. into the abdominal wall (spontaneous metastasis). Various concentrations of the compound to be tested can be admixed with the tumor cells in Matrigel prior to injection. Daily i.p. injections of the test compound are made either on days 1-6 or days 7-13 after tumor inoculation. The animals are sacrificed about three or four weeks after tumor inoculation, and the lung tumor

-86-

colonies are counted. Evaluation of the resulting data permits a determination as to efficacy of the test compound, optimal dosing and route of administration.

The activity of the tested compounds toward decreasing tumor volume and metastasis can be evaluated in model described in Rabbani et al., *Int. J.*

- 5 *Cancer* 63:840-845 (1995) to evaluate their inhibitor. There, Mat LyLu tumor cells were injected into the flank of Copenhagen rats. The animals were implanted with osmotic minipumps to continuously administer various doses of test compound for up to three weeks. The tumor mass and volume of experimental and control animals were evaluated during the experiment, as were
- 10 metastatic growths. Evaluation of the resulting data permits a determination as to efficacy of the test compound, optimal dosing, and route of administration. Some of these authors described a related protocol in Xing et al., *Canc. Res.*, 57:3585-3593 (1997).

- To evaluate the inhibitory activity of a compound, a rabbit cornea
- 15 neovascularization model can be employed. Avery et al., *Arch. Ophthalmol.*, 108:1474-1475 (1990) describe anesthetizing New Zealand albino rabbits and then making a central corneal incision and forming a radial corneal pocket. A slow release prostaglandin pellet was placed in the pocket to induce neovascularization. Test compound was administered i.p. for five days, at which
- 20 time the animals were sacrificed. The effect of the test compound is evaluated by review of periodic photographs taken of the limbus, which can be used to calculate the area of neovascular response and, therefore, limbal neovascularization. A decreased area of neovascularization as compared with appropriate controls indicates the test compound was effective at decreasing or
- 25 inhibiting neovascularization.

- An angiogenesis model used to evaluate the effect of a test compound in preventing angiogenesis is described by Min et al., *Canc. Res.*, 56:2428-2433 (1996). C57BL6 mice receive subcutaneous injections of a Matrigel mixture containing bFGF, as the angiogenesis-inducing agent, with and without the test
- 30 compound. After five days, the animals are sacrificed and the Matrigel plugs, in which neovascularization can be visualized, are photographed. An experimental animal receiving Matrigel and an effective dose of test compound will exhibit

-87-

less vascularization than a control animal or an experimental animal receiving a less- or non-effective dose of compound.

An *in vivo* system designed to test compounds for their ability to limit the spread of primary tumors is described by Crowley et al., *Proc. Natl. Acad. Sci.*, 90:5021-5025 (1993). Nude mice are injected with tumor cells (PC3) engineered to express CAT (chloramphenicol acetyltransferase). Compounds to be tested for their ability to decrease tumor size and/or metastases are administered to the animals, and subsequent measurements of tumor size and/or metastatic growths are made. In addition, the level of CAT detected in various organs provides an indication of the ability of the test compound to inhibit metastasis; detection of less CAT in tissues of a treated animal versus a control animal indicates less CAT-expressing cells migrated to that tissue.

In vivo experimental modes designed to evaluate the inhibitory potential of a test serine protease inhibitors, using a tumor cell line F3II, the to be highly invasive, are described by Alonso et al., *Breast Canc. Res. Treat.*, 40:209-223 (1996). This group describes *in vivo* studies for toxicity determination, tumor growth, invasiveness, spontaneous metastasis, experimental lung metastasis, and an angiogenesis assay.

The CAM model (chick embryo chorioallantoic membrane model), first described by L. Ossowski in 1998 (*J. Cell Biol.*, 107:2437-2445 (1988)), provides another method for evaluating the urokinase inhibitory activity of a test compound. In the CAM model, tumor cells invade through the chorioallantoic membrane containing CAM with tumor cells in the presence of several serine protease inhibitors results in less or no invasion of the tumor cells through the membrane. Thus, the CAM assay is performed with CAM and tumor cells in the presence and absence of various concentrations of test compound. The invasiveness of tumor cells is measured under such conditions to provide an indication of the compound's inhibitory activity. A compound having inhibitory activity correlates with less tumor invasion.

The CAM model is also used in a standard assay of angiogenesis (*i.e.*, effect on formation of new blood vessels (Brooks et al., *Methods in Molecular Biology*, 129:257-269 (1999)). According to this model, a filter disc containing

-88-

an angiogenesis inducer, such as basic fibroblast growth factor (bFDG) is placed onto the CAM. Diffusion of the cytokine into the CAM induces local angiogenesis, which may be measured in several ways such as by counting the number of blood vessel branch points within the CAM directly below the filter disc. The ability of identified compounds to inhibit cytokine-induced angiogenesis can be tested using this model. A test compound can either be added to the filter disc that contains the angiogenesis inducer, be placed directly on the membrane or be administered systemically. The extent of new blood vessel formation in the presence and/or absence of test compound can be compared using this model. The formation of fewer new blood vessels in the presence of a test compound would be indicative of anti-angiogenesis activity. Demonstration of anti-angiogenesis activity for inhibitors of an MTSP protein indicates a role in angiogenesis for that MTSP protein.

b. Known serine protease inhibitors

Compounds for screening can be serine protease inhibitors, which can be tested for their ability to inhibit the activity of an MTSP, particularly an MTSP3, MTSP4, or MTSP6.

Exemplary, but not limiting serine proteases, include the following known serine protease inhibitors are used in the screening assays: Serine Protease Inhibitor 3 (SPI-3) (Chen, M.C., et al., *Cytokine*, 11(11):856-862 (1999)); Aprotinin (Iijima, R., et al., *J. Biochem. (Tokyo)*, 126(5):912-916 (1999)); Kazal-type serine protease inhibitor-like proteins (Niimi, T., et al., *Eur. J. Biochem.*, 266(1):282-292 (1999)); Kunitz-type serine protease inhibitor (Ravichandran, S., et al., *Acta Crystallogr. D. Biol. Crystallogr.*, 55(11):1814-1821 (1999)); Tissue factor pathway inhibitor-2/Matrix-associated serine protease inhibitor (TFPI-2/MSPI), (Liu, Y., et al., *Arch. Biochem. Biophys.*, 370(1):112-8 (1999)); Bukunin, (Cui, C.Y., et al., *J. Invest. Dermatol.*, 113(2):182-8 (1999)); Nafmostat mesilate (Ryo, R., et al., *Vox Sang.*, 76(4):241-6 (1999)); TPCK (Huang, Y., et al., *Oncogene*, 18(23):3431-9 (1999)); A synthetic cotton-bound serine protease inhibitor (Edwards, J.V., et al., *Wound Repair Regen.*, 7(2):106-18 (1999)); FUT-175 (Sawada, M., et al., *Stroke*, 30(3):644-50 (1999)); Combination of serine protease inhibitor FUT-0175 and thromboxane synthetase

- inhibitor OKY-046 (Kaminogo, M., et al., *Neurol. Med. Chir. (Tokyo)*, 38(11):704-8; discussion 708-9 (1998)); The rat serine protease inhibitor 2.1 gene (LeCam, A., et al., *Biochem. Biophys. Res. Commun.*, 253(2):311-4 (1998)); A new intracellular serine protease inhibitor expressed in the rat
- 5 pituitary gland complexes with granzyme B (Hill, R.M., et al., *FEBS Lett.*, 440(3):361-4 (1998)); 3,4-Dichloroisocoumarin (Hammed, A., et al., *Proc. Soc. Exp. Biol. Med.*, 219(2):132-7 (1998)); LEX032 (Bains, A.S., et al., *Eur. J. Pharmacol.*, 356(1):67-72 (1998)); N-tosyl-L-phenylalanine chloromethyl ketone (Dryjanski, M., et al., *Biochemistry*, 37(40):14151-6 (1998)); Mouse gene for
- 10 the serine protease inhibitor neuroserpin (P112) (Berger, P., et al., *Gene*, 214(1-2):25-33 (1998)); Rat serine protease inhibitor 2.3 gene (Paul, C., et al., *Eur. J. Biochem.*, 254(3):538-46 (1998)); Ecotin (Yang, S.Q., et al., *J. Mol. Biol.*, 279(4):945-57 (1998)); A 14 kDa plant-related serine protease inhibitor (Roch, P., et al., *Dev. Comp. Immunol.*, 22(1):1-12 (1998)); Matrix-associated serine
- 15 protease inhibitor TFPI-2/33 kDa MSPI (Rao, C.N., et al., *Int. J. Cancer*, 76(5):749-56 (1998)); ONO-3403 (Hiwasa, T., et al., *Cancer Lett.*, 126(2):221-5 (1998)); Bdelastasin (Moser, M., et al., *Eur. J. Biochem.*, 253(1):212-20 (1998)); Bikunin (Xu, Y., et al., *J. Mol. Biol.*, 276(5):955-66 (1998));
- Nafamostat mesilate (Mellgren, K., et al., *Thromb. Haemost.*, 79(2):342-7
- 20 (1998)); The growth hormone dependent serine protease inhibitor, Spi 2.1 (Maake, C., et al., *Endocrinology*, 138(12):5630-6 (1997)); Growth factor activator inhibitor type 2, a Kunitz-type serine protease inhibitor (Kawaguchi, T., et al., *J. Biol. Chem.*, 272(44):27558-64 (1997)); Heat-stable serine protease inhibitor protein from ovaries of the desert locust, *Schistocerca gregaria*
- 25 (Hamdaoui, A., et al., *Biochem. Biophys. Res. Commun.*, 238(2):357-60 (1997)); Bikunin, (Delaria, K.A., et al., *J. Biol. Chem.*, 272(18):12209-14 (1997)); Human placental bikunin (Marlor, C.W., et al., *J. Biol. Chem.*, 272(10):12202-8 (1997)); Hepatocyte growth factor activator inhibitor, a novel Kunitz-type serine protease inhibitor (Shimomura, T., et al., *J. Biol. Chem.*, 272(10):6370-6 (1997)); FUT-187, oral serine protease inhibitor, (Shiozaki, H., et al., *Gan To Kagaku Ryoho*, 23(14): 1971-9 (1996)); Extracellular matrix-associated serine protease inhibitors (Mr 33,000, 31,000, and 27,000 (Rao,
- 30

-90-

- C.N., et al., *Arch. Biochem. Biophys.*, 335(1):82-92 (1996)); An irreversible isocoumarin serine protease inhibitor (Palencia, D.D., et al., *Biol. Reprod.*, 55(3):536-42 (1996)); 4-(2-aminoethyl)-benzenesulfonyl fluoride (AEBSF) (Nakabo, Y., et al., *J. Leukoc. Biol.*, 60(3):328-36 (1996)); Neuroserpin
- 5 (Osterwalder, T., et al., *EMBO J.*, 15(12):2944-53 (1996)); Human serine protease inhibitor alpha-1-antitrypsin (Forney, J.R., et al., *J. Parasitol.*, 82(3):496-502 (1996)); Rat serine protease inhibitor 2.3 (Simar-Blanchet, A.E., et al., *Eur. J. Biochem.*, 236(2):638-48 (1996)); Gebaxate mesilate (parodi, F., et al., *J. Cardiothorac. Vasc. Anesth.*, 10(2):235-7 (1996)); Recombinant serine
- 10 protease inhibitor, CPTI II (Stankiewicz, M., et al., *Acta Biochim. Pol.*, 43(3):525-9 (1996)); A cysteine-rich serine protease inhibitor (Guamerin II) (Kim, D.R., et al., *J. Enzym. Inhib.*, 10(2):81-91 (1996)); Diisopropylfluorophosphate (Lundqvist, H., et al., *Inflamm. Res.*, 44(12):510-7 (1995)); Nexin 1 (Yu, D.W., et al., *J. Cell Sci.*, 108(Pt 12):3867-74 (1995)); LEX032 (Scalia, R., et al.,
- 15 *Shock*, 4(4):251-6 (1995)); Protease nexin I (Houenou, L.J., et al., *Proc. Natl. Acad. Sci. U.S.A.*, 92(3):895-9 (1995)); Chymase-directed serine protease inhibitor (Woodard S.L., et al., *J. Immunol.*, 153(11):5016-25 (1994)); N-alpha-tosyl-L-lysyl-chloromethyl ketone (TLCK) (Bourinbaier, A.S., et al., *Cell Immunol.*, 155(1):230-6 (1994)); Smpi56 (Ghendler, Y., et al., *Exp. Parasitol.*, 78(2):121-
- 20 31 (1994)); Schistosoma haematobium serine protease (Blanton, R.E., et al., *Mol. Biochem. Parasitol.*, 63(1):1-11 (1994)); Spi-1 (Warren, W.C., et al., *Mol. Cell Endocrinol.*, 98(1):27-32 (1993)); TAME (Jessop, J.J., et al., *Inflammation*, 17(5):613-31 (1993)); Antithrombin III (Kalaria, R.N., et al., *Am. J. Pathol.*, 143(3):886-93 (1993)); FOY-305 (Ohkoshi, M., et al., *Anticancer Res.*,
- 25 13(4):963-6 (1993)); Camostat mesilate (Senda, S., et al., *Intern. Med.*, 32(4):350-4 (1993)); Pigment epithelium-derived factor (Steele, F.R., et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90(4):1526-30 (1993)); Antistasin (Holstein, T.W., et al., *FEBS Lett.*, 309(3):288-92 (1992)); The vaccinia virus K2L gene encodes a serine protease inhibitor (Zhou, J., et al., *Virology*, 189(2):678-86
- 30 (1992)); Bowman-Birk serine-protease inhibitor (Werner, M.H., et al., *J. Mol. Biol.*, 225(3):873-89 (1992)); FUT-175 (Yanamoto, H., et al., *Neurosurgery*, 30(3):358-63 (1992)); FUT-175; (Yanamoto, H., et al., *Neurosurgery*,

-91-

- 30(3):351-6, discussion 356-7 (1992)); PAI-I (Yreadwell, B.V., et al., *J. Orthop. Res.*, 9(3):309-16 (1991)); 3,4-Dichloroisocoumarin (Rusbridge, N.M., et al., *FEBS Lett.*, 268(1):133-6 (1990)); Alpha 1-antichymotrypsin (Lindmark, B.E., et al., *Am. Rev. Respir. Dis.*, 141(4 Pt 1):884-8 (1990)); P-toluenesulfonyl-L-
- 5 arginine methyl ester (TAME) (Scuderi, P., *J. Immunol.*, 143(1):168-73 (1989)); Aprotinin (Seto, S., et al., *Adv. Exp. Med. Biol.*, 247B:49-54 (1989)); Alpha 1-antichymotrypsin (Abraham, C.R., et al., *Cell*, 52(4):487-501 (1988)); Contrapsin (Modha, J., et al., *Parasitology*, 96 (Pt 1):99-109 (1988)); (FOY-305) (Yamauchi, Y., et al., *Hiroshima J. Med. Sci.*, 36(1):81-7 No abstract available
- 10 (1987)); Alpha 2-antiplasmin (Holmes, W.E., et al., *J. Biol. Chem.*, 262(4):1659-64 (1987)); 3,4-dichloroisocoumarin (Harper, J.W., et al., *Biochemistry*, 24(8):1831-41 (1985)); Diisoprophylfluorophosphate (Tsutsui, K., et al., *Biochem. Biophys. Res. Commun.*, 123(1):271-7 (1984)); Gabexate mesilate (Hesse, B., et al., *Pharmacol. Res. Commun.*, 16(7):637-45 (1984)); Phenyl
- 15 methyl sulfonyl fluoride (Dufer, J., et al., *Scand. J. Haematol.*, 32(1):25-32 (1984)); Aprotinin (Seto, S., et al., *Hypertension*, 5(6):893-9 (1983)); Protease inhibitor Cl-2 (McPhalen, C.A., et al., *J. Mol. Biol.*, 168(2):445-7 (1983)); Phenylmethylsulfonyl fluoride (Sekar V., et al., *Biochem. Biophys. Res. Commun.*, 89(2):474-8 (1979)); PGE1 (Feinstein, M.D., et al., *Prostaglandine*, 14(6):1075-93 (1977)
- 20

c. Combinatorial libraries and other libraries

- The source of compounds for the screening assays, can be libraries, including, but are not limited to, combinatorial libraries. Methods for synthesizing combinatorial libraries and characteristics of such combinatorial
- 25 libraries are known in the art (See generally, *Combinatorial Libraries: Synthesis, Screening and Application Potential* (Cortese Ed.) Walter de Gruyter, Inc., 1995; Tietze and Lieb, *Curr. Opin. Chem. Biol.*, 2(3):363-71 (1998); Lam, *Anticancer Drug Des.*, 12(3):145-67 (1997); Blaney and Martin, *Curr. Opin. Chem. Biol.*, 1(1):54-9 (1997); and Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43
- 30 (1996)).

Methods and strategies for generating diverse libraries, primarily peptide- and nucleotide-based oligomer libraries, have been developed using molecular

-92-

- biology methods and/or simultaneous chemical synthesis methodologies (see, e.g., Dower et al., *Annu. Rep. Med. Chem.*, 26:271-280 (1991); Fodor et al., *Science*, 251:767-773 (1991); Jung et al., *Angew. Chem. Int. Ed. Engl.*, 31:367-383 (1992); Zuckerman et al., *Proc. Natl. Acad. Sci. USA*, 89:4505-4509 (1992); Scott et al., *Science*, 249:386-390 (1990); Devlin et al., *Science*, 249:404-406 (1990); Cwirla et al., *Proc. Natl. Acad. Sci. USA*, 87:6378-6382 (1990); and Gallop et al., *J. Medicinal Chemistry*, 37:1233-1251 (1994)). The resulting combinatorial libraries potentially contain millions of compounds and that can be screened to identify compounds that exhibit a selected activity.
- 10 The libraries fall into roughly three categories: fusion-protein-displayed peptide libraries in which random peptides or proteins are presented on the surface of phage particles or proteins expressed from plasmids; support-bound synthetic chemical libraries in which individual compounds or mixtures of compounds are presented on insoluble matrices, such as resin beads (see, e.g.,
- 15 Lam et al., *Nature*, 354:82-84 (1991)) and cotton supports (see, e.g., Eichler et al., *Biochemistry* 32:11035-11041 (1993)); and methods in which the compounds are used in solution (see, e.g., Houghten et al., *Nature*, 354:84-86 (1991); Houghten et al., *BioTechniques*, 31:412-421 (1992); and Scott et al., *Curr. Opin. Biotechnol.*, 5:40-48 (1994)). There are numerous examples of
- 20 synthetic peptide and oligonucleotide combinatorial libraries and there are many methods for producing libraries that contain non-peptidic small organic molecules. Such libraries can be based on basis set of monomers that are combined to form mixtures of diverse organic molecules or that can be combined to form a library based upon a selected pharmacophore monomer.
- 25 Either a random or a deterministic combinatorial library can be screened by the presently disclosed and/or claimed screening methods. In either of these two libraries, each unit of the library is isolated and/or immobilized on a solid support. In the deterministic library, one knows *a priori* a particular unit's location on each solid support. In a random library, the location of a particular
- 30 unit is not known *a priori* although each site still contains a single unique unit. Many methods for preparing libraries are known to those of skill in this art (see, e.g., Geysen et al., *Proc. Natl. Acad. Sci. USA*, 81:3998-4002 (1984),

-93-

- Houghten et al., *Proc. Natl. Acad. Sci. USA*, 81:5131-5135 (1985)).
- Combinatorial library generated by the any techniques known to those of skill in the art are contemplated (see, e.g., Table 1 of Schultz and Schultz, *Biotechnol. Prog.*, 12(6):729-43 (1996)) for screening; Bartel et al., *Science*, 261:1411-1418 (1993); Baumbach et al. *BioPharm*, (May):24-35 (1992); Bock et al. *Nature*, 355:564-566 (1992); Borman, S., Combinatorial chemists focus on small molecules molecular recognition, and automation, *Chem. Eng. News*, 2(12):29 (1996); Boublik, et al., Eukaryotic Virus Display: Engineering the Major Surface Glycoproteins of the Autographa California Nuclear Polyhedrosis Virus (ACNPV) for the Presentation of Foreign Proteins on the Virus Surface, *Bio/Technology*, 13:1079-1084 (1995); Brenner, et al., Encoded Combinatorial Chemistry, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5381-5383 (1992); Caflisch, et al., Computational Combinatorial Chemistry for *De Novo* Ligand Design: Review and Assessment, *Perspect. Drug Discovery Des.*, 3:51-84 (1995); Cheng, et al., Sequence-Selective Peptide Binding with a Peptido-A,B-*trans*-steroidal Receptor Selected from an Encoded Combinatorial Library, *J. Am. Chem. Soc.*, 118:1813-1814 (1996); Chu, et al., Affinity Capillary Electrophoresis to Identify the Peptide in A Peptide Library that Binds Most Tightly to Vancomycin, *J. Org. Chem.*, 58:648-652 (1993); Clackson, et al., Making Antibody Fragments Using Phage Display Libraries, *Nature*, 352:624-628 (1991); Combs, et al., Protein Structure-Based Combinatorial Chemistry: Discovery of Non-Peptide Binding Elements to Src SH3 Domain, *J. Am. Chem. Soc.*, 118:287-288 (1996); Cwirla, et al., Peptides On Phage: A Vast Library of Peptides for Identifying Ligands, *Proc. Natl. Acad. Sci. U.S.A.*, 87:6378-6382 (1990); Ecker, et al., Combinatorial Drug Discovery: Which Method will Produce the Greatest Value, *Bio/Technology*, 13:351-360 (1995); Ellington, et al., *In Vitro* Selection of RNA Molecules That Bind Specific Ligands, *Nature*, 346:818-822 (1990); Ellman, J.A., Variants of Benzodiazepines, *J. Am. Chem. Soc.*, 114:10997 (1992); Erickson, et al., *The Proteins*; Neurath, H., Hill, R.L., Eds.: Academic: New York, 1976; pp. 255-257; Felici, et al., *J. Mol. Biol.*, 222:301-310 (1991); Fodor, et al., Light-Directed, Spatially Addressable Parallel Chemical Synthesis, *Science*, 251:767-773 (1991); Francisco, et al., Transport and Anchoring of Beta-

RECTIFIED SHEET (RULE 91)

- Lactamase to the External Surface of *E. Coli.*, *Proc. Natl. Acad. Sci. U.S.A.*, 89:2713-2717 (1992); Georgiou, et al., Practical Applications of Engineering Gram-Negative Bacterial Cell Surfaces, *TIBTECH*, 11:6-10 (1993); Geysen, et al., Use of peptide synthesis to probe viral antigens for epitopes to a resolution of a single amino acid, *Proc. Natl. Acad. Sci. U.S.A.*, 81:3998-4002 (1984); Glaser, et al., Antibody Engineering by Condon-Based Mutagenesis in a Filamentous Phage Vector System, *J. Immunol.*, 149:3903-3913 (1992); Gram, et al., In vitro selection and affinity maturation of antibodies from a naive combinatorial immunoglobulin library, *Proc. Natl. Acad. Sci.*, 89:3576-3580 (1992); Han, et al., Liquid-Phase Combinatorial Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 92:6419-6423 (1995); Hoogenboom, et al., Multi-Subunit Proteins on the Surface of Filamentous Phage: Methodologies for Displaying Antibody (Fab) Heavy and Light Chains, *Nucleic Acids Res.*, 19:4133-4137 (1991); Houghten, et al., General Method for the Rapid Solid-Phase Synthesis of Large Numbers of Peptides: Specificity of Antigen-Antibody Interaction at the Level of Individual Amino Acids, *Proc. Natl. Acad. Sci. U.S.A.*, 82:5131-5135 (1985); Houghten, et al., The Use of Synthetic Peptide Combinatorial Libraries for the Determination of Peptide Ligands in Radio-Receptor Assays-Opioid-Peptides, *Bioorg. Med. Chem. Lett.*, 3:405-412 (1993); Houghten, et al., Generation and Use of Synthetic Peptide Combinatorial Libraries for Basic Research and Drug Discovery, *Nature*, 354:84-86 (1991); Huang, et al., Discovery of New Ligand Binding Pathways in Myoglobin by Random Mutagenesis, *Nature Struct. Biol.*, 1:226-229 (1994); Huse, et al., Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire In Phage Lambda, *Science*, 246:1275-1281 (1989); Janda, K.D., New Strategies for the Design of Catalytic Antibodies, *Biotechnol. Prog.*, 6:178-181 (1990); Jung, et al., Multiple Peptide Synthesis Methods and Their Applications, *Angew. Chem. Int. Ed. Engl.*, 31:367-486 (1992); Kang, et al., Linkage of Recognition and Replication Functions By Assembling Combinatorial Antibody Fab Libraries Along Phage Surfaces, *Proc. Natl. Acad. Sci. U.S.A.*, 88:4363-4366 (1991a); Kang, et al., Antibody Redesign by Chain Shuffling from Random Combinatorial Immunoglobulin Libraries, *Proc. Natl. Acad. Sci. U.S.A.*, 88:11120-11123 (1991b); Kay, et al., An M13 Phage Library

-95-

- Displaying Random 38-Amino-Acid-Peptides as a Source of Novel Sequences with Affinity to Selected Targets Genes, *Gene*, 128:59-65 (1993); Lam, et al., A new type of synthetic peptide library for identifying ligand-binding activity, *Nature*, 354:82-84 (1991) (published errata appear in *Nature*, 358:434 (1992)
- 5 and *Nature*, 360:768 (1992); Lebl, et al., One Bead One Structure Combinatorial Libraries, *Biopolymers (Pept. Sci.)*, 37:177-198 (1995); Lerner, et al., Antibodies without Immunization, *Science*, 258:1313-1314 (1992); Li, et al., Minimization of a Polypeptide Hormone, *Science*, 270:1657-1660 (1995); Light, et al., Display of Dimeric Bacterial Alkaline Phosphatase on the Major Coat Protein of
- 10 Filamentous Bacteriophage, *Bioorg. Med. Chem. Lett.*, 3:1073-1079 (1992); Little, et al., Bacterial Surface Presentation of Proteins and Peptides: An Alternative to Phage Technology, *Trends Biotechnol.*, 11:3-5 (1993); Marks, et al., By-Passing Immunization. Human Antibodies from V-Gene Libraries Displayed on Phage, *J. Mol. Biol.*, 222:581-597 (1991); Matthews, et al.,
- 15 Substrate Phage: Selection of Protease Substrates by Monovalent Phage Display, *Science*, 260:1113-1117 (1993); McCafferty, et al., Phage Enzymes: Expression and Affinity Chromatography of Functional Alkaline Phosphatase on the Surface of Bacteriophage, *Protein Eng.*, 4:955-961 (1991); Menger, et al., Phosphatase Catalysis Developed Via Combinatorial Organic Chemistry, *J. Org. Chem.*,
- 20 60:6666-6667 (1995); Nicolaou, et al., *Angew. Chem. Int. Ed. Engl.*, 34:2289-2291 (1995); Oldenburg, et al., Peptide Ligands for A Sugar-Binding Protein Isolated from a Random Peptide Library, *Proc. Natl. Acad. Sci. U.S.A.*, 89:5393-5397 (1992); Parmley, et al., Antibody-Selectable Filamentous fd Phage Vectors: Affinity Purification of Target Genes, *Genes*, 73:305-318 (1988); Pinilla, et al.,
- 25 Synthetic Peptide Combinatorial Libraries (SPCLS)--Identification of the Antigenic Determinant of Beta-Endorphin Recognized by Monoclonal Antibody-3E7, *Gene*, 128:71-76 (1993); Pinilla, et al., Review of the Utility of Soluble Combinatorial Libraries, *Biopolymers*, 37:221-240 (1995); Pistor, et al., Expression of Viral Hemagglutinin On the Surface of *E. Coli.*, *Klin. Wochenschr.*, 66:110-116
- 30 (1989); Pollack, et al., Selective Chemical Catalysis by an Antibody, *Science*, 234:1570-1572 (1986); Rigler, et al., Fluorescence Correlations, Single Molecule Detection and Large Number Screening: Applications in Biotechnology, *J.*

-96-

- Biotechnol.*, 41:177-186 (1995); Sarvetnick, et al., Increasing the Chemical Potential of the Germ-Line Antibody Repertoire, *Proc. Natl. Acad. Sci. U.S.A.*, 90:4008-4011 (1993); Sastry, et al., Cloning of the Immunological Repertoire in *Escherichia Coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library, *Proc. Natl. Acad. Sci. U.S.A.*, 86:5728-5732 (1989); Scott, et al., Searching for Peptide Ligands with an Epitope Library, *Science*, 249:386-390 (1990); Sears, et al., Engineering Enzymes for Bioorganic Synthesis: Peptide Bond Formation, *Biotechnol. Prog.*, 12:423-433 (1996); Simon, et al., Peptides: A Modular Approach to Drug Discovery, *Proc. Natl. Acad. Sci. U.S.A.*, 89:9367-9371 (1992); Still, et al., Discovery of Sequence-Selective Peptide Binding by Synthetic Receptors Using Encoded Combinatorial Libraries, *Acc. Chem. Res.*, 29:155-163 (1996); Thompson, et al., Synthesis and Applications of Small Molecule Libraries, *Chem. Rev.*, 96:555-600 (1996); Tramontano, et al., Catalytic Antibodies, *Science*, 234:1566-1570 (1986); Wrighton, et al., Small Peptides as Potent Mimetics of the Protein Hormone Erythropoietin, *Science*, 273:458-464 (1996); York, et al., Combinatorial mutagenesis of the reactive site region in plasminogen activator inhibitor I, *J. Biol. Chem.*, 266:8595-8600 (1991); Zebedee, et al., Human Combinatorial Antibody Libraries to Hepatitis B Surface Antigen, *Proc. Natl. Acad. Sci. U.S.A.*, 89:3175-3179 (1992); Zuckerman, et al., Identification of Highest-Affinity Ligands by Affinity Selection from Equimolar Peptide Mixtures Generated by Robotic Synthesis, *Proc. Natl. Acad. Sci. U.S.A.*, 89:4505-4509 (1992).

For example, peptides that bind to an MTSP protein or a protease domain of an MTSP protein can be identified using phage display libraries. In an exemplary embodiment, this method can include a) contacting phage from a phage library with the MTSP protein or a protease domain thereof; (b) isolating phage that bind to the protein; and (c) determining the identity of at least one peptide coded by the isolated phage to identify a peptide that binds to an MTSP protein.

-97-

H. Modulators of the activity of MTSP proteins

Provided herein are compounds, identified by screening or produced using the MTSP proteins or protease domain in other screening methods, that modulate the activity of an MTSP. These compounds act by directly interacting
5 with the MTSP protein or by altering transcription or translation thereof. Such molecules include, but are not limited to, antibodies that specifically react with an MTSP protein, particularly with the protease domain thereof, antisense nucleic acids that alter expression of the MTSP protein, antibodies, peptide mimetics and other such compounds.

10 1. Antibodies

Antibodies, including polyclonal and monoclonal antibodies, that specifically bind to the MTSP proteins provided herein, particularly to the single chain protease domains thereof are provided. Preferably, the antibody is a monoclonal antibody, and preferably, the antibody specifically binds to the
15 protease domain of the MTSP protein. In particular embodiments, antibodies to each of the single chain of protease domain of MTSP1, MTSP3, MTSP4 and MTSP6. Also provided are antibodies that specifically bind to any domain of MTSP3 or MTSP4, and to double chain forms thereof.

The MTSP protein and domains, fragments, homologs and derivatives
20 thereof may be used as immunogens to generate antibodies that specifically bind such immunogens. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments, and an Fab expression library. In a specific embodiment, antibodies to human MTSP protein are produced. In another embodiment, complexes formed from fragments of MTSP
25 protein, which fragments contain the serine protease domain, are used as immunogens for antibody production.

Various procedures known in the art may be used for the production of polyclonal antibodies to MTSP protein, its domains, derivatives, fragments or analogs. For production of the antibody, various host animals can be immunized
30 by injection with the native MTSP protein or a synthetic version, or a derivative of the foregoing, such as a cross-linked MTSP protein. Such host animals include but are not limited to rabbits, mice, rats, etc. Various adjuvants can be

-98-

used to increase the immunological response, depending on the host species, and include but are not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, and potentially useful human adjuvants such as bacille Calmette-Guerin (BCG) and corynebacterium parvum.

For preparation of monoclonal antibodies directed towards an MTSP protein or domains, derivatives, fragments or analogs thereof, any technique that provides for the production of antibody molecules by continuous cell lines in culture may be used. Such techniques include but are not restricted to the hybridoma technique originally developed by Kohler and Milstein (*Nature* 256:495-497 (1975)), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., *Immunology Today* 4:72 (1983)), and the EBV hybridoma technique to produce human monoclonal antibodies (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). In an additional embodiment, monoclonal antibodies can be produced in germ-free animals utilizing recent technology (PCT/US90/02545). Human antibodies may be used and can be obtained by using human hybridomas (Cote et al., *Proc. Natl. Acad. Sci. USA* 80:2026-2030 (1983)). Or by transforming human B cells with EBV virus *in vitro* (Cole et al., in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985)). Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci. USA* 81:6851-6855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) by splicing the genes from a mouse antibody molecule specific for the MTSP protein together with genes from a human antibody molecule of appropriate biological activity can be used.

Techniques described for the production of single chain antibodies (U.S. patent 4,946,778) can be adapted to produce MTSP protein-specific single chain antibodies. An additional embodiment uses the techniques described for the construction of Fab expression libraries (Huse et al., *Science* 246:1275-1281 (1989)) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity for MTSP protein or MTSP protein, or domains,

-99-

derivatives, or analogs thereof. Non-human antibodies can be "humanized" by known methods (*see, e.g.*, U.S. Patent No. 5,225,539).

Antibody fragments that contain the idiotypes of MTSP protein can be generated by techniques known in the art. For example, such fragments include but are not limited to: the F(ab')₂ fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments that can be generated by reducing the disulfide bridges of the F(ab')₂ fragment, the Fab fragments that can be generated by treating the antibody molecular with papain and a reducing agent, and Fv fragments.

10 In the production of antibodies, screening for the desired antibody can be accomplished by techniques known in the art, *e.g.*, ELISA (enzyme-linked immunosorbent assay). To select antibodies specific to a particular domain of the MTSP protein one may assay generated hybridomas for a product that binds to the fragment of the MTSP protein that contains such a domain

15 The foregoing antibodies can be used in methods known in the art relating to the localization and/or quantitation of MTSP proteins, *e.g.*, for imaging these proteins, measuring levels thereof in appropriate physiological samples, in diagnostic methods, etc.

In another embodiment, (*see infra*), anti-MTSP protein antibodies, or
20 fragments thereof, containing the binding domain are used as therapeutic agents.

2. Peptides and Peptide Mimetics

Provided herein are methods for identifying molecules that bind to and modulate the activity of MTSP proteins. Included among molecules that bind to MTSPs, particularly the single chain protease domain or catalytically active
25 fragments thereof, are peptides and peptide mimetics. Peptide mimetics are molecules or compounds that mimic the necessary molecular conformation of a ligand or polypeptide for specific binding to a target molecule such as, *e.g.*, an MTSP protein. In an exemplary embodiment, the peptides or peptide mimetics bind to the protease domain of the MTSP protein. Such peptides and peptide
30 mimetics include those of antibodies that specifically bind an MTSP protein and, preferably, bind to the protease domain of an MTSP protein. The peptides and

-100-

peptide mimetics identified by methods provided herein can be agonists or antagonists of MTSP proteins.

Such peptides and peptide mimetics are useful for diagnosing, treating, preventing, and screening for a disease or disorder associated with MTSP protein activity in a mammal. In addition, the peptides and peptide mimetics are useful for identifying, isolating, and purifying molecules or compounds that modulate the activity of an MTSP protein, or specifically bind to an MTSP protein, preferably, the protease domain of an MTSP protein. Low molecular weight peptides and peptide mimetics can have strong binding properties to a target molecule, *e.g.*, an MTSP protein or, preferably, to the protease domain of an MTSP protein.

Peptides and peptide mimetics that bind to MTSP proteins as described herein can be administered to mammals, including humans, to modulate MTSP protein activity. Thus, methods for therapeutic treatment and prevention of neoplastic diseases comprise administering a peptide or peptide mimetic compound in an amount sufficient to modulate such activity are provided. Thus, also provided herein are methods for treating a subject having such a disease or disorder in which a peptide or peptide mimetic compound is administered to the subject in a therapeutically effective dose or amount.

Compositions containing the peptides or peptide mimetics can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, compositions can be administered to a patient already suffering from a disease, as described above, in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Amounts effective for this use will depend on the severity of the disease and the weight and general state of the patient.

In prophylactic applications, compositions containing the peptides and peptide mimetics are administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective dose". In this use, the precise amounts again depend on the patient's state of health and weight.

-101-

Accordingly, the peptides and peptide mimetics that bind to an MTSP protein can be used generating pharmaceutical compositions containing, as an active ingredient, at least one of the peptides or peptide mimetics in association with a pharmaceutical carrier or diluent. The compounds can be administered, for example, by oral, pulmonary, parental (intramuscular, intraperitoneal, intravenous (IV) or subcutaneous injection), inhalation (via a fine powder formulation), transdermal, nasal, vaginal, rectal, or sublingual routes of administration and can be formulated in dosage forms appropriate for each route of administration (see, e.g., International PCT application Nos. WO 93/25221 and WO 94/17784; and European Patent Application 613,683).

Peptides and peptide mimetics that bind to MTSP proteins are useful *in vitro* as unique tools for understanding the biological role of MTSP proteins, including the evaluation of the many factors thought to influence, and be influenced by, the production of MTSP protein. Such peptides and peptide mimetics are also useful in the development of other compounds that bind to and modulate the activity of an MTSP protein, because such compounds provide important information on the relationship between structure and activity that should facilitate such development.

The peptides and peptide mimetics are also useful as competitive binders in assays to screen for new MTSP proteins or MTSP protein agonists. In such assay embodiments, the compounds can be used without modification or can be modified in a variety of ways; for example, by labeling, such as covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the materials thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups such as: radiolabels such as ¹²⁵I enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups. The compounds may also include spacers or linkers in cases where the compounds are to be attached to a solid support.

-102-

Moreover, based on their ability to bind to an MTSP protein, the peptides and peptide mimetics can be used as reagents for detecting MTSP proteins in living cells, fixed cells, in biological fluids, in tissue homogenates, in purified, natural biological materials, etc. For example, by labelling such peptides and peptide mimetics, one can identify cells having MTSP proteins. In addition, based on their ability to bind an MTSP protein, the peptides and peptide mimetics can be used in situ staining, FACS (fluorescence-activated cell sorting), Western blotting, ELISA, etc. In addition, based on their ability to bind to an MTSP protein, the peptides and peptide mimetics can be used in purification of MTSP protein polypeptides or in purifying cells expressing the MTSP protein polypeptides, *e.g.*, a polypeptide encoding the protease domain of an MTSP protein.

The peptides and peptide mimetics can also be used as commercial reagents for various medical research and diagnostic uses.

The activity of the peptides and peptide mimetics can be evaluated either *in vitro* or *in vivo* in one of the numerous models described in McDonald (1992) *Am. J. of Pediatric Hematology/Oncology*, 14:8-21, which is incorporated herein by reference.

3. Peptide and peptide mimetic therapy

Peptides and peptide mimetics that can bind to MTSP proteins or the protease domain of MTSP proteins and modulate the activity thereof, or have MTSP protein activity, can be used for treatment of neoplastic diseases. The peptides and peptide mimetics may be delivered, *in vivo* or *ex vivo*, to the cells of a subject in need of treatment. Further, peptides which have MTSP protein activity can be delivered, *in vivo* or *ex vivo*, to cells which carry mutant or missing alleles encoding the MTSP protein gene. Any of the techniques described herein or known to the skilled artisan can be used for preparation and *in vivo* or *ex vivo* delivery of such peptides and peptide mimetics that are substantially free of other human proteins. For example, the peptides can be readily prepared by expression in a microorganism or synthesis *in vitro*.

The peptides or peptide mimetics can be introduced into cells, *in vivo* or *ex vivo*, by microinjection or by use of liposomes, for example. Alternatively, the

-103-

peptides or peptide mimetics may be taken up by cells, *in vivo* or *ex vivo*, actively or by diffusion. In addition, extracellular application of the peptide or peptide mimetic may be sufficient to effect treatment of a neoplastic disease.

- Other molecules, such as drugs or organic compounds, that: 1) bind to an
5 MTSP protein or protease domain thereof; or 2) have a similar function or activity to an MTSP protein or protease domain thereof, may be used in methods for treatment.

4. Rational drug design

- The goal of rational drug design is to produce structural analogs of
10 biologically active polypeptides or peptides of interest or of small molecules or peptide mimetics with which they interact (e.g., agonists, antagonists, inhibitors) in order to fashion drugs which are, e.g., more active or stable forms thereof; or which, e.g., enhance or interfere with the function of a polypeptide *in vivo* (e.g., an MTSP protein). In one approach, one first determines the three-dimensional
15 structure of a protein of interest (e.g., an MTSP protein or polypeptide having a protease domain) or, for example, of a MTSP protein-ligand complex, by X-ray crystallography, by computer modeling or most typically, by a combination of approaches (see, e.g., Erickson *et al.* 1990). Also, useful information regarding the structure of a polypeptide may be gained by modeling based on the structure
20 of homologous proteins. In addition, peptides can be analyzed by an alanine scan. In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

- 25 Also, a polypeptide or peptide that binds to an MTSP protein or, preferably, the protease domain of an MTSP protein, can be selected by a functional assay, and then the crystal structure of this polypeptide or peptide can be determined. The polypeptide can be, for example, an antibody specific for an MTSP protein or the protein domain of an MTSP protein. This approach
30 can yield a pharmacore upon which subsequent drug design can be based. Further, it is possible to bypass the crystallography altogether by generating anti-idiotypic polypeptides or peptides, (anti-ids) to a functional,

-104-

pharmacologically active polypeptide or peptide that binds to an MTSP protein or protease domain of an MTSP protein. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original target molecule, e.g., an MTSP protein or polypeptide having an MTSP protein. The

5 anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

Thus, one may design drugs which have, e.g., improved activity or stability or which act as modulators (e.g., inhibitors, agonists, antagonists, etc.)

10 of MTSP protein activity, and are useful in the methods, particularly the methods for diagnosis, treatment, prevention, and screening of a neoplastic disease. By virtue of the availability of cloned MTSP protein sequences, sufficient amounts of the MTSP protein polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, the knowledge of the

15 amino acid sequence of an MTSP protein or the protease domain thereof, e.g., the protease domain encoded by the amino acid sequence of SEQ ID NO: 2, can provide guidance on computer modeling techniques in place of, or in addition to, X-ray crystallography.

20 **Methods of identifying peptides and peptide mimetics that bind to MTSP proteins**

Peptides having a binding affinity to the MTSP protein polypeptides provided herein (e.g., an MTSP protein or a polypeptide having a protease domain of an MTSP protein) can be readily identified, for example, by random peptide diversity generating systems coupled with an affinity enrichment

25 process. Specifically, random peptide diversity generating systems include the "peptides on plasmids" system (see, e.g., U.S. Patent Nos. 5,270,170 and 5,338,665); the "peptides on phage" system (see, e.g., U.S. Patent No. 6,121,238 and Cwirla, *et al.* (1990) *Proc. Natl. Acad. Sci. U.S.A.* 87:6378-6382); the "polysome system;" the "encoded synthetic library (ESL)"

30 system; and the "very large scale immobilized polymer synthesis" system (see, e.g., U.S. Patent No. 6,121,238; and Dower *et al.* (1991) *An. Rep. Med. Chem.* 26:271-280

-105-

For example, using the procedures described above, random peptides can generally be designed to have a defined number of amino acid residues in length (*e.g.*, 12). To generate the collection of oligonucleotides encoding the random peptides, the codon motif (NNK)*x*, where N is nucleotide A, C, G, or T

- 5 (equimolar; depending on the methodology employed, other nucleotides can be employed), K is G or T (equimolar), and *x* is an integer corresponding to the number of amino acids in the peptide (*e.g.*, 12) can be used to specify any one of the 32 possible codons resulting from the NNK motif: 1 for each of 12 amino acids, 2 for each of 5 amino acids, 3 for each of 3 amino acids, and only one of
10 the three stop codons. Thus, the NNK motif encodes all of the amino acids, encodes only one stop codon, and reduces codon bias.

- The random peptides can be presented, for example, either on the surface of a phage particle, as part of a fusion protein containing either the pIII or the pVIII coat protein of a phage fd derivative (peptides on phage) or as a fusion
15 protein with the LacI peptide fusion protein bound to a plasmid (peptides on plasmids). The phage or plasmids, including the DNA encoding the peptides, can be identified and isolated by an affinity enrichment process using immobilized MTSP protein polypeptide having a protease domain. The affinity enrichment process, sometimes called "panning," typically involves multiple rounds of
20 incubating the phage, plasmids, or polysomes with the immobilized MTSP protein polypeptide, collecting the phage, plasmids, or polysomes that bind to the MTSP protein polypeptide (along with the accompanying DNA or mRNA), and producing more of the phage or plasmids (along with the accompanying LacI-peptide fusion protein) collected.

25 **Characteristics of peptides and peptide mimetics**

- Typically, the molecular weight of preferred peptides or peptide mimetics is from about 250 to about 8,000 daltons. If the peptides are oligomerized, dimerized and/or derivatized with a hydrophilic polymer (*e.g.*, to increase the affinity and/or activity of the compounds), the molecular weights of such
30 peptides can be substantially greater and can range anywhere from about 500 to about 120,000 daltons, more preferably from about 8,000 to about 80,000 daltons. Such peptides can comprise 9 or more amino acids wherein the amino

-106-

acids are naturally occurring or synthetic (non-naturally occurring) amino acids. One skilled in the art would know how to determine the affinity and molecular weight of the peptides and peptide mimetics suitable for therapeutic and/or diagnostic purposes (*e.g.*, see Dower *et al.*, U.S. Patent No. 6,121,238).

5 The peptides may be covalently attached to one or more of a variety of hydrophilic polymers. Suitable hydrophilic polymers include, but are not limited to, polyalkylethers as exemplified by polyethylene glycol and polypropylene glycol, polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol, polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran
10 derivatives, etc. When the peptide compounds are derivatized with such polymers, their solubility and circulation half-lives can be increased with little, if any, diminishment in their binding activity. The peptide compounds may be dimerized and each of the dimeric subunits can be covalently attached to a hydrophilic polymer. The peptide compounds can be PEGylated, *i.e.*, covalently
15 attached to polyethylene glycol (PEG).

Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compounds are termed "peptide mimetics" or "peptidomimetics" (Luthman *et al.*, *A Textbook of Drug Design and*
20 *Development*, 14:386-406, 2nd Ed., Harwood Academic Publishers (1996); Joachim Grante (1994) *Angew. Chem. Int. Ed. Engl.*, 33:1699-1720; Fauchere (1986) *J. Adv. Drug Res.*, 15:29; Veber and Freidinger (1985) *TINS*, p. 392; and Evans *et al.* (1987) *J. Med. Chem.* 30:1229). Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an
25 equivalent or enhanced therapeutic or prophylactic effect. Preparation of peptidomimetics and structures thereof are known to those of skill in this art.

Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) may be used to generate more stable peptides. In addition, constrained
30 peptides containing a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo *et al.* (1992) *An. Rev. Biochem.*, 61:387, incorporated herein by reference); for

-107-

example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

Those skilled in the art would appreciate that modifications may be made to the peptides and mimetics without deleteriously effecting the biological or functional activity of the peptide. Further, the skilled artisan would know how to design non-peptide structures in three dimensional terms, that mimic the peptides that bind to a target molecule, *e.g.*, an MTSP protein or, preferably, the protease domain of MTSP proteins (see, *e.g.*, Eck and Sprang (1989) *J. Biol. Chem.*, 26: 17605-18795).

When used for diagnostic purposes, the peptides and peptide mimetics may be labeled with a detectable label and, accordingly, the peptides and peptide mimetics without such a label can serve as intermediates in the preparation of labeled peptides and peptide mimetics. Detectable labels can be molecules or compounds, which when covalently attached to the peptides and peptide mimetics, permit detection of the peptide and peptide mimetics *in vivo*, for example, in a patient to whom the peptide or peptide mimetic has been administered, or *in vitro*, *e.g.*, in a sample or cells. Suitable detectable labels are well known in the art and include, by way of example, radioisotopes, fluorescent labels (*e.g.*, fluorescein), and the like. The particular detectable label employed is not critical and is selected relative to the amount of label to be employed as well as the toxicity of the label at the amount of label employed. Selection of the label relative to such factors is well within the skill of the art.

Covalent attachment of a detectable label to the peptide or peptide mimetic is accomplished by conventional methods well known in the art. For example, when the ^{125}I radioisotope is employed as the detectable label, covalent attachment of ^{125}I to the peptide or the peptide mimetic can be achieved by incorporating the amino acid tyrosine into the peptide or peptide mimetic and then iodinating the peptide (see, *e.g.*, Weaner *et al.* (1994) *Synthesis and Applications of Isotopically Labelled Compounds*, pp. 137-140). If tyrosine is not present in the peptide or peptide mimetic, incorporation of tyrosine to the N or C terminus of the peptide or peptide mimetic can be achieved by well known chemistry. Likewise, ^{32}P can be incorporated onto the peptide or peptide

RECTIFIED SHEET (RULE 91)

-108-

mimetic as a phosphate moiety through, for example, a hydroxyl group on the peptide or peptide mimetic using conventional chemistry.

Labeling of peptidomimetics usually involves covalent attachment of one or more labels, directly or through a spacer (*e.g.*, an amide group), to non-interfering position(s) on the peptidomimetic that are predicted by quantitative structure-activity data and/or molecular modeling. Such non-interfering positions generally are positions that do not form direct contacts with the macromolecules(s) to which the peptidomimetic binds to produce the therapeutic effect. Derivatization (*e.g.*, labeling) of peptidomimetics should not substantially interfere with the desired biological or pharmacological activity of the peptidomimetic.

6. Methods of preparing peptides and peptide mimetics

Peptides that bind to MTSP proteins can be prepared by classical methods known in the art, for example, by using standard solid phase techniques. The standard methods include exclusive solid phase synthesis, partial solid phase synthesis methods, fragment condensation, classical solution synthesis, and even by recombinant DNA technology (see, *e.g.*, Merrifield (1963) *J. Am. Chem. Soc.*, 85:2149, incorporated herein by reference.)

Using the "encoded synthetic library" or "very large scale immobilized polymer synthesis" systems (see, *e.g.*, U.S. Patent No. 5,925,525, and 5,902,723); one can not only determine the minimum size of a peptide with the activity of interest, one can also make all of the peptides that form the group of peptides that differ from the preferred motif (or the minimum size of that motif) in one, two, or more residues. This collection of peptides can then be screened for ability to bind to the target molecule, *e.g.*, and MTSP protein or, preferably, the protease domain of an MTSP protein. This immobilized polymer synthesis system or other peptide synthesis methods can also be used to synthesize truncation analogs and deletion analogs and combinations of truncation and deletion analogs of the peptide compounds.

These procedures can also be used to synthesize peptides in which amino acids other than the 20 naturally occurring, genetically encoded amino acids are substituted at one, two, or more positions of the peptide. For instance,

-109-

naphthylalanine can be substituted for tryptophan, facilitating synthesis. Other synthetic amino acids that can be substituted into the peptides include L-hydroxypropyl, L-3, 4-dihydroxy-phenylalanyl, D amino acids such as L-D-hydroxylysyl and D-D-methylalanyl, L- α -methylalanyl, β amino acids, and
5 isoquinolyl. D amino acids and non-naturally occurring synthetic amino acids can also be incorporated into the peptides (see, *e.g.*, Roberts *et al.* (1983) *Unusual Amino/Acids in Peptide Synthesis*, 5(6):341-449).

The peptides may also be modified by phosphorylation (see, *e.g.*, W. Bannwarth *et al.* (1996) *Biorganic and Medicinal Chemistry Letters*,
10 6(17):2141-2146), and other methods for making peptide derivatives (see, *e.g.*, Hruby *et al.* (1990) *Biochem. J.*, 268(2):249-262). Thus, peptide compounds also serve as a basis to prepare peptide mimetics with similar biological activity.

Those of skill in the art recognize that a variety of techniques are available for constructing peptide mimetics with the same or similar desired
15 biological activity as the corresponding peptide compound but with more favorable activity than the peptide with respect to solubility, stability, and susceptibility to hydrolysis and proteolysis (see, *e.g.*, Morgan *et al.* (1989) *An. Rep. Med. Chem.*, 24:243-252). Methods for preparing peptide mimetics modified at the N-terminal amino group, the C-terminal carboxyl group, and/or
20 changing one or more of the amido linkages in the peptide to a non-amido linkage are known to those of skill in the art.

Amino terminus modifications include alkylating, acetylating, adding a carbobenzoyl group, forming a succinimide group, etc. (see, *e.g.*, Murray *et al.* (1995) *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., Vol. 1,
25 Manfred E. Wolf, ed., John Wiley and Sons, Inc.). C-terminal modifications include mimetics wherein the C-terminal carboxyl group is replaced by an ester, an amide or modifications to form a cyclic peptide.

In addition to N-terminal and C-terminal modifications, the peptide compounds, including peptide mimetics, can advantageously be modified with or
30 covalently coupled to one or more of a variety of hydrophilic polymers. It has been found that when peptide compounds are derivatized with a hydrophilic polymer, their solubility and circulation half-lives may be increased and their

-110-

immunogenicity is masked, with little, if any, diminishment in their binding activity. Suitable nonproteinaceous polymers include, but are not limited to, polyalkylethers as exemplified by polyethylene glycol and polypropylene glycol, polylactic acid, polyglycolic acid, polyoxyalkenes, polyvinylalcohol,

- 5 polyvinylpyrrolidone, cellulose and cellulose derivatives, dextran and dextran derivatives, etc. Generally, such hydrophilic polymers have an average molecular weight ranging from about 500 to about 100,000 daltons, more preferably from about 2,000 to about 40,000 daltons and, even more preferably, from about 5,000 to about 20,000 daltons. The hydrophilic polymers also can
10 have an average molecular weights of about 5,000 daltons, 10,000 daltons and 20,000 daltons.

- Methods for derivatizing peptide compounds or for coupling peptides to such polymers have been described (see, *e.g.*, Zallipsky (1995) *Bioconjugate Chem.*, 6:150-165; Monfardini *et al.* (1995) *Bioconjugate Chem.*, 6:62-69; U.S.
15 Pat. No. 4,640,835; U.S. Pat. No. 4,496,689; U.S. Pat. No. 4,301,144; U.S. Pat. No. 4,670,417; U.S. Pat. No. 4,791,192; U.S. Pat. No. 4,179,337 and WO 95/34326, all of which are incorporated by reference in their entirety herein).

- Other methods for making peptide derivatives are described, for example, in Hruby *et al.* (1990), *Biochem J.*, 268(2):249-262, which is incorporated
20 herein by reference. Thus, the peptide compounds also serve as structural models for non-peptidic compounds with similar biological activity. Those of skill in the art recognize that a variety of techniques are available for constructing compounds with the same or similar desired biological activity as a particular peptide compound but with more favorable activity with respect to
25 solubility, stability, and susceptibility to hydrolysis and proteolysis (see, *e.g.*, Morgan *et al.* (1989) *An. Rep. Med. Chem.*, 24:243-252, incorporated herein by reference). These techniques include replacing the peptide backbone with a backbone composed of phosphonates, amidates, carbamates, sulfonamides, secondary amines, and N-methylamino acids.

- 30 Peptide compounds may exist in a cyclized form with an intramolecular disulfide bond between the thiol groups of the cysteines. Alternatively, an intermolecular disulfide bond between the thiol groups of the cysteines can be

-111-

produced to yield a dimeric (or higher oligomeric) compound. One or more of the cysteine residues may also be substituted with a homocysteine.

I. CONJUGATES

- A conjugate, containing: a) a single chain protease domain (or
- 5 proteolytically active portion thereof) of an MTSP protein or an MTSP3, MTSP4 or MTSP6 full length zymogen, activated form thereof, or double or single chain protease domain thereof; and b) a targeting agent linked to the MTSP protein directly or via a linker, wherein the agent facilitates: i) affinity isolation or purification of the conjugate; ii) attachment of the conjugate to a surface; iii)
- 10 detection of the conjugate; or iv) targeted delivery to a selected tissue or cell, is provided herein. The conjugate can be a chemical conjugate or a fusion protein mixture thereof.

- The targeting agent is preferably a protein or peptide fragment, such as a tissue specific or tumor specific monoclonal antibody or growth factor or
- 15 fragment thereof linked either directly or via a linker to an MTSP protein or a protease domain thereof. The targeting agent may also be a protein or peptide fragment that contains a protein binding sequence, a nucleic acid binding sequence, a lipid binding sequence, a polysaccharide binding sequence, or a metal binding sequence, or a linker for attachment to a solid support. In a
- 20 particular embodiment, the conjugate contains a) the MTSP or portion thereof, as described herein; and b) a targeting agent linked to the MTSP protein directly or via a linker.

- Conjugates, such as fusion proteins and chemical conjugates, of the MTSP protein with a protein or peptide fragment (or plurality thereof)
- 25 that functions, for example, to facilitate affinity isolation or purification of the MTSP protein domain, attachment of the MTSP protein domain to a surface, or detection of the MTSP protein domain are provided. The conjugates can be produced by chemical conjugation, such as via thiol linkages, but are preferably produced by recombinant means as fusion proteins. In the fusion protein, the
- 30 peptide or fragment thereof is linked to either the N-terminus or C-terminus of the MTSP protein domain. In chemical conjugates the peptide or fragment thereof may be linked anywhere that conjugation can be effected, and there may

-112-

be a plurality of such peptides or fragments linked to a single MTSP protein domain or to a plurality thereof.

The targeting agent is preferably for *in vitro* delivery to a cell or tissue, and includes agents such as cell or tissue-specific antibodies, growth factors and
5 other factors expressed on specific cells; and other cell or tissue specific agents the promote directed delivery of a linked protein.

Most preferably the targeting agent specifically delivers the MTSP protein to selected cells by interaction with a cell surface protein and internalization of conjugate or MTSP protein portion thereof. These conjugate are used in a variety
10 of methods and are particularly suited for use in methods of activation of prodrugs, such as prodrugs that upon cleavage by the particular MTSP protein are cytotoxic. The prodrugs are administered prior to simultaneously with or subsequently to the conjugate. Upon delivery to the targeted cells, the protease activates the prodrug, which then exhibits is therapeutic effect, such as a
15 cytotoxic effect.

1. Conjugation

Conjugates with linked MTSP protein domains can be prepared either by chemical conjugation, recombinant DNA technology, or combinations of recombinant expression and chemical conjugation. The MTSP protein domains
20 and the targeting agent may be linked in any orientation and more than one targeting agents and/or MTSP protein domains may be present in a conjugate.

a. Fusion proteins

Fusion proteins are proved herein. A fusion protein contains: a) one or a plurality of domains of an MTSP proteins and b) a targeting agent. The fusion
25 proteins are preferably produced by recombinant expression of nucleic acids that encode the fusion protein.

b. Chemical conjugation

To effect chemical conjugation herein, the MTSP protein domain is linked via one or more selected linkers or directly to the targeting agent. Chemical
30 conjugation must be used if the targeted agent is other than a peptide or protein, such a nucleic acid or a non-peptide drug. Any means known to those of skill in the art for chemically conjugating selected moieties may be used.

-113-

2. Linkers

Linkers for two purposes are contemplated herein. The conjugates may include one or more linkers between the MTSP protein portion and the targeting agent. Additionally, linkers are used for facilitating or enhancing immobilization of an MTSP protein or portion thereof on a solid support, such as a microtiter plate, silicon or silicon-coated chip, glass or plastic support, such as for high throughput solid phase screening protocols.

Any linker known to those of skill in the art for preparation of conjugates may be used herein. These linkers are typically used in the preparation of chemical conjugates; peptide linkers may be incorporated into fusion proteins.

Linkers can be any moiety suitable to associate a domain of MTSP protein and a targeting agent. Such linkers and linkages include, but are not limited to, peptidic linkages, amino acid and peptide linkages, typically containing between one and about 60 amino acids, more generally between about 10 and 30 amino acids, chemical linkers, such as heterobifunctional cleavable cross-linkers, including but are not limited to, N-succinimidyl (4-iodoacetyl)-aminobenzoate, sulfosuccinimidyl (4-iodoacetyl)-aminobenzoate, 4-succinimidyl-oxycarbonyl-a-(2-pyridyldithio)toluene, sulfosuccinimidyl-6- [a-methyl-a-(pyridyldithiol)-toluamido] hexanoate, N-succinimidyl-3-(-2-pyridyldithio) - propionate, succinimidyl 6[3-(-2-pyridyldithio)-propionamido] hexanoate, sulfosuccinimidyl 6[3-(-2-pyridyldithio)-propionamido] hexanoate, 3-(2-pyridyldithio)-propionyl hydrazide, Ellman's reagent, dichlorotriazinic acid, and S-(2-thiopyridyl)-L-cysteine. Other linkers include, but are not limited to peptides and other moieties that reduce steric hindrance between the domain of MTSP protein and the targeting agent, intracellular enzyme substrates, linkers that increase the flexibility of the conjugate, linkers that increase the solubility of the conjugate, linkers that increase the serum stability of the conjugate, photocleavable linkers and acid cleavable linkers.

-114-

Other exemplary linkers and linkages that are suitable for chemically linked conjugates include, but are not limited to, disulfide bonds, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds are produced using

5 heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups to which reactive maleimido groups or thiol groups can be attached on the other. Other linkers include, acid cleavable linkers, such as bismaleimideoxy propane, acid labile-transferrin conjugates

10 and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; cross linkers that are cleaved upon exposure to UV or visible light and linkers, such as the various domains, such as C_H1, C_H2, and C_H3, from the constant region of human IgG, (see, Batra *et al. Molecular Immunol.*, 30:379-386 (1993)). In some embodiments, several linkers may be included in

15 order to take advantage of desired properties of each linker.

Chemical linkers and peptide linkers may be inserted by covalently coupling the linker to the domain of MTSP protein and the targeting agent. The heterobifunctional agents, described below, may be used to effect such covalent coupling. Peptide linkers may also be linked by expressing DNA encoding the

20 linker and TA, linker and targeted agent, or linker, targeted agent and TA as a fusion protein. Flexible linkers and linkers that increase solubility of the conjugates are contemplated for use, either alone or with other linkers are also contemplated herein.

a) Acid cleavable, photocleavable and heat sensitive linkers

25 Acid cleavable linkers, photocleavable and heat sensitive linkers may also be used, particularly where it may be necessary to cleave the domain of MTSP protein to permit it to be more readily accessible to reaction. Acid cleavable linkers include, but are not limited to, bismaleimideoxy propane; and adipic acid dihydrazide linkers (see, *e.g.*, Fattom *et al. (1992) Infection & Immun.*

30 *60*:584-589) and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see, *e.g.*, Welhöner *et al. (1991) J. Biol. Chem.* 266:4309-4314).

RECTIFIED SHEET (RULE 91)

-115-

Photocleavable linkers are linkers that are cleaved upon exposure to light (see, *e.g.*, Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which linkers are herein incorporated by reference), thereby releasing the targeted agent upon exposure to light. Photocleavable linkers that are cleaved upon exposure to light are known (see, *e.g.*, Hazum *et al.* (1981) in *Pept., Proc. Eur. Pept. Symp.*, 16th, Brunfeldt, K (Ed), pp. 105-110, which describes the use of a nitrobenzyl group as a photocleavable protective group for cysteine; Yen *et al.* (1989) *Makromol. Chem* 190:69-82, which describes water soluble photocleavable copolymers, including hydroxypropylmethacrylamide copolymer, glycine copolymer, fluorescein copolymer and methylrhodamine copolymer; Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which describes a cross-linker and reagent that undergoes photolytic degradation upon exposure to near UV light (350 nm); and Senter *et al.* (1985) *Photochem. Photobiol* 42:231-237, which describes nitrobenzyloxycarbonyl chloride cross linking reagents that produce photocleavable linkages), thereby releasing the targeted agent upon exposure to light. Such linkers would have particular use in treating dermatological or ophthalmic conditions that can be exposed to light using fiber optics. After administration of the conjugate, the eye or skin or other body part can be exposed to light, resulting in release of the targeted moiety from the conjugate. Such photocleavable linkers are useful in connection with diagnostic protocols in which it is desirable to remove the targeting agent to permit rapid clearance from the body of the animal.

b) Other linkers for chemical conjugation

Other linkers, include trityl linkers, particularly, derivatized trityl groups to generate a genus of conjugates that provide for release of therapeutic agents at various degrees of acidity or alkalinity. The flexibility thus afforded by the ability to preselect the pH range at which the therapeutic agent will be released allows selection of a linker based on the known physiological differences between tissues in need of delivery of a therapeutic agent (see, *e.g.*, U.S. Patent No. 5,612,474). For example, the acidity of tumor tissues appears to be lower than that of normal tissues.

-116-

c) Peptide linkers

The linker moieties can be peptides. Peptide linkers can be employed in fusion proteins and also in chemically linked conjugates. The peptide typically has from about 2 to about 60 amino acid residues, for example from about 5 to about 40, or from about 10 to about 30 amino acid residues. The length selected will depend upon factors, such as the use for which the linker is included.

Peptide linkers are advantageous when the targeting agent is proteinaceous. For example, the linker moiety can be a flexible spacer amino acid sequence, such as those known in single-chain antibody research. Examples of such known linker moieties include, but are not limited to, peptides, such as $(\text{Gly}_m\text{Ser})_n$ and $(\text{Ser}_m\text{Gly})_n$, in which n is 1 to 6, preferably 1 to 4, more preferably 2 to 4, and m is 1 to 6, preferably 1 to 4, more preferably 2 to 4, enzyme cleavable linkers and others.

Additional linking moieties are described, for example, in Huston *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883, 1988; Whitlow, M., *et al.*, *Protein Engineering* 6:989-995, 1993; Newton *et al.*, *Biochemistry* 35:545-553, 1996; A. J. Cumber *et al.*, *Bioconj. Chem.* 3:397-401, 1992; Ladurner *et al.*, *J. Mol. Biol.* 273:330-337, 1997; and U.S. Patent. No. 4,894,443. In some embodiments, several linkers may be included in order to take advantage of desired properties of each linker.

3. Targeting agents

Any agent that facilitates detection, immobilization, or purification of the conjugate is contemplated for use herein. For chemical conjugates any moiety that has such properties is contemplated; for fusion proteins, the targeting agent is a protein, peptide or fragment thereof that sufficient to effects the targeting activity. Preferred targeting agents are those that deliver the MTSP protein or portion thereof to selected cells and tissues. Such agents include tumor specific monoclonal antibodies and portions thereof, growth factors, such as FGF, EGF, PDGF, VEGF, cytokines, including chemokines, and other such agents.

4. Nucleic acids, plasmids and cells

-117-

Isolated nucleic acid fragments encoding fusion proteins are provided. The nucleic acid fragment that encodes the fusion protein includes: a) nucleic acid encoding a protease domain of an MTSP protein encoded by a nucleic acid that hybridizes to a nucleic acid having the nucleotide sequence set forth in the SEQ. ID NO:1; and b) nucleic acid encoding a protein, peptide or effective fragment thereof that facilitates: i) affinity isolation or purification of the fusion protein; ii) attachment of the fusion protein to a surface; or iii) detection of the fusion protein. Preferably, the nucleic acid is DNA..

Plasmids for replication and vectors for expression that contain the above nucleic acid fragments are also provided. Cells containing the plasmids and vectors are also provided. The cells can be any suitable host including, but are not limited to, bacterial cells, yeast cells, fungal cells, plant cells, insect cell and animal cells. The nucleic acids, plasmids, and cells containing the plasmids can be prepared according to methods known in the art including any described herein.

Also provided are methods for producing the above fusion proteins. An exemplary method includes the steps of growing, i.e. culturing the cells so that the proliferate, cells containing a plasmid encoding the fusion protein under conditions whereby the fusion protein is expressed by the cell, and recovering the expressed fusion protein. Methods for expressing and recovering recombinant proteins are well known in the art (*See generally, Current Protocols in Molecular Biology* (1998) § 16, John Wiley & Sons, Inc.) and such methods can be used for expressing and recovering the expressed fusion proteins. Preferably, the recombinant expression and recovery methods disclosed in Section B can be used.

The recovered fusion proteins can be isolated or purified by methods known in the art such as centrifugation, filtration, chromatograph, electrophoresis, immunoprecipitation, etc., or by a combination thereof (*See generally, Current Protocols in Molecular Biology* (1998) § 10, John Wiley & Sons, Inc.). Preferably, the recovered fusion protein is isolated or purified through affinity binding between the protein or peptide fragment of the fusion protein and an affinity binding moiety. As discussed in the above sections

-118-

regarding the construction of the fusion proteins, any affinity binding pairs can be constructed and used in the isolation or purification of the fusion proteins. For example, the affinity binding pairs can be protein binding sequences/protein, DNA binding sequences/DNA sequences, RNA binding sequences/RNA sequences, lipid binding sequences/lipid, polysaccharide binding sequences/polysaccharide, or metal binding sequences/metal.

5. Immobilization and supports or substrates therefor

In certain embodiments, where the targeting agents are designed for linkage to surfaces, the MTSP protein can be attached by linkage such as ionic or covalent, non-covalent or other chemical interaction, to a surface of a support or matrix material. Immobilization may be effected directly or via a linker. The MTSP protein may be immobilized on any suitable support, including, but are not limited to, silicon chips, and other supports described herein and known to those of skill in the art. A plurality of MTSP protein or protease domains thereof may be attached to a support, such as an array (*i.e.*, a pattern of two or more) of conjugates on the surface of a silicon chip or other chip for use in high throughput protocols and formats.

It is also noted that the domains of the MTSP protein can be linked directly to the surface or via a linker without a targeting agent linked thereto. Hence chips containing arrays of the domains of the MTSP protein.

The matrix material or solid supports contemplated herein are generally any of the insoluble materials known to those of skill in the art to immobilize ligands and other molecules, and are those that used in many chemical syntheses and separations. Such supports are used, for example, in affinity chromatography, in the immobilization of biologically active materials, and during chemical syntheses of biomolecules, including proteins, amino acids and other organic molecules and polymers. The preparation of and use of supports is well known to those of skill in this art; there are many such materials and preparations thereof known. For example, naturally-occurring support materials, such as agarose and cellulose, may be isolated from their respective sources, and processed according to known protocols, and synthetic materials may be prepared in accord with known protocols.

-119-

The supports are typically insoluble materials that are solid, porous, deformable, or hard, and have any required structure and geometry, including, but not limited to: beads, pellets, disks, capillaries, hollow fibers, needles, solid fibers, random shapes, thin films and membranes. Thus, the item may be
5 fabricated from the matrix material or combined with it, such as by coating all or part of the surface or impregnating particles.

Typically, when the matrix is particulate, the particles are at least about 10-2000 μ M, but may be smaller or larger, depending upon the selected application. Selection of the matrices will be governed, at least in part, by their
10 physical and chemical properties, such as solubility, functional groups, mechanical stability, surface area swelling propensity, hydrophobic or hydrophilic properties and intended use.

If necessary, the support matrix material can be treated to contain an appropriate reactive moiety. In some cases, the support matrix material already
15 containing the reactive moiety may be obtained commercially. The support matrix material containing the reactive moiety may thereby serve as the matrix support upon which molecules are linked. Materials containing reactive surface moieties such as amino silane linkages, hydroxyl linkages or carboxysilane linkages may be produced by well established surface chemistry techniques
20 involving silanization reactions, or the like. Examples of these materials are those having surface silicon oxide moieties, covalently linked to gamma-amino-propylsilane, and other organic moieties; N-[3-(triethoxysilyl)propyl]phthelamic acid; and bis-(2-hydroxyethyl)aminopropyltriethoxysilane. Exemplary of readily available materials containing amino group reactive functionalities, include, but
25 are not limited to, para-aminophenyltriethoxysilane. Also derivatized polystyrenes and other such polymers are well known and readily available to those of skill in this art (*e.g.*, the Tentagel® Resins are available with a multitude of functional groups, and are sold by Rapp Polymere, Tubingen, Germany; see, U.S. Patent No. 4,908,405 and U.S. Patent No. 5,292,814; *see, also* Butz et al.,
30 *Peptide Res.*, 7:20-23 (1994); and Kleine et al., *Immunobiol.*, 190:53-66 (1994)).

-120-

These matrix materials include any material that can act as a support matrix for attachment of the molecules of interest. Such materials are known to those of skill in this art, and include those that are used as a support matrix.

These materials include, but are not limited to, inorganics, natural polymers, and
5 synthetic polymers; including, but are not limited to: cellulose, cellulose derivatives, acrylic resins, glass, silica gels, polystyrene, gelatin, polyvinyl pyrrolidone, co-polymers of vinyl and acrylamide, polystyrene cross-linked with divinylbenzene and others (*see*, Merrifield, *Biochemistry*, 3:1385-1390 (1964)), polyacrylamides, latex gels, polystyrene, dextran, polyacrylamides, rubber,
10 silicon, plastics, nitrocellulose, celluloses, natural sponges. Of particular interest herein, are highly porous glasses (*see, e.g.*, U.S. Patent No. 4,244,721) and others prepared by mixing a borosilicate, alcohol and water.

Synthetic supports include, but are not limited to: acrylamides, dextran-derivatives and dextran co-polymers, agarose-polyacrylamide blends, other
15 polymers and co-polymers with various functional groups, methacrylate derivatives and co-polymers, polystyrene and polystyrene copolymers (*see, e.g.*, Merrifield, *Biochemistry*, 3:1385-1390 (1964); Berg et al., in *Innovation Perspect. Solid Phase Synth. Collect. Pap.*, Int. Symp., 1st, Epton, Roger (Ed), pp. 453-459 (1990); Berg et al., *Pept., Proc. Eur. Pept. Symp.*, 20th, Jung, G.
20 *et al.* (Eds), pp. 196-198 (1989); Berg et al., *J. Am. Chem. Soc.*, 111:8024-8026 (1989); Kent et al., *Isr. J. Chem.*, 17:243-247 (1979); Kent et al., *J. Org. Chem.*, 43:2845-2852 (1978); Mitchell et al., *Tetrahedron Lett.*, 42:3795-3798 (1976); U.S. Patent No. 4,507,230; U.S. Patent No. 4,006,117; and U.S. Patent No. 5,389,449). Such materials include those made from
25 polymers and co-polymers such as polyvinylalcohols, acrylates and acrylic acids such as polyethylene-co-acrylic acid, polyethylene-co-methacrylic acid, polyethylene-co-ethylacrylate, polyethylene-co-methyl acrylate, polypropylene-co-acrylic acid, polypropylene-co-methyl-acrylic acid, polypropylene-co-ethylacrylate, polypropylene-co-methyl acrylate, polyethylene-co-vinyl acetate, poly-
30 propylene-co-vinyl acetate, and those containing acid anhydride groups such as polyethylene-co-maleic anhydride and polypropylene-co-maleic anhydride.

-121-

Liposomes have also been used as solid supports for affinity purifications (Powell et al. *Biotechnol. Bioeng.*, 33:173 (1989)).

Numerous methods have been developed for the immobilization of proteins and other biomolecules onto solid or liquid supports (*see, e.g.,*

- 5 Mosbach, *Methods in Enzymology*, 44 (1976); Weetall, *Immobilized Enzymes, Antigens, Antibodies, and Peptides*, (1975); Kennedy et al., *Solid Phase Biochemistry, Analytical and Synthetic Aspects*, Scouten, ed., pp. 253-391 (1983); *see, generally*, Affinity Techniques. *Enzyme Purification: Part B. Methods in Enzymology*, Vol. 34, ed. W. B. Jakoby, M. Wilchek, Acad. Press, N.Y. (1974); and *Immobilized Biochemicals and Affinity Chromatography, Advances in Experimental Medicine and Biology*, vol. 42, ed. R. Dunlap, Plenum Press, N.Y. (1974)).

- Among the most commonly used methods are absorption and adsorption or covalent binding to the support, either directly or via a linker, such as the
- 15 numerous disulfide linkages, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups, known to those of skill in art (*see, e.g.,* the PIERCE CATALOG, ImmunoTechnology Catalog & Handbook, 1992-1993, which describes the preparation of and use of such reagents and provides a commercial source for
- 20 such reagents; Wong, *Chemistry of Protein Conjugation and Cross Linking*, CRC Press (1993); *see also* DeWitt et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90:6909 (1993); Zuckermann et al., *J. Am. Chem. Soc.*, 114:10646 (1992); Kurth et al., *J. Am. Chem. Soc.*, 116:2661 (1994); Ellman et al., *Proc. Natl. Acad. Sci. U.S.A.*, 91:4708 (1994); Sucholeiki, *Tetrahedron Lttrs.*, 35:7307 (1994); Su-
- 25 Sun Wang, *J. Org. Chem.*, 41:3258 (1976); Padwa et al., *J. Org. Chem.*, 41:3550 (1976); and Vedejs et al., *J. Org. Chem.*, 49:575 (1984), which describe photosensitive linkers).

- To effect immobilization, a composition containing the protein or other biomolecule is contacted with a support material such as alumina, carbon, an
- 30 ion-exchange resin, cellulose, glass or a ceramic. Fluorocarbon polymers have been used as supports to which biomolecules have been attached by adsorption

-122-

(see, U.S. Patent No. 3,843,443; Published International PCT Application WO/86 03840).

J. Prognosis and diagnosis

MTSP protein proteins, domains, analogs, and derivatives thereof, and
5 encoding nucleic acids (and sequences complementary thereto), and anti-MTSP
protein antibodies, can be used in diagnostics. Such molecules can be used in
assays, such as immunoassays, to detect, prognose, diagnose, or monitor
various conditions, diseases, and disorders affecting MTSP protein expression, or
monitor the treatment thereof. For purposes herein, the presence of MTSPs in
10 body fluids or tumor tissues are of particular interest.

In particular, such an immunoassay is carried out by a method including
contacting a sample derived from a patient with an anti-MTSP protein antibody
under conditions such that specific binding can occur, and detecting or
measuring the amount of any specific binding by the antibody. In a specific
15 aspect, such binding of antibody, in tissue sections, can be used to detect
aberrant MTSP protein localization or aberrant (*e.g.*, low or absent) levels of
MTSP protein. In a specific embodiment, antibody to MTSP protein can be used
to assay in a patient tissue or serum sample for the presence of MTSP protein
where an aberrant level of MTSP protein is an indication of a diseased condition.

20 The immunoassays which can be used include but are not limited to
competitive and non-competitive assay systems using techniques such as
western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent
assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin
reactions, gel diffusion precipitin reactions, immunodiffusion assays,
25 agglutination assays, complement-fixation assays, immunoradiometric assays,
fluorescent immunoassays, protein A immunoassays, to name but a few.

MTSP protein genes and related nucleic acid sequences and
subsequences, including complementary sequences, can also be used in
hybridization assays. MTSP protein nucleic acid sequences, or subsequences
30 thereof containing about at least 8 nucleotides, preferably 14 or 16 or 30 or
more contiguous nucleotides, can be used as hybridization probes.
Hybridization assays can be used to detect, prognose, diagnose, or monitor

-123-

conditions, disorders, or disease states associated with aberrant changes in MTSP protein expression and/or activity as described herein. In particular, such a hybridization assay is carried out by a method by contacting a sample containing nucleic acid with a nucleic acid probe capable of hybridizing to MTSP protein encoding DNA or RNA, under conditions such that hybridization can occur, and detecting or measuring any resulting hybridization.

- 5
- 10
- 15
- In a specific embodiment, a method of diagnosing a disease or disorder characterized by detecting an aberrant level of an MTSP protein in a subject is provided herein by measuring the level of the DNA, RNA, protein or functional activity of the epithelial MTSP protein at least partially encoded by a nucleic acid that hybridizes to a nucleic acid having the nucleotide sequence set forth in the SEQ. ID NO:1 in a sample derived from the subject, wherein an increase or decrease in the level of the DNA, RNA, protein or functional activity of the MTSP protein, relative to the level of the DNA, RNA, protein or functional activity found in an analogous sample not having the disease or disorder indicates the presence of the disease or disorder in the subject.

- Kits for diagnostic use are also provided, that contain in one or more containers an anti-MTSP protein antibody, particularly anti-MTSP3 or anti-MTSP4, and, optionally, a labeled binding partner to the antibody.
- 20
- A kit is also provided that includes in one or more containers a nucleic acid probe capable of hybridizing to MTSP protein-encoding RNA. In a specific embodiment, a kit can comprise in one or more containers a pair of primers (*e.g.*, each in the size range of 6-30 nucleotides) that are capable of priming amplification [*e.g.*, by polymerase chain reaction (see *e.g.*, Innis et al., 1990, PCR Protocols, Academic Press, Inc., San Diego, CA), ligase chain reaction (see EP 320,308) use of Q β replicase, cyclic probe reaction, or other methods known in the art under appropriate reaction conditions of at least a portion of an MTSP protein-encoding nucleic acid. A kit can optionally further comprise in a container a predetermined amount of a purified MTSP protein or nucleic acid, *e.g.*, for use as a standard or control.
- 25
- 30

-124-

K. PHARMACEUTICAL COMPOSITIONS AND MODES OF ADMINISTRATION**1. Components of the compositions**

Pharmaceutical compositions containing the identified compounds that modulate the activity of an MTSP protein are provided herein. Also provided are combinations of a compound that modulates the activity of an MTSP protein and another treatment or compound for treatment of a neoplastic disorder, such as a chemotherapeutic compound.

The MTSP protein modulator and the anti-tumor agent can be packaged as separate compositions for administration together or sequentially or intermittently. Alternatively, they can be provided as a single composition for administration or as two compositions for administration as a single composition. The combinations can be packaged as kits.

a. MTSP protein inhibitors

Any MTSP protein inhibitors, including those described herein when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with neoplastic diseases, including undesired and/or uncontrolled angiogenesis, can be used in the present combinations.

In one embodiment, the MTSP protein inhibitor is an antibody or fragment thereof that specifically reacts with an MTSP protein or the protease domain thereof, an inhibitor of the MTSP protein production, an inhibitor of the epithelial MTSP protein membrane-localization, or any inhibitor of the expression of or, especially, the activity of an MTSP protein.

25 b. Anti-angiogenic agents and anti-tumor agents

Any anti-angiogenic agents and anti-tumor agents, including those described herein, when used alone or in combination with other compounds, that can alleviate, reduce, ameliorate, prevent, or place or maintain in a state of remission of clinical symptoms or diagnostic markers associated with undesired and/or uncontrolled angiogenesis and/or tumor growth and metastasis, particularly solid neoplasms, vascular malformations and cardiovascular disorders, chronic inflammatory diseases and aberrant wound repairs, circulatory

-125-

disorders, crest syndromes, dermatological disorders, or ocular disorders, can be used in the combinations. Also contemplated are anti-tumor agents for use in combination with an inhibitor of an MTSP protein.

c. Anti-tumor agents and anti-angiogenic agents

- 5 The compounds identified by the methods provided herein or provided herein can be used in combination with anti-tumor agents and/or anti-angiogenesis agents.

2. Formulations and route of administration

- 10 The compounds herein and agents are preferably formulated as pharmaceutical compositions, preferably for single dosage administration. The concentrations of the compounds in the formulations are effective for delivery of an amount, upon administration, that is effective for the intended treatment. Typically, the compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of a compound or mixture thereof is
- 15 dissolved, suspended, dispersed or otherwise mixed in a selected vehicle at an effective concentration such that the treated condition is relieved or ameliorated. Pharmaceutical carriers or vehicles suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration.
- 20 In addition, the compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients. Liposomal suspensions, including tissue-targeted liposomes, may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art. For
- 25 example, liposome formulations may be prepared as described in U.S. Patent No. 4,522,811.

- The active compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. The therapeutically
- 30 effective concentration may be determined empirically by testing the compounds in known in vitro and in vivo systems, such as the assays provided herein.

-126-

The concentration of active compound in the drug composition will depend on absorption, inactivation and excretion rates of the active compound, the physicochemical characteristics of the compound, the dosage schedule, and amount administered as well as other factors known to those of skill in the art.

5 Typically a therapeutically effective dosage is contemplated. The amounts administered may be on the order of 0.001 to 1 mg/ml, preferably about 0.005-0.05 mg/ml, more preferably about 0.01 mg/ml, of blood volume. Pharmaceutical dosage unit forms are prepared to provide from about 1 mg to about 1000 mg and preferably from about 10 to about 500 mg, more preferably
10 about 25-75 mg of the essential active ingredient or a combination of essential ingredients per dosage unit form. The precise dosage can be empirically determined.

The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is
15 understood that the precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by extrapolation from in vivo or in vitro test data. It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular
20 subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or use of the claimed compositions and combinations containing them.

25 Preferred pharmaceutically acceptable derivatives include acids, salts, esters, hydrates, solvates and prodrug forms. The derivative is typically selected such that its pharmacokinetic properties are superior to the corresponding neutral compound.

-127-

Thus, effective concentrations or amounts of one or more of the compounds provided herein or pharmaceutically acceptable derivatives thereof are mixed with a suitable pharmaceutical carrier or vehicle for systemic, topical or local administration to form pharmaceutical compositions. Compounds are included in an amount effective for ameliorating or treating the disorder for which treatment is contemplated. The concentration of active compound in the composition will depend on absorption, inactivation, excretion rates of the active compound, the dosage schedule, amount administered, particular formulation as well as other factors known to those of skill in the art.

- 5 Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers, such as acetates, citrates and phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. Parenteral preparations can be enclosed in ampules, disposable syringes or single or multiple dose vials made of glass, plastic or other suitable material.
- 10 In instances in which the compounds exhibit insufficient solubility, methods for solubilizing compounds may be used. Such methods are known to those of skill in this art, and include, but are not limited to, using cosolvents, such as dimethylsulfoxide (DMSO), using surfactants, such as Tween®, or dissolution in aqueous sodium bicarbonate. Derivatives of the compounds, such as prodrugs of the compounds may also be used in formulating effective pharmaceutical compositions. For ophthalmic indications, the compositions are formulated in an ophthalmically acceptable carrier. For the ophthalmic uses herein, local administration, either by topical administration or by injection is preferred. Time release formulations are also desirable. Typically, the compositions are formulated for single dosage administration, so that a single dose administers an effective amount.

-128-

Upon mixing or addition of the compound with the vehicle, the resulting mixture may be a solution, suspension, emulsion or other composition. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the compound in the selected carrier or vehicle. If necessary, pharmaceutically acceptable salts or other derivatives of the compounds are prepared.

The compound is included in the pharmaceutically acceptable carrier in an amount sufficient to exert a therapeutically useful effect in the absence of undesirable side effects on the patient treated. It is understood that number and degree of side effects depends upon the condition for which the compounds are administered. For example, certain toxic and undesirable side effects are tolerated when treating life-threatening illnesses that would not be tolerated when treating disorders of lesser consequence.

The compounds can also be mixed with other active materials, that do not impair the desired action, or with materials that supplement the desired action known to those of skill in the art. The formulations of the compounds and agents for use herein include those suitable for oral, rectal, topical, inhalational, buccal (*e.g.*, sublingual), parenteral (*e.g.*, subcutaneous, intramuscular, intradermal, or intravenous), transdermal administration or any route. The most suitable route in any given case will depend on the nature and severity of the condition being treated and on the nature of the particular active compound which is being used. The formulations are provided for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, and oral solutions or suspensions, and oil-water emulsions containing suitable quantities of the compounds or pharmaceutically acceptable derivatives thereof. The pharmaceutically therapeutically active compounds and derivatives thereof are typically formulated and administered in unit-dosage forms or multiple-dosage forms. Unit-dose forms as used herein refers to physically discrete units suitable for human and animal subjects and packaged individually as is known in the art. Each unit-dose contains a predetermined quantity of the therapeutically active compound sufficient to produce the desired therapeutic effect, in association

-129-

with the required pharmaceutical carrier, vehicle or diluent. Examples of unit-dose forms include ampoules and syringes and individually packaged tablets or capsules. Unit-dose forms may be administered in fractions or multiples thereof. A multiple-dose form is a plurality of identical unit-dosage forms

- 5 packaged in a single container to be administered in segregated unit-dose form. Examples of multiple-dose forms include vials, bottles of tablets or capsules or bottles of pints or gallons. Hence, multiple dose form is a multiple of unit-doses which are not segregated in packaging.

- The composition can contain along with the active ingredient: a diluent
10 such as lactose, sucrose, dicalcium phosphate, or carboxymethylcellulose; a lubricant, such as magnesium stearate, calcium stearate and talc; and a binder such as starch, natural gums, such as gum acaciagelatin, glucose, molasses, polvinylpyrrolidine, celluloses and derivatives thereof, povidone, crospovidones and other such binders known to those of skill in the art. Liquid
15 pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be
20 administered may also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, or solubilizing agents, pH buffering agents and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such
25 dosage forms are known, or will be apparent, to those skilled in this art (see, *e.g.*, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 15th Edition, 1975). The composition or formulation to be administered will contain a quantity of the active compound in an amount sufficient to alleviate the symptoms of the treated subject.
- 30 Dosage forms or compositions containing active ingredient in the range of 0.005% to 100% with the balance made up from non-toxic carrier may be prepared. For oral administration, the pharmaceutical compositions may take the

-130-

form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.*, pregelatinized maize starch, polyvinyl pyrrolidone or hydroxypropyl methylcellulose); fillers (*e.g.*, lactose, microcrystalline cellulose or calcium 5 hydrogen phosphate); lubricants (*e.g.*, magnesium stearate, talc or silica); disintegrants (*e.g.*, potato starch or sodium starch glycolate); or wetting agents (*e.g.*, sodium lauryl sulphate). The tablets may be coated by methods well-known in the art.

The pharmaceutical preparation may also be in liquid form, for example, 10 solutions, syrups or suspensions, or may be presented as a drug product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.*, sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (*e.g.*, lecithin or 15 acacia); non-aqueous vehicles (*e.g.*, almond oil, oily esters, or fractionated vegetable oils); and preservatives (*e.g.*, methyl or propyl-p-hydroxybenzoates or sorbic acid).

Formulations suitable for rectal administration are preferably presented as unit dose suppositories. These may be prepared by admixing the active 20 compound with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

Formulations suitable for topical application to the skin or to the eye preferably take the form of an ointment, cream, lotion, paste, gel, spray, aerosol and oil. Carriers which may be used include vaseline, lanoline, polyethylene 25 glycols, alcohols, and combinations of two or more thereof. The topical formulations may further advantageously contain 0.05 to 15 percent by weight of thickeners selected from among hydroxypropyl methyl cellulose, methyl cellulose, polyvinylpyrrolidone, polyvinyl alcohol, poly (alkylene glycols), poly(hydroxyalkyl), (meth)acrylates or poly(meth)acrylamides. A topical 30 formulation is often applied by instillation or as an ointment into the conjunctival sac. It can also be used for irrigation or lubrication of the eye, facial sinuses, and external auditory meatus. It may also be injected into the anterior eye

-131-

chamber and other places. The topical formulations in the liquid state may be also present in a hydrophilic three-dimensional polymer matrix in the form of a strip, contact lens, and the like from which the active components are released.

- For administration by inhalation, the compounds for use herein can be
- 5 delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and
- 10 cartridges of, *e.g.*, gelatin, for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

- Formulations suitable for buccal (sublingual) administration include, for example, lozenges containing the active compound in a flavored base, usually
- 15 sucrose and acacia or tragacanth; and pastilles containing the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

- The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or in multi-dose
- 20 containers, with an added preservative. The compositions may be suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water or other solvents, before use.

- 25 Formulations suitable for transdermal administration may be presented as discrete patches adapted to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. Such patches suitably contain the active compound as an optionally buffered aqueous solution of, for example, 0.1 to 0.2 M concentration with respect to the active compound. Formulations
- 30 suitable for transdermal administration may also be delivered by iontophoresis (*see, e.g.*, Pharmaceutical Research 3 (6), 318 (1986)) and typically take the form of an optionally buffered aqueous solution of the active compound.

-132-

The pharmaceutical compositions may also be administered by controlled release means and/or delivery devices (see, *e.g.*, in U.S. Patent Nos. 3,536,809; 3,598,123; 3,630,200; 3,845,770; 3,847,770; 3,916,899; 4,008,719; 4,687,610; 4,769,027; 5,059,595; 5,073,543; 5,120,548; 5,354,566; 5 5,591,767; 5,639,476; 5,674,533 and 5,733,566).

Desirable blood levels may be maintained by a continuous infusion of the active agent as ascertained by plasma levels. It should be noted that the attending physician would know how to and when to terminate, interrupt or adjust therapy to lower dosage due to toxicity, or bone marrow, liver or kidney 10 dysfunctions. Conversely, the attending physician would also know how to and when to adjust treatment to higher levels if the clinical response is not adequate (precluding toxic side effects).

The efficacy and/or toxicity of the MTSP protein inhibitor(s), alone or in combination with other agents can also be assessed by the methods known in 15 the art (See generally, O'Reilly, *Investigational New Drugs*, 15:5-13 (1997)).

The active compounds or pharmaceutically acceptable derivatives may be prepared with carriers that protect the compound against rapid elimination from the body, such as time release formulations or coatings.

Kits containing the compositions and/or the combinations with 20 instructions for administration thereof are provided. The kit may further include a needle or syringe, preferably packaged in sterile form, for injecting the complex, and/or a packaged alcohol pad. Instructions are optionally included for administration of the active agent by a clinician or by the patient.

Finally, the compounds or MTSP proteins or protease domains thereof or 25 compositions containing any of the preceding agents may be packaged as articles of manufacture containing packaging material, a compound or suitable derivative thereof provided herein, which is effective for treatment of a diseases or disorders contemplated herein, within the packaging material, and a label that indicates that the compound or a suitable derivative thereof is for treating the 30 diseases or disorders contemplated herein. The label can optionally include the disorders for which the therapy is warranted.

-133-

L. METHODS OF TREATMENT

The compounds identified by the methods herein are used for treating or preventing neoplastic diseases in an animal, particularly a mammal, including a human, is provided herein. In one embodiment, the method includes

- 5 administering to a mammal an effective amount of an inhibitor of an MTSP protein, whereby the disease or disorder is treated or prevented. In a preferred embodiment, the MTSP protein inhibitor used in the treatment or prevention is administered with a pharmaceutically acceptable carrier or excipient. The mammal treated can be a human.
- 10 The inhibitors provided herein are those identified by the screening assays. In addition, antibodies and antisense nucleic acids are contemplated.

- The treatment or prevention method can further include administering an anti-angiogenic treatment or agent or anti-tumor agent simultaneously with, prior to or subsequent to the MTSP protein inhibitor, which can be any compound
- 15 identified that inhibits the activity of an MTSP protein, and includes an antibody or a fragment or derivative thereof containing the binding region thereof against the MTSP protein, an antisense nucleic acid encoding the MTSP protein, and a nucleic acid containing at least a portion of a gene encoding the MTSP protein into which a heterologous nucleotide sequence has been inserted such that the
- 20 heterologous sequence inactivates the biological activity of at least a portion of the gene encoding the MTSP protein, in which the portion of the gene encoding the MTSP protein flanks the heterologous sequence so as to promote homologous recombination with a genomic gene encoding the MTSP protein.

1. Antisense treatment

- 25 In a specific embodiment, as described hereinabove, MTSP protein function is reduced or inhibited by MTSP protein antisense nucleic acids, to treat or prevent neoplastic disease. The therapeutic or prophylactic use of nucleic acids of at least six nucleotides that are antisense to a gene or cDNA encoding MTSP protein or a portion thereof. An MTSP protein "antisense" nucleic acid as
- 30 used herein refers to a nucleic acid capable of hybridizing to a portion of an MTSP protein RNA (preferably mRNA) by virtue of some sequence complementarity. The antisense nucleic acid may be complementary to a coding

-134-

and/or noncoding region of an MTSP protein mRNA. Such antisense nucleic acids have utility as therapeutics that reduce or inhibit MTSP protein function, and can be used in the treatment or prevention of disorders as described *supra*.

- The MTSP protein antisense nucleic acids are of at least six nucleotides
5 and are preferably oligonucleotides (ranging from 6 to about 150 nucleotides, or more preferably 6 to 50 nucleotides). In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or at least 125 nucleotides. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-
10 stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger et al., *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556 (1989); Lemaitre et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652 (1987);
15 PCT Publication No. WO 88/09810, published December 15, 1988) or blood-brain barrier (see, *e.g.*, PCT Publication No. WO 89/10134, published April 25, 1988), hybridization-triggered cleavage agents (see, *e.g.*, Krol et al., *BioTechniques* 6:958-976 (1988)) or intercalating agents (see, *e.g.*, Zon, *Pharm. Res.* 5:539-549 (1988)).
- 20 The MTSP protein antisense nucleic acid is preferably an oligonucleotide, more preferably of single-stranded DNA. In a preferred aspect, the oligonucleotide includes a sequence antisense to a portion of human MTSP protein. The oligonucleotide may be modified at any position on its structure with substituents generally known in the art.
- 25 The MTSP protein antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil,
30 dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine,

-135-

- 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil,
- 5 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

- In another embodiment, the oligonucleotide includes at least one modified sugar moiety selected from the group including but not limited to arabinose,
- 10 2-fluoroarabinose, xylulose, and hexose. The oligonucleotide can include at least one modified phosphate backbone selected from a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.
- 15 The oligonucleotide can be an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which the strands run parallel to each other (Gautier et al., *Nucl. Acids Res.* 15:6625-6641 (1987)).

- The oligonucleotide may be conjugated to another molecule, *e.g.*, a
- 20 peptide, hybridization triggered cross-linking agent, transport agent and hybridization-triggered cleavage agent.

- The oligonucleotides may be synthesized by standard methods known in the art, *e.g.* by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples,
- 25 phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (*Nucl. Acids Res.* 16:3209 (1988)), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., *Proc. Natl. Acad. Sci. U.S.A.* 85:7448-7451 (1988)), etc.

- In a specific embodiment, the MTSP protein antisense oligonucleotide
- 30 includes catalytic RNA, or a ribozyme (see, *e.g.*, PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al., *Science* 247:1222-1225 (1990)). In another embodiment, the oligonucleotide is a 2'-O-

-136-

methylribonucleotide (Inoue et al., *Nucl. Acids Res.* 15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., *FEBS Lett.* 215:327-330 (1987)).

- In an alternative embodiment, the MTSP protein antisense nucleic acid is produced intracellularly by transcription from an exogenous sequence. For
- 5 example, a vector can be introduced *in vivo* such that it is taken up by a cell, within which cell the vector or a portion thereof is transcribed, producing an antisense nucleic acid (RNA). Such a vector would contain a sequence encoding the MTSP protein antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce
- 10 the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the MTSP protein antisense RNA can be by any promoter known in the art to act in mammalian, preferably human, cells.
- 15 Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, *Nature* 290:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell* 22:787-797 (1980), the herpes thymidine kinase promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature* 296:39-42 (1982), etc.
- 20

The antisense nucleic acids include sequence complementary to at least a portion of an RNA transcript of an MTSP protein gene, preferably a human MTSP protein gene. Absolute complementarity, although preferred, is not required.

- 25 The amount of MTSP protein antisense nucleic acid that will be effective in the treatment or prevention of neoplastic disease will depend on the nature of the disease, and can be determined empirically by standard clinical techniques. Where possible, it is desirable to determine the antisense cytotoxicity in cells *in vitro*, and then in useful animal model systems prior to testing and use in
- 30 humans.

-137-

2. Gene Therapy

- In an exemplary embodiment, nucleic acids that include a sequence of nucleotides encoding an MTSP protein or functional domains or derivative thereof, are administered to promote MTSP protein function, by way of gene therapy. Gene therapy refers to therapy performed by the administration of a nucleic acid to a subject. In this embodiment, the nucleic acid produces its encoded protein that mediates a therapeutic effect by promoting MTSP protein function. Any of the methods for gene therapy available in the art can be used (see, Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *An. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *An. Rev. Biochem.* 62:191-217 (1993); *TIBTECH* 11(5):155-215 (1993). For example, one therapeutic composition for gene therapy includes an MTSP protein-encoding nucleic acid that is part of an expression vector that expresses an MTSP protein or domain, fragment or chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the MTSP protein coding region, the promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the MTSP protein coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the MTSP protein nucleic acid (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).
- Delivery of the nucleic acid into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.
- In a specific embodiment, the nucleic acid is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by

-138-

constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by infection using a defective or attenuated retroviral or other viral vector (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see *e.g.*, Wu and Wu, *J. Biol. Chem.* 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand is a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, *e.g.*, PCT Publications WO 92/06180 dated April 16, 1992 (Wu et al.); WO 92/22635 dated December 23, 1992 (Wilson et al.); WO92/20316 dated November 26, 1992 (Findeis et al.); WO93/14188 dated July 22, 1993 (Clarke et al.), WO 93/20221 dated October 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989)).

In a specific embodiment, a viral vector that contains the MTSP protein nucleic acid is used. For example, a retroviral vector can be used (see Miller et al., *Meth. Enzymol.* 217:581-599 (1993)). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The *MTSP protein* nucleic acid to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references

-139-

illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

- 5 Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the
- 10 advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of
- 15 adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); and Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993).

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., *Proc. Soc. Exp. Biol. Med.* 204:289-300 (1993)).

- 20 Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are
- 25 expressing the transferred gene. Those cells are then delivered to a patient.

- In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or
- 30 bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the

-140-

introduction of foreign genes into cells (see *e.g.*, Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92 (1985)) and may be used, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. In a preferred embodiment, epithelial cells are injected, *e.g.*, subcutaneously. In another embodiment, recombinant skin cells may be applied as a skin graft onto the patient. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient. In an embodiment in which recombinant cells are used in gene therapy, an MTSP protein nucleic acid is introduced into the cells such that it is expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment. Such stem cells include but are not limited to hematopoietic stem cells (HSC), stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (PCT Publication WO

-141-

94/08598, dated April 28, 1994), and neural stem cells (Stemple and Anderson, *Cell* 71:973-985 (1992)).

- Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures (Rheinwald, *Meth. Cell Bio.* 21A:229 (1980)). In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture
- 10 (Rheinwald, *Meth. Cell Bio.* 21A:229 (1980); Pittelkow and Scott, *Mayo Clinic Proc.* 61:771 (1986)). If the ESCs are provided by a donor, a method for suppression of host versus graft reactivity (*e.g.*, irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used.

- With respect to hematopoietic stem cells (HSC), any technique which
- 15 provides for the isolation, propagation, and maintenance *in vitro* of HSC can be used in this embodiment. Techniques by which this may be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which may be allogeneic or xenogeneic.
- 20 Non-autologous HSC are used preferably in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a particular embodiment, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, *e.g.*, Kodo et al., *J. Clin. Invest.* 73:1377-1384 (1984)). In a preferred embodiment, the HSCs can be made
- 25 highly enriched or in substantially pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques known in the art. Long-term cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., *J. Cell Physiol.* 91:335 (1977) or Witlock-Witte culture
- 30 techniques (Witlock and Witte, *Proc. Natl. Acad. Sci. USA* 79:3608-3612 (1982)).

-142-

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy includes an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5 **3. Prodrugs**

A method for treating tumors is provided. The method is practiced by administering a prodrug that is specifically cleaved by an MTSP to release an active drug. Upon contact with a cell that expresses MTSP activity, the prodrug is converted into an active drug. The prodrug can be a conjugate that contains
10 the active agent, such as an anti-tumor drug, such as a cytotoxic agent, or other atherapeutic agent, linked, linked to a substrate for the targeted MTSP, such that the drug or agent is inactive or unable to enter a cell, in the conjugate, but is activated upon cleavage. The prodrug, for example, can contain an oligopeptide, preferably a relatively short, less than about 10 amino acids peptide, that is
15 selectively proteolytically cleaved by the targeted MTSP. Cytotoxic agents, include, but are not limited to, alkylating agents, antiproliferative agents and tubulin binding agents. Others include, vinca drugs, mitomycins, bleomycins and taxanes.

M. ANIMAL MODELS

20 Transgenic animal models are provided herein. Such an animal can be initially produced by promoting homologous recombination between an MTSP protein gene in its chromosome and an exogenous MTSP protein gene that has been rendered biologically inactive (preferably by insertion of a heterologous sequence, *e.g.*, an antibiotic resistance gene). In a preferred aspect, this
25 homologous recombination is carried out by transforming embryo-derived stem (ES) cells with a vector containing the insertionally inactivated MTSP protein gene, such that homologous recombination occurs, followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of the chimeric animal ("knockout animal") in which an
30 MTSP protein gene has been inactivated (see Cápecchi, *Science* 244:1288-1292 (1989)). The chimeric animal can be bred to produce additional knockout animals. Such animals can be mice, hamsters, sheep, pigs, cattle, etc., and are

-143-

preferably non-human mammals. In a specific embodiment, a knockout mouse is produced.

Such knockout animals are expected to develop or be predisposed to developing neoplastic diseases and thus can have use as animal models of such diseases *e.g.*, to screen for or test molecules for the ability to treat or prevent such diseases or disorders. Hence, the animal models for are provided. Such an animal can be initially produced by promoting homologous recombination between an MTSP gene in its chromosome and an exogenous MTSP protein gene that would be over-expressed or mis-expressed (preferably by expression under a strong promoter). In a preferred aspect, this homologous recombination is carried out by transforming embryo-derived stem (ES) cells with a vector containing the over-expressed or mis-expressed MTSP protein gene, such that homologous recombination occurs, followed by injecting the ES cells into a blastocyst, and implanting the blastocyst into a foster mother, followed by the birth of the chimeric animal in which an MTSP gene has been over-expressed or mis-expressed (see Capecchi, *Science* 244:1288-1292 (1989)). The chimeric animal can be bred to produce additional animals with over-expressed or mis-expressed MTSP protein. Such animals can be mice, hamsters, sheep, pigs, cattle, etc., and are preferably non-human mammals. In a specific embodiment, a mouse with over-expressed or mis-expressed MTSP protein is produced.

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

25

EXAMPLE 1**Cloning of MTSP3, cloning and mutagenesis of the Protease domain of MTSP3****1. Identification and cloning of MTSP3****a. Identification of EST clones AI924527 and AI924182 as part of a serine protease MTSP3**

30

DNA encoding the protease domain of the protease designated MTSP1 was independently cloned from the human prostatic adenocarcinoma cell line, PC-3, using degenerate oligonucleotide primers, then sequenced and

-144-

characterized (see EXAMPLE 6). The sequence of the sense degenerate primer used in cloning MTSP1 was 5'-TGGRT(I)VT(I)WS(I)GC(I)RC(I)CAYTG-3' (SEQ ID No: 13), and that of the anti-sense was

5'-(I)GG(I)CC(I)CC(I)SWRTC(I)CCYT(I)RCA(I)GHRTC-3' (SEQ ID No:14), where

- 5 R = A,G; V = G,A,C; W = A,T; S = G,C; Y = C,T; H = A,T,C. The primer sequences correspond to two highly conserved regions in all serine proteases and should amplify PCR products ranging from 400 to 500 base pairs. MTSP1 was subsequently found to be identical to matriptase (Genbank accession number AF118224; see also Takeuchi et al., *Proc. Natl. Acad. Sci. USA*,
10 96(20):11054-61 (1999); and Lin et al., *J. Biol. Chem.*, 274(26):18231-6 (1999).

- Using the protein sequence of the protease domain of the serine protease MTSP1, the EST database (dbEST) at the National Center for Biotechnology Information (Bethesda, MD; www.ncbi.nlm.nih.gov) was searched for EST
15 clones that contain similar or identical sequences to MTSP1 using the search algorithm tblastn. The tblastn algorithm compares a protein query sequence against a nucleotide sequence database dynamically translated in all six reading frames (both strands). The sequences for two identical EST clones (NCI_CGAP_Lu19 AI924527 and AI924182) derived from human lung tumor
20 tissue showed 43% identity with the MTSP1 protein sequence. Subsequent search of GenBank and SwissProt database for the EST sequence AI924527 and AI924182 did not show any matching sequence to MTSP1, indicating that the sequence contained in these EST clones AI924527 and AI924182 may be portions of a new serine protease.

- 25 b. **PCR cloning of a cDNA fragment of another membrane type serine protease MTSP3**

- The double-stranded Marathon-Ready(tm) cDNA library derived from human lung carcinoma (LX-1) was obtained from Clontech (Palo Alto, CA; catalog # 7495-1) and used as a template. Two primers,
30 5'-TCACCGAGAAGATGATGTGTGTCAGGCATCC-3' (SEQ ID No:15) (sense primer), and 5'-GGGACAGGGGCTGTAAGGCAGGGAATGAG-3' (SEQ ID No:16) (antisense primer), were used to amplify a ~360 bp DNA fragment. The PCR

-145-

product was separated on a 2% agarose gel and purified using a gel extraction kit (catalog number 28706; QIAquick gel extraction kit; Qiagen). The purified DNA fragment was ligated into TA vectors (catalog number K4500-01; TOPO-TA cloning kit, Invitrogen, Carlsbad, CA). After transformation into *E. coli* cells, plasmids were isolated and analyzed by digestion with EcoRI restriction enzyme. Clones that had inserted DNA were further characterized by sequencing using a fluorescent dye-based DNA sequencing method (catalog number 4303149; BigDye terminator cycle sequencing kit with AmpliTaq DNA polymerase; Perkin Elmer, Lincoln, CA).

- 10 The DNA sequence obtained was analyzed and has 43% identity with the MTSP1 protein sequence. This indicates that the LX-1 cDNA library contains a desired nucleic acid molecule. It was used to isolate a cDNA clone encompassing a full length protease.

c. 5'- and 3'- rapid amplification of cDNA ends (RACE)

- 15 To obtain the full-length cDNA that encoded this serine protease, hereafter called MTSP3, 5'- and 3'-RACE reactions were performed. The Marathon-Ready cDNA library from human lung carcinoma (LX-1) was used to isolate the 5' and 3' ends of the cDNA encoding MTSP3. Marathon-Ready cDNA is specifically made for RACE reactions. Two gene specific primers were used: 5'-CCCGCAGCCATAGCCCCAGCTAACG-3' (SEQ ID No. 17) for 5'-RACE reaction and 5'-GCAGACGATGCGTACCAGGGGAAGTC-3' (SEQ ID No. 18) for 3'-RACE reaction. Two fragments, approximately 1.8 kbp and 0.85 kbp, were isolated that correspond to the missing 5' and 3' end sequences, respectively. These fragments were subcloned as described above. They were further confirmed by Southern analysis using an internal cDNA fragment encompassing the 2 primers used in the RACE reactions as probe and by DNA sequence analysis.

d. PCR amplification of cDNA encoding full-length protease domain of MTSP3

- 30 To obtain the cDNA fragment encoding the protease domain of MTSP3, an end-to-end PCR amplification using gene-specific primers was used. The two primers used were:

-146-

5'-CTCGAGAAAAGAGTGGTGGGTGGGGAGGAGGCCTCTGTG-3' (SEQ ID No. 19) for the 5' end and 5'-GCGGCCGCATTACAGCTCAGCCTTCCAGAC-3' (SEQ ID No. 20) for the 3' end. The 5' primer contains the sequence (underlined) that encodes the start of the MTSP3 protease domain (VVGGEASV). The 3' primer contains the stop codon (underlined) of MTSP3. A ~700-bp fragment was amplified and subcloned into a *Pichia pastoris* expression vector, pPIC9K.

e. C310S mutagenesis of MTSP3

To eliminate the free cysteine (at position 310 in SEQ ID No. 4) that exists when the protease domain of the MTSP3 protein is expressed or the zymogen is activated, the free cysteine at position 310 (see SEQ ID No. 3), which is Cys122 if a chymotrypsin numbering scheme is used, was replaced with a serine. The resulting vector was designated pPIC9K:MTSP3C122S.

The gene encoding the protease domain of MTSP3 was mutagenized by PCR SOE (PCR-based splicing by overlap extension) to replace the unpaired cysteine at position 310 (122 chymotrypsin numbering system) with a serine. Two overlapping gene fragments, each containing the TCT codon for serine at position 310 were PCR amplified using the following primers: for the 5' gene fragment, TCTCTCGAGAAAAGAGTGGTGGGTGGGTGGGGAGGAGGCCTCTGTG SEQ ID No. 51 and GCTCCTCATCAAAGAAGGGCAGAGAGATGGGCCTGACTGTGCC SEQ ID No. 52; for the 3' gene fragment, ATTCGCGGCCGCATTACAGCTCAGCCTTCCAGAC (SEQ ID No. 53) and GGCACAGTCAGGCCCATCTCTCTGCCCTTCTTTGATGAGGAGC (SEQ ID No. 54). The amplified gene fragments were purified on a 1% agarose gel, mixed and reamplified by PCR to produce the full length coding sequence for MTSP3 C122S. This sequence was then cut with restriction enzymes NotI and XhoI, and ligated into vector pPic9K.

2. Sequence analysis

All derived DNA and protein sequences were analyzed using MacVector (version 6.5; Oxford Molecular Ltd., Madison, WI). The full-length cDNA encoding MTSP3 is composed of 2,137 base pairs containing the longest open reading frame of 1,314 base pairs which translate to a 437-amino acid protein

-147-

sequence. The cDNA fragment (nt 873-1,574) encoding the protease domain of MTSP3 is composed of 702 base pairs which translate to a 233-amino acid protein sequence plus the stop codon. The DNA sequence and the translated protein sequence of MTSP3 are shown in SEQ ID Nos. 3 and 4, respectively.

3. Construction of the Expression Vectors

DNA encoding MTSP3 full length protein containing the C310S point mutation (i.e., MTSP3C122S) was cloned from pPIC9K:MTSP3C122S. The primers MTSP3:

5' GAATTCCATATGCCGCGCTTTAAAGTGGTGGGTGGGGAGGAGGCC SEQ ID

No. 47 (containing a NdeI restriction site) and MTSP3-

3' GGGATACCCGTTACAGCTCAGCCTTCCAGAC 5' SEQ ID No. 48 (containing a BamHI restriction site) were used to PCR amplify the human MTSP3C122S

protease domain utilizing a plasmin recognition sequence (PRFK) for zymogen activation. Amplification was conducted in a total volume of 50 μ l containing 10

mM KCl, 20 mM Tris-HCl (pH 8.8 at 25 °C), 10 mM $(\text{NH}_4)_2\text{SO}_4$, 2.0 mM MgSO_4 , 0.1% Triton X-100, 0.3 mM dNTPs, 5.0 units of vent DNA polymerase, and 100 pmol of primers. The reaction mixtures were heated to 95 °C for 5 min, followed by 25–30 cycles of 95, 60, and 75 °C for 30 s each and a final extension at 75 °C for 2 min.

PCR products were purified using a QIAquick PCR purification kit (QIAGEN Inc., Chatsworth, CA). Full-length oligonucleotides were doubly digested with 10 units BamHI and 20 units NdeI for 2 h at 37°C. The digested fragments were purified on a 1.3% agarose gel and stained with ethidium bromide. The band containing the MTSP3C122S encoding DNA was excised and purified using a QIAEX II gel extraction kit.

The MTSP3C122S encoding DNA was then cloned into the NdeI and BamHI sites of the pET19b vector (Novagen) using standard methods. This vector allows the fusion of a HIS_6 tag for purification by metal affinity chromatography (MAC). Competent XL1 Blue cells (Stratagene) were transformed with the pET19b-MTSP3C122S vector and used to produce plasmid stocks. Proper insertion and DNA sequence were confirmed by fluorescent thermal dye DNA sequencing methods as well as restriction digests.

-148-

4. Protein Expression, Purification, and Refolding

Overexpression of the gene product was achieved in *E. coli* strain BL21(DE3) (Novagen, Madison WI) containing the DNAY plasmid for rare codon optimization (see, *e.g.*, Garcia *et al.* (1986) *Cell* 45:453-459). Cells were grown at 37 °C in (2xYT) media supplemented with carbenicillin and kanamycin to a final concentrations of 50 ug/ml and 34 ug/ml, respectively. One liter cultures were inoculated with 10 mL of an overnight culture grown in the same media. Cells were allowed to grow to a density of 0.6 – 1.0 OD₆₀₀ before the addition of IPTG (final concentration 1.0 mM). Cells were grown an additional 4 hours before harvesting.

The cell pellet was resuspended in 20 mL of lysis buffer (50 mM Na₂HPO₄, 300 mM NaCl, pH 7.4). The cell suspension was treated with 10-20 mg lysozyme and incubated at 37°C for 1 hour. DNaseI was then added (1-2mg) with mixing until the solution was no longer viscous. The solution was then transferred to a Rosette flask and sonicated, on ice, at high power for 15 min. Inclusion bodies were pelleted by centrifugation at 20K rpm (~48,000 g) at 4°C for 30 min.

Inclusion bodies were washed by douncing 2 times in 50 mM Na₂HPO₄, 300 mM NaCl, 5% LADO, pH 7.4 followed by 2 times in 50 mM Na₂HPO₄, 300 mM NaCl, pH 7.4. Inclusion bodies (~ 500 mg) are solubilized in 25 mL 6 M GuHCl, 100 mM tris-HCl, 20 mM βMe, pH 8.0. This solution was spun at 20K rpm for 30 minutes to pull down any particulate matter. This solution was passed through a 0.2 μM filter and diluted to 100 mL in solubilization buffer.

MTSP3C122S was refolded by slowly adding the inclusion body mixture to 8 L of refolding buffer (100 mM tris-HCl, 150 mM NaCl, 5 mM GSH, 0.05 mM GSSG, 1 M arginine, pH 8.0) using a peristaltic pump. The refolding mixture was allowed to stir at 4°C for 7 days or until the thiol concentration was below 1 mM as detected by Ellman's reagent. The solution was filtered through a 5 μM filter, concentrated by ultrafiltration and the buffer exchanged into MAC equilibration buffer (50 mM Na₂HPO₄, 300 mM NaCl, 10 mM imidazole, pH 8.0) by crossflow filtration. The resulting solution was passed through a 0.2 μM filter and further purified on a FPLC (Amersham-Pharmacia)

RECTIFIED SHEET (RULE 91)

-149-

using Pharmacia chelating sepharose. The solution was loaded onto the nickel loaded MAC at a flow rate of 1.0 mL/min and eluted with a linear gradient of 1.0 mM -1.0 M imidazole in 50 mM Na₂HPO₄, 300 mM NaCl, pH 8.0. Protein containing fractions were determined by SDS-PAGE and subsequently pooled
5 and frozen at -80°C.

Small amounts of purified MTSP3C122S were activated using plasmin sepharose for 30 min. at 37°C. The resin was spun down at 14K rpm for 5 min. and the protein solution removed. The resulting solution was screened for activity against of series of protease substrates; spec-tpa, spec-pl, spec-UK,
10 spec-fXIIa (American Diagnostica), S-2238, S-2266 (Kabi Diagnostica), S-2586, S-2366, S-2444, S-2288, S-2251, S-2302, S-2765, S-2222, spec-THE (Chromogenix), spec-fVIIa (Pentapharm). MTSP3C122S cleaved several of these substrates efficiently but was most active towards Spec-fXIIa, Spec-tPA, S-2765, Spec-fVIIa and S-2444.

15 5. **Gene expression profile of the serine protease MTSP3 in normal and tumor tissues**

To obtain information regarding the tissue distribution of the MTSP3 transcripts, the DNA insert encoding the MTSP3 protease domain was used to
20 probe a RNA blot composed of 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH, Palo Alto, CA). The expression pattern observed in decreasing signal level was: trachea = colon (descending) = esophagus > colon (ascending) > colon (transverse) = rectum > ileum > duodenum > jejunum > bladder > ilocecum > stomach >
25 kidney > appendix. It is also expressed less abundantly in fetal kidney, and in two tumor cell lines, HeLa S3 and leukemia, K-562. Northern analysis using RNA blots (catalog numbers 7780-1, 7765-1 & 7782-1; human 12-lane, human muscle and human digestive system multiple tissue northern (MTN) blots; CLONTECH) confirmed that the expression was detected most abundantly in the
30 colon, moderately in the esophagus, small intestine, bladder and kidney, and less abundantly in stomach and rectum. A single transcript of ~2.2 kb was detected.

-150-

Amplification of the MTSP3 transcript in several human primary tumors xenografted in mouse was performed using gene-specific primers. The MTSP3 transcript was detected in lung carcinoma (LX-1), colon adenocarcinoma (CX-1), colon adenocarcinoma (GI-112) and ovarian carcinoma (GI-102). No apparent
5 signal was detected in another form of lung carcinoma (GI-117), breast carcinoma (GI-101), pancreatic adenocarcinoma (GI-103) and prostatic adenocarcinoma (PC3).

EXAMPLE 2

Identification of genomic clone of MTSP4

10 Using the nucleotide sequence encoding the protease domain of the serine protease MTSP1 (also called matriptase), the protein database (SWISSPROT) at the National Center for Biotechnology Information (Bethesda, MD; <<http://www.ncbi.nlm.nih.gov>>) was searched for similar or identical
15 sequence to MTSP1 using the search algorithm blastx. The blastx algorithm compares the six-frame conceptual translation products of a nucleotide query sequence (both strands) against a protein sequence database. A protein encoding sequence (CAA18442) that has 37% identity to the MTSP1 protein sequence that was found to include a putative LDL-receptor domain and a trypsin-like serine protease domain was identified. This protein-encoding
20 sequence (hereinafter referred to as MTSP4) was found to be encoded by a genomic clone (AL022314) derived from human chromosome 22 sequenced by the Sanger Centre Chromosome 22 Mapping Group and deposited into the public database as part of the Human Genome Project. Subsequent search of the GenBank database showed that no identical sequence has been deposited. A
25 search of the EST database also did not show any matching human sequence, indicating that no human EST clone exists in the public database. Mouse EST clones (A1391417 and AA208793) are present and showed 88% identity to the serine protease at the nucleotide level.

30 PCR cloning of a genomic DNA fragment of MTSP4 for use as hybridization probe

In order to obtain tissue distribution profile of MTSP4 as well as to identify a tissue source for subsequent cloning of the cDNA, a genomic fragment

-151-

was amplified from human genomic DNA using two gene-specific primers, 5'-CCTCCACGGTGCTGTGGACCGTGTTCC-3' (5' primer) SEQ ID No. 21 and 5'-CCTCGCGCAAGGCGCCCCAGCCCG-3' (3' primer) SEQ ID No. 22. These two primers amplified a 265-base pair fragment within a single exon of MTSP4.

- 5 The fragment was then used as a hybridization probe on human tissue northern blot (human 12-lane multiple tissue northern (MTN) blot (catalog number 7780-1); CLONTECH, Palo Alto, CA). A prominent band (~2.6 kb) was detected in liver. Relatively weaker signals were obtained from the brain, heart, skeletal muscle and kidney. Since human liver showed a very strong signal, this
- 10 tissue was selected for the amplification of the MTSP4 cDNA.

5'- and 3'- rapid amplification of cDNA ends (RACE)

To obtain a full-length clone encoding MTSP4, 5'- and 3'-RACE reactions were performed. The Marathon-Ready cDNA library from human liver (CLONTECH) was used to isolate the 5' and 3' ends of the cDNA encoding

- 15 MTSP4. Marathon-Ready cDNA clones are specifically made for RACE reactions. Two gene specific primers were used:
- 5'-GCGTGGCGTCACCTGGTAGCGATAGACCTCGC -3' (SEQ ID No. 23) for 5'-RACE reaction and 5'-CCTCCACGGTGCTGTGGACCGTGTTCC-3' (SEQ ID No. 24) for 3'-RACE reaction. No fragment was obtained from the initial 5'-RACE
- 20 reaction.

- The 3'-RACE reaction, however, produced a ~1.5 kbp fragment. A nested PCR reaction was used on the initial 5'-RACE reaction products to obtain part of the 5' end of MTSP4. The nested 5' gene-specific primer used was 5'-CCTCGCGCAAGGCGCCCCAGCCCG-3' (SEQ ID No. 25) and produced a
- 25 ~0.8 kbp fragment. The fragments were subcloned into pCR2.1-TOPO TA cloning vector (Invitrogen, Carlsbad, CA). The resulting clones were analyzed by Southern analysis using the internal genomic fragment encompassing the primers used in the RACE reactions as probe and by DNA sequence analysis. Sequence analysis of the 5'-RACE product showed that the potential initiation codon was
- 30 still missing.

To obtain the 5' cDNA end that encodes the N terminus of MTSP4, the publicly available genomic sequence of chromosome 22 was searched for

-152-

sequence corresponding to the sequence obtained in the 5'-RACE clone. The resulting genomic sequence was translated and the protein sequence was compared to that derived from the translated sequence of the 5'-RACE clone. After determining the overlapping sequences, a gene-specific oligonucleotide primer (5'-TCATCGGCCAGAGGGTGATCAGTGAG-3') SEQ ID No. 26 corresponding to the sequence upstream of the potential initiation codon and another gene-specific oligonucleotide primer (5'-CCTCCTCAGTGCATAGGCATCAAACCAG-3') SEQ ID No. 27 corresponding to a sequence within the overlapping region were used to amplify the missing 5' cDNA of MTSP4 from the human liver cDNA library.

Splice variants and domain organization of MTSP4

At least two cDNA fragments were consistently obtained during PCR amplification, indicating multiple splice variants of MTSP4. Subcloning and sequence analysis revealed that a longer, more abundant form, MTSP4-L and a shorter form, MTSP4-S. The encoded proteins are multi-domain, type II membrane-type serine proteases and include a transmembrane domain at the N terminus followed by a CUB domain, 3 LDLR domains and a trypsin-like serine protease domain at the C terminus. The difference between these two forms of MTSP4 is the absence in MTSP4-S of a 432-bp nucleotide sequence between the transmembrane and the CUB domains (see FIGURE 2).

PCR amplification of cDNA encoding full-length protease domain of MTSP4

To obtain a cDNA fragment encoding the protease domain of MTSP4, an end-to-end PCR amplification using gene-specific primers and the Marathon-Ready cDNA library from human liver was used. The two primers used were: 5'-TCTCTCGAGAAAAGAATTGTTGGTGGAGCTGTGTCCTCCGAG-3' (SEQ ID No. 28) for the 5' end and 5'-AGGTGGGCCTTGCTTTGCAGGGGGGCGAGTTC-3' for the 3' end SEQ ID NO. 29). The 5' primer contained the sequence that encodes the start of the MTSP4 protease domain (IVGGAVSSE). The 3' primer corresponds to the sequence just downstream of the stop codon. A ~740-bp fragment was amplified, subcloned into pCR2.1-TOPO TA cloning vector and sequenced.

Gene expression profile of MTSP4 in normal and tumor tissues

-153-

To obtain information regarding the gene expression profile of the MTSP4 transcript, a DNA fragment encoding part of the LDL receptor domain and the protease domain was used to probe an RNA blot composed of 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH). As in the northern analysis of gel blot, a very strong signal was observed in the liver. Signals in other tissues were observed in (decreasing signal level): fetal liver > heart = kidney = adrenal gland = testis = fetal heart and kidney = skeletal muscle = bladder = placenta > brain = spinal cord = colon = stomach = spleen = lymph node = bone marrow = trachea = uterus = pancreas = salivary gland = mammary gland = lung. MTSP4 is also expressed less abundantly in several tumor cell lines including HeLa S3 = leukemia K-562 = Burkitt's lymphomas (Raji and Daudi) = colorectal adenocarcinoma (SW480) > lung carcinoma (A549) = leukemia MOLT-4 = leukemia HL-60. PCR of the MTSP4 transcript from cDNA libraries made from several human primary tumors xenografted in nude mice (human tumor multiple tissue cDNA panel, catalog number K1522-1, CLONTECH) was performed using MTSP4-specific primers. The MTSP4 transcript was detected in breast carcinoma (GI-101), lung carcinoma (LX-1), colon adenocarcinoma (GI-112) and pancreatic adenocarcinoma (GI-103). No apparent signal was detected in another form of lung carcinoma (GI-117), colon adenocarcinoma (CX-1), ovarian carcinoma (GI-102). and prostatic adenocarcinoma (PC3). The MTSP4 transcript was also detected in LNCaP and PC-3 prostate cancer cell lines as well as in HT-1080 human fibrosarcoma cell line.

Sequence analysis

MTSP4 DNA and protein sequences were analyzed using MacVector (version 6.5; Oxford Molecular Ltd., Madison, WI). The ORF of MTSP4-L includes 2,409 bp, which translate to a 802-amino acid protein, while the ORF of MTSP4-S is composed of 1,977 bp which translate to a 658-amino acid protein. The cDNA encoding the protease domain in both forms is composed of 708 bp which translate to a 235-amino acid protein sequence (see, SEQ ID No. 6) The DNA sequences and the translated protein sequences of MTSP4-L and

-154-

MTSP4-S, and of the protease domain of MTSP4 are set forth in SEQ ID Nos. 8, 10 and 6, respectively.

EXAMPLE 3

Cloning of MTSP6

5 Identification of genomic clone of MTSP6

Using the protein sequence of the protease domain of the serine protease MTSP4 (see EXAMPLE 2), the non-redundant database (all non-redundant GenBank CDS translations + PDB + SwissProt + PIR + PRF) at the National Center for Biotechnology Information (Bethesda, MD;

- 10 <<http://www.ncbi.nlm.nih.gov>>) was searched for sequences that were similar or identical to MTSP4 using the search algorithm tblastn. The tblastn algorithm compares a protein query sequence against a nucleotide sequence database dynamically translated in all reading frames. A protein (55 amino acids), which has 60% identity with the query MTSP4 sequence (55 amino acids), was
- 15 obtained from the translation of genomic sequence of AC015555 (nucleotide #15553 to 15717). This protein hereafter is referred to as MTSP6. Subsequent search of the GenBank database showed that no cDNA encoding MTSP6 has been deposited.

- The gene exhibiting highest homology to MTSP6 was human
- 20 transmembrane serine protease 2 (GenBank accession number U75329; Swissprot accession number O15393), which showed 66% identity to MTSP6 within the 45 amino acid regions compared. Consequently, the nucleotide sequence encoding the MTSP6 protease domain was obtained by comparing the protein sequence of human transmembrane serine protease 2 protease domain
- 25 with the nucleotide sequence of AC015555 translated in six reading frames. The protein sequence obtained from the translated nucleotide sequence of MTSP6 revealed an overall 50% identity with human transmembrane serine protease 2. A search of the EST database indicated the presence of seven MTSP6 EST clones (AA883068, AW591433, AI978874, AI469095, AI935487,
- 30 AA534591 and AI758271).

-155-

Cloning of human MTSP6 full-length cDNA

To obtain cDNA encoding the region of the MTSP6 protease domain identified by database searches described above, two gene-specific primers, Ch17-NSP-1, 5'-TCACGCATCGTGGGTGGAACATGTCC-3' (5' primer) SEQ ID NO. 30 and Ch17-NSP-2AS, 5'-ACCCACCTCCATCTGCTCGTGGATCC-3' SEQ ID NO. 31 (3' primer), were used for PCR. These two primers amplified a 708-base pair fragment from human mammary gland carcinoma cDNA (Clontech Marathon-Ready cDNA, Cat. No. 7493-1).

- To obtain the remaining, unknown cDNA of MTSP6, 5'- and 3'-RACE reactions were performed on the human mammary gland carcinoma. Marathon-Ready cDNA is specifically made for RACE reactions. The first RACE reactions were performed by PCR using Marathon cDNA adaptor primer 1 (AP1) with gene specific primers, Ch17-NSP-2AS, 5'-ACCCACCTCCATCTGCTCGTGGATCC-3' SEQ ID NO. 31 for 5'-RACE reaction and Ch17-NSP-1, 5'-TCACGCATCGTGGGTGGAACATGTCC-3' SEQ ID NO. 30 for 3'-RACE reaction. The PCR products were purified from agarose gel. A second nested PCR was then performed using Marathon cDNA adaptor primer 2 (AP2) with gene specific primer Ch17-NSP-3AS, 5'-CCACAGCCTCCTCTCTTGACACACCAG-3' SEQ ID No. 32 for 5'-RACE reaction (using first 5'-RACE product as template) and Ch17-NSP-3 5'-ACGCCCTGTGGATCATCACTGCTGC-3' SEQ ID No. 33 for 3'-RACE reaction (using first 3'-RACE product as template). First 5'- and 3'-RACE products were also used as template for PCR reactions using primers Ch17-NSP-3 and Ch17-NSP-4AS to obtain a cDNA fragment for use as a probe. PCR products from RACE reactions which were larger than 700 bp were cut out and purified from agarose gel and subcloned into pCR2.1-TOPO cloning vector (Invitrogen, Carlsbad, CA). Colony hybridization was then performed to identify positive colonies containing MTSP6 sequence. Positive clones were identified by colony hybridization using the 495 bp DNA fragment obtained from PCR reaction (with primers Ch17-NSP-3 and Ch17-NSP-4AS) and by DNA sequencing.

-156-

- Sequence analysis of the 5'-RACE products indicated that an additional 420 bp of upstream sequence were obtained. The potential initial codon was not present in the 5'-RACE sequence. Another round of nested 5'-RACE reaction was performed using AP2 and a gene specific primer (designed based
- 5 on the new RACE sequence) Ch17-NSP-5AS
5'-TCCCTCCCTCACATATACTGAGTGGTG-3' SEQ ID No. 34, using the PCR products obtained from the first 5'-RACE as template. A PCR product of 367 bp using Ch17-NSP-6 5'-CGACTGCTCAGGGAAGTCAGATGTCG-3' SEQ ID NO. 35 (designed based on the new 5'-RACE sequence) and Ch17-NSP-5AS was used
- 10 to identify the positive clones. An additional sequence of 480 bp was obtained from the second 5'-RACE products. A potential ATG start codon was observed within a sequence of GTCACCATGG (nucleotides 262-272 of SEQ ID No. 12, which appears to be a Kozak sequence (GCC (A/G) CCAUGG), indicating that this ATG is likely the initiation codon for MTSP6.
- 15 The 3'-RACE reaction to obtain the rest of the 3' end of MTSP6 was not successful using Marathon Ready human mammary gland carcinoma cDNA. The sequence of the 3'-RACE products obtained was exclusively that of an MTSP6 cDNA truncated with the Marathon AP2 primer sequence within the coding region.
- 20 The 3'-end sequence of MTSP6 was obtained by PCR using Ch17-NSP-3 (5'-ACGCCCTGTGGATCATCACTGCTGC-3'; SEQ ID NO. 33) and Ch17-NSP-4 (5'-CTGGTGTGTCAAGAGAGGAGGCTGTGG-3'; SEQ ID NO. 37) with an antisense primer Ch17-NSP-7AS (5'-ACTCAGGTGGCTACTTATCCCCTTCCTC-3'; SEQ ID NO. 38) designed based
- 25 on the sequence of an EST clone AA883068, which apparently covers the 3'-end of MTSP6 sequence, and human small intestine cDNA (Clontech) as template. Two PCR products (650 bp and 182 bp, respectively) were obtained and DNA sequence analysis indicated that both PCR products contained a stop codon.
- 30 **Sequence analysis and domain organization of MTSP6**
- The MTSP6 DNA and protein sequences were analyzed using DNA Strider (version 1.2). The ORF of MTSP6 is composed of 1,362 bp, which translate

-157-

into a 453-amino acid protein. Protein sequence analysis using the SMART (Simple Modular Architecture Research Tool) program at <http://smart.embl-heidelberg.de> predicts that MTSP6 is a multi-domain, type-II membrane-type serine protease containing of a transmembrane domain (amino acids 48-68) at the N terminus followed by a LDLRa domain (LDL receptor domain class a) (amino acids 72-108), a SR domain (Scavenger receptor Cys-rich domain)(amino acids 109-205), and a trypsin-like serine protease domain (amino acids 216-443) (see FIGURE 3).

Gene expression profile of MTSP6 in normal and tumor tissues

- 10 To obtain information regarding the gene expression profile of the MTSP6 transcript, a 495 bp DNA fragment obtained from PCR reaction with primers Ch17-NSP-3 and NSP-4AS was used to probe an RNA blot composed of 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH). The strongest signal was observed in
- 15 duodenum. Signal in other tissues were observed in (decreased signal level): Stomach > trachea = mammary gland = thyroid gland = salivary gland = pituitary gland = pancreas > kidney > lung > jejunum = ileum = ileocecum = appendix = fetal kidney > fetal lung. Very weak signals can also be detected in several other tissues. MTSP6 is also expressed in several tumor cell lines
- 20 including HeLa S3 > colorectal adenocarcinoma (SW480) > leukemia MOLT-4 > leukemia K-562. PCR analysis of the MTSP6 transcript from cDNA libraries made from several human primary tumors xenografted in nude mice (human tumor multiple tissue cDNA panel, catalog number K1522-1, CLONTECH) was performed using MTSP6-specific primers (Ch17-NSP-3 and Ch17-NSP2AS). The
- 25 MTSP6 transcript was strongly detected in lung carcinoma (LX-1), moderately detected in pancreatic adenocarcinoma (GI-103), weakly detected in ovarian carcinoma (GI-102); and very weakly detected in colon adenocarcinoma (GI-112 and CX-1), breast carcinoma (GI-101), lung carcinoma (GI-117) and prostatic adenocarcinoma (PC3). The MTSP6 transcript was also detected in breast
- 30 cancer cell line MDA-MB-231, prostate cancer cell line PC-3, but not in HT-1080 human fibrosarcoma cell line. MTSP6 is also expressed in mammary gland carcinoma cDNA (Clontech).

-158-

EXAMPLE 4**Expression of the protease MTSP domains**

The DNA encoding each of the MTSP 3 and 4 protease domains was cloned into a derivative of the *Pichia pastoris* vector pPIC9K (available from
5 Invitrogen; see SEQ ID NO. 45). Plasmid pPIC9k features include the 5' AOX1 promoter fragment at 1-948; 5' AOX1 primer site at 855-875; alpha-factor secretion signal(s) at 949-1218; alpha-factor primer site at 1152-1172; multiple cloning site at 1192-1241; 3' AOX1 primer site at 1327-1347; 3' AOX1 transcription termination region at 1253-1586; HIS4 ORF at 4514-1980;
10 kanamycin resistance gene at 5743-4928; 3' AOX1 fragment at 6122-6879; ColE1 origin at 7961-7288; and the ampicillin resistance gene at 8966-8106. The plasmid used herein is derived from pPIC9K by eliminating the XhoI site in the kanamycin resistance gene and the resulting vector is herein designated pPIC9KX.

- 15 **Primers used for PCR amplification of protease domain and subcloning into the XhoI/NotI sites of Pichia vector**

MTSP3

5' primer (with XhoI site [underlined]) SEQ ID No. 39

5' TCTCTCGAGAAAAGAGTGGTGGGTGGGGAGGAGGCCTCTGTG 3'

- 20 3' primer (with NotI site [underlined]) SEQ ID No. 40

5' ATTCGCGGCCGCGATTACAGCTCAGCCTTCCAGAC 3'

MTSP4-S and MTSP4-L

5' primer (with XhoI site [underlined]) SEQ ID No. 41

5' TCTCTCGAGAAAAGAATTGTTGGTGGAGCTGTGTCCTCCGAG

- 25 3' primer with NotI site SEQ ID No. 42

5' ATTCGCGGCCGCTCAGGTCACCACTTGCTGGATCCAG 3'

MTSP6

- MTSP6 was cloned into the *E. coli* TOPO vector (pcR[®] 2.1 TOPO[™], SEQ
ID No. 46, Invitrogen, Carlsbad, CA; the TOPO[®] TA Cloning[®] Kit is designed
30 form cloning Taq-amplified PRCR products).

5' primer (with XhoI site [underlined]) SEQ ID No. 43

5' CTCGAGAAACGCATCGTGGGTGGAAACATGTCCTTG 3'

-159-

3' primer NotI site comes from E. coli TOPO vector SEQ ID No. 44:

5' ACTCAGGTGGCTACTTATCCCCTTCCTC 3'

EXAMPLE 5

5 Assays for identification of candidate compounds that modulate that activity of an MTSP

Assay for identifying inhibitors

The ability of test compounds to act as inhibitors of catalytic activity of an MTSP, including MTSP1, MTSP3, MTSP4, MTSP6 can be assessed in an amidolytic assay. The inhibitor-induced inhibition of amidolytic activity by a recombinant MTSP or the protease domain portions thereof, can be measured by IC50 values in such an assay.

An exemplary assay buffer is HBSA (10 mM Hepes, 150mM sodium chloride, pH 7.4, 0.1% bovine serum albumin). All reagents were from Sigma Chemical Co. (St. Louis, MO), unless otherwise indicated. Two IC50 assays at 30-minute (a 30-minute preincubation of test compound and enzyme) and at 0-minutes (no preincubation of test compound and enzyme) are conducted. For the IC50 assay at 30-minute, the following reagents are combined in appropriate wells of a Corning microtiter plate: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering a broad concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the MTSP or protease domain thereof diluted in buffer, yielding a final enzyme concentration of about 100-500 pM. Following a 30-minute incubation at ambient temperature, the assay is initiated by the addition of 50 microliters of a substrate for the particular MTSP (see, *e.g.*, table and discussion below) and reconstituted in deionized water, followed by dilution in HBSA prior to the assay) were added to the wells, yielding a final volume of 200 microliters and a final substrate concentration of 300 μ M (about 1.5-times K_m).

For an IC50 assay at 0-minute, the same reagents are combined: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering the identical concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the substrate, such as a chromogenic substrate. The assay is initiated by the addition of 50 microliters of MTSP. The

-160-

final concentrations of all components are identical in both IC50 assays (at 30- and 0-minute incubations).

The initial velocity of the substrate hydrolysis is measured in both assays by, for example for a chromogenic substrate, as the change of absorbance at a particular wavelength, using a Thermo Max₂ Kinetic Microplate Reader (Molecular Devices) over a 5 minute period, in which less than 5% of the added substrate was used. The concentration of added inhibitor, which caused a 50% decrease in the initial rate of hydrolysis was defined as the respective IC50 value in each of the two assays (30-and 0-minute).

10 **Another assay for identifying inhibitors**

Test compounds for inhibition of the protease activity of the protease domain of is assayed in Costar 96 well tissue culture plates (Corning NY). Approximately 2-3 nM the MTSP or protease domain thereof is mixed with varying concentrations of inhibitor in 29.2 mM Tris, pH 8.4, 29.2 mM imidazole, 15 217 mM NaCl (100 mL final volume), and allowed to incubate at room temperature for 30 minutes. 400 mM substrate is added, and the reaction monitored in a SpectraMAX Plus microplate reader (Molecular Devices, Sunnyvale CA) by following the change in a parameter correlated with hydrolysis, such as absorbance for a chromogenic substrate for 1 hour at 37° C.

20

ASSAY FOR SCREENING MTSP6

The protease domain of MTSP6 expressed in Pichia pastoris is assayed for inhibition by various compounds in Costar 96 well tissue culture plates (Corning NY). Approximately 1-20 nM MTSP6 is mixed with varying 25 concentrations of inhibitor in 29.2 mM Tris, pH 8.4, 29.2 mM Imidazole, 217 mM NaCl (100 μ L final volume), and allowed to incubate at room temperature for 30 minutes. 500 μ M substrate Spectrozyme t-PA (American Diagnostica,, Greenwich, CT) is added, and the reaction is monitored in a SpectraMAX Plus microplate reader (Molecular Devices, Sunnyvale CA) by measuring the change 30 in absorbance at 405 nm for 30 minutes at 37°C.

Identification of substrates

-161-

Particular substrates for use in the assays can be identified empirically by testing substrates. The following list of substrates are exemplary of those that can be tested.

	Substrate name	Structure
5	S 2366	pyroGlu-Pro-Arg-pNA.HCl
	spectrozyme t-PA	CH ₃ SO ₂ -D-HHT-Gly-Arg-pNA.AcOH
	N-p-tosyl-Gly-Pro-Arg-pNA	N-p-tosyl-Gly-Pro-Arg-pNA
	Benzoyl-Val-Gly-Arg-pNA	Benzoyl-Val-Gly-Arg-pNA
	Pefachrome t-PA	CH ₃ SO ₂ -D-HHT-Gly-Arg-pNA
10	S 2765	N- α -Z-D-Arg-Gly-Arg-pNA.2HCl
	S 2444	pyroGlu-Gly-Arg-pNA.HCl
	S 2288	H-D-Ile-Pro-Arg-pNA.2HCl
	spectrozyme UK	Cbo-L-(γ)Glu(α -t-BuO)-Gly-Arg-pNA.2AcOH
	S 2302	H-D-Pro-Phe-Arg-pNA.2HCl
15	S 2266	H-D-Val-Leu-Arg-pNA.2HCl
	S 2222	Bz-Ile-Glu(g-OR)-Gly-Arg-pNA.HCl R = H(50%) and R = CH ₃ (50%)
	Chromozyme PK	Benzoyl-Pro-Phe-Arg-pNA
	S 2238	H-D-Phe-Pip-Arg-pNA.2HCl
	S 2251	H-D-Val-Leu-Lys-pNA.2HCl
20	Spectrozyme PI	H-D-Nle-HHT-Lys-pNA.2AcOH
		Pyr-Arg-Thr-Lys-Arg-AMC
		H-Arg-Gln-Arg-Arg-AMC
		Boc-Gln-Gly-Arg-AMC
		Z-Arg-Arg-AMC
25	Spectrozyme THE	H-D-HHT-Ala-Arg-pNA.2AcOH
	Spectrozyme fXIIa	H-D-CHT-Gly-Arg-pNA.2AcOH
		CVS 2081-6 (MeSO ₂ -dPhe-Pro-Arg-pNA)
		Pefachrome fVIIa (CH ₃ SO ₂ -D-CHA-But-Arg-pNA)

pNA = para-nitranilide (chromogenic)

30 AMC = amino methyl coumarin (fluorescent)

If none of the above substrates are cleaved, a coupled assay, described above, can be used. Briefly, test the ability of the protease to activate and enzyme, such as plasminogen and trypsinogen. To perform these assays, the single chain protease is incubated with a zymogen, such as plasminogen or
35 trypsinogen, in the presence of the a known substrate, such, lys-plasminogen,

-162-

for the zymogen. If the single chain activates the zymogen, the activated enzyme, such as plasmin and trypsin, will degrade the substrate therefor.

EXAMPLE 6

Isolation and Cloning of Matriptase

5 A. Cell type and growth of cells

Human prostate adenocarcinoma cell line, PC-3, was purchased from ATCC (catalog number CRL-1435; Manassas, VA). The cells were cultured at 37°C, 5% CO₂ in Ham's F-12K growth medium (catalog number 9077; Irvine) supplemented with 2 mM L-glutamine and 10% fetal bovine serum. All
10 subsequent cell manipulations were carried out according to the manufacturer's instructions. PC-3 cells were allowed to grow to about 90% confluence, and were then washed briefly with 1x phosphate buffered saline.

B. Isolation of total RNA, and purification and enrichment of polyA⁺ RNA

15 PC-3 cells were lysed in Trizol reagent (catalog number 15596; Life Technologies, Rockville, MD) and total RNA was isolated according to the manufacturer's protocol. The concentration of total RNA was estimated from absorbance reading at 260 nm. PolyA⁺ RNA was purified and enriched using oligo-dT beads (catalog number 70061; Oligotex, Qiagen, Valencia, CA).

20 C. Reverse-transcription and polymerase chain reaction (PCR)

PC-3-derived polyA⁺ RNA was converted to single-stranded cDNA (sscDNA) by reverse transcription using ProSTAR first-strand RT-PCR kit (catalog number 200420; Stratagene, La Jolla, CA) and SuperScript II RNase H⁻ reverse transcriptase (catalog number 18064-022; Life Technologies). After reverse
25 transcription, an aliquot of PC-3 sscDNA (4 µL) was subjected to PCR using 2 mM each of the sense and anti-sense degenerate oligonucleotide primers and *Taq* polymerase (catalog number 201203; Qiagen). Total reaction volume was 100 µL. The sequence of the sense primer was 5'-
TGGRT(I)VT(I)WS(I)GC(I)RC(I)CAYTG-3' (SEQ ID No. 13) and that of the anti-
30 sense was 5'(I)GG(I)CC(I)CC(I)SWRTC(I)CCYT(I)RCA(I)GHRTC-3' (SEQ ID No. 14), where R=A,G; V=G,A,C; W=A,T; S=G,C; Y=C,T; H=A,T,C. The primer sequences correspond to two highly conserved regions in all

-163-

chymotrypsin-like serine proteases and amplify PCR products ranging from approximately 400 to 500 base pairs.

D. Clone screening and sequencing

The PCR products were separated on a 2% agarose gel and purified using
5 a gel extraction kit (catalog number 28706; QIAquick gel extraction kit; Qiagen). The purified DNA fragments were ligated into pCR2.1-TOPO (catalog number K4500-01; Invitrogen, Carlsbad, CA). After transformation into *E. coli* cells, plasmid DNA was isolated and analyzed by digestion with *EcoRI* restriction enzyme. Clones that had inserted nucleic acid were further characterized by
10 sequencing using a fluorescent dye-based DNA sequencing method (catalog number 4303149; BigDye terminator cycle sequencing kit with AmpliTaq DNA polymerase; Perkin Elmer, Lincoln, CA). A total of 31 clones were sequenced and analyzed. All sequences were analyzed by a multiple nucleotide sequence alignment algorithm (blastn) (www.ncbi.nlm.nih.gov/blast) to identify identical or
15 closely related DNA deposited in GenBank (NCBI, Bethesda, MD). Those that did not show significant homology were further analyzed using blastx, which compares the six-frame conceptual translation products of a nucleotide sequence (both strands) against a protein sequence database (SwissProt). Eight clones yielded identical cDNA fragments that encode MTSP1. MTSP1 was
20 subsequently found to be identical to matriptase (GenBank accession number AF118224).

E. Rapid amplification of cDNA ends (RACE) and gene-specific amplification of MTSP1

To obtain DNA encoding the complete protease domain of MTSP1, RACE
25 and gene-specific amplification reactions were performed. A human prostate Marathon-Ready cDNA (catalog # 7418-1; Clontech) was used to isolate part of the cDNA encoding MTSP1. Marathon-Ready cDNA is prepared to contain a known hybridization sequence at the 5' and 3' ends of the ssDNA. The 3' region of MTSP1 cDNA was obtained by a 3'-RACE reaction using a gene
30 specific primer, 5'-CACCCCTTCTTCAATGACTTCACCTTCG-3' (SEQ ID No. 55). The 5' end of the MTSP1 protease domain was obtained by gene-specific amplification reaction using two MTSP1-specific primers, 5'-

-164-

TACCTCTCCTACGACTCC-3' (SEQ ID No. 56) for the sense primer and 5'-GAGGTTCTCGCAGGTGGTCTGGTTG-3' (SEQ ID No. 57) for the antisense primer. The sequences for these two primers were obtained from the human SNC19 mRNA sequence. The 3'-RACE reaction and gene-specific PCR produced

5 DNA fragments that were > 1 kbp in size. These fragments were subcloned into pCR2.1-TOPO (Invitrogen, San Diego, CA). After transformation into *E. coli* cells, plasmid DNA was isolated and analyzed by digestion with *EcoRI* restriction enzyme. Clones that had inserts were characterized by Southern blot analysis (using the internal cDNA fragment as probe) and by DNA sequence analysis.

10 **F. PCR amplification of cDNA encoding the protease domain of MTSP1**

To obtain a cDNA fragment encoding the entire protease domain of MTSP1, an end-to-end PCR amplification using gene-specific primers was used. The two primers used were: 5'-

15 CTCGAGAAAAGAGTTGTTGGGGGCACGGATGCGGATGAG-3' (SEQ ID No. 58) for the 5' end and 5'-GCGGCCGCACTATACCCAGTGTTCTCTTTGATCCA-3' (SEQ ID No. 36 for the 3' end. The 5' primer contained the sequence that encodes the start of the MTSP1 protease domain (VVGGTDADE) (SEQ. ID. NO. 10). The 3' primer contained the stop codon of MTSP1. A ~800-bp

20 fragment was amplified, purified and subcloned into the *Pichia pastoris* expression vector, pPIC9K, resulting in pPIC9K-MTSP1.

G. Gene expression profile of MTSP1 in normal tissues, cancer cells and cancer tissues

To obtain information regarding the tissue distribution and gene

25 expression level of MTSP1, the DNA insert from pPIC9K-MTSP1 was used to probe a blot containing RNA from 76 different human tissues (catalog number 7775-1; human multiple tissue expression (MTE) array; CLONTECH, Palo Alto, CA). Significant expression was observed in the colon (ascending, transverse and descending), rectum, trachea, esophagus and duodenum. Moderate

30 expression levels were observed in the jejunum, ileum, ileocecum, stomach, prostate, pituitary gland, appendix, kidney, lung, placenta, pancreas, thyroid gland, salivary gland, mammary gland, fetal kidney, and fetal lung. Lower

-165-

expression levels were seen in the spleen, thymus, peripheral blood leukocyte, lymph node, bone marrow, bladder, uterus, liver, adrenal gland, fetal heart, fetal liver, fetal spleen, and fetal thymus. A significant amount of the MTSP1 transcript was also detected in colorectal adenocarcinoma cell line (SW480),
5 Burkitt's lymphoma cell line (Daudi), and leukemia cell line (HL-60). RT-PCR of the MTSP1 transcript in several human primary tumors xenografted in athymic nude mice was performed using gene-specific primers. A high level of MTSP1 transcript was detected in colon adenocarcinoma (CX-1) and pancreatic adenocarcinoma (GI-103). Moderate levels were observed in another colon
10 adenocarcinoma (GI-112), ovarian carcinoma (GI-102), lung carcinoma (LX-1), and breast carcinoma (GI-101). Another lung carcinoma (GI-117) expressed a low level of the MTSP1 transcript. A similar RT-PCR was performed to detect the presence of the MTSP1 transcript in PC-3 and LNCaP cell lines. Both cell lines expressed significant amounts of MTSP1 transcript.

15 H. Sequence analysis

All derived DNA and protein sequences were analyzed using MacVector (version 6.5; Oxford Molecular Ltd., Madison, WI). The cDNA encoding the protease domain of MTSP1 is composed of 726 base pairs which translate into a 241-amino acid protein sequence (rMAP) (see SEQ ID No. 1, 2, 49 and 50).

20 EXAMPLE 7

Production of Recombinant Serine Protease Domain of Matriptase or MTSP1 (rMAP)

A. Fermentation

The production of multi-milligram amounts of rMAP was carried out by
25 fermentation in a BioFlo 3000 fermentor (New Brunswick Scientific, NJ) equipped with a 3.3 L capacity bioreactor using a SMD1168/pPIC9K:MTSP1 Sac SC1 clone. ZA001 complex media (10 g/L yeast extract, 20 g/L peptone, 40 g/L glycerol, 5 g/L ammonium sulfate, 0.2 g/L calcium sulfate dihydrate, 2 g/L magnesium sulfate heptahydrate, 2 g/L potassium sulfate, 25 g/L sodium
30 hexametaphosphate, 4.35 ml/L PTM1) was inoculated with 100 ml of an overnight culture of the *P. pastoris* transformant. The culture was supplemented

-166-

with 50% glycerol by fed-batch phase and induced for 18-24 hours with methanol controlled at 0.025%.

B. Purification of Recombinant Serine Protease Domain of Matriptase or MTSP1 (rMAP)

- 5 The rMAP was secreted into the culture medium, so the first step of the purification involved the removal of cells and cell debris by centrifugation at 5000 g for 30 minutes. The resulting supernatant was decanted, adjusted to pH 8.0 with 10 N NaOH, and filtered through a Sartobran 300 0.45 + 0.2 μ m capsule. This supernatant was concentrated to 1 L by ultrafiltration using a 10
- 10 kDa ultrafiltration cartridge (NC SRT UF system with AG/Technologies UFP-10-C-5A filter), and the buffer was exchanged by crossflow filtration into 50 mM tris-HCl, 50 mM NaCl, 0.05% tween-80, pH 8.0 (buffer A). The filtration unit was rinsed once with 1 L buffer A which was combined with the concentrate.
- 15 The concentrated rMAP-containing solution was passed over a 150 ml benzamidine column that had been equilibrated with buffer A, at a flow rate of 8 ml/min. The column was washed with 3 column volumes of 50 mM tris-HCl, 1.0 M NaCl, 0.05% tween-80, pH 8.0 (buffer B) and eluted with 3 column
- 20 volumes of 50 mM tris-HCl, 1.0 M L-arginine, 0.05% tween-80, pH 8.0 (buffer C). Fractions containing rMAP were identified by activity assay and pooled. This pooled material was concentrated to 10 ml using a JumboSep concentrator (Pall Gelman) and a 10 kDa cutoff membrane. Once concentrated to 10 ml, the buffer was exchanged into 50 mM Na₂HPO₄, 125 mM NaCl, pH 5.5 (buffer D) and the volume adjusted to 5-10 ml. The retentate was removed and the
- 25 concentrator washed with buffer D which was added to the concentrate. The total sample volume was adjusted 15 ml.
- The partially purified rMAP was passed through a 5 ml Q-sepharose Fast Flow HiTrap column (Amersham-Pharmacia Biotech) pre-equilibrated with 15 ml of buffer D. The flow through was collected. The HiTrap column was washed
- 30 with an additional 10 ml of buffer D. Both flow throughs were pooled, and the protein concentration was determined by measurement of OD₂₈₀ (using an extinction coefficient of 2.012 mg/OD₂₈₀). Purified rMAP was then

-167-

deglycosylated by the addition 0.1 μ l of Endoglycosidase H (ProZyme, 5 U/ml) per mg of protein and incubating overnight at 4°C with gentle swirling.

- The conductivity of the deglycosylated pool was adjusted to 2.0-3.0 mS/cm with Nanopure H₂O and the pH adjusted to 6.5 (-200-300 mL final volume). The rMAP was then further purified by anion exchange chromatography by loading directly onto a Pharmacia Akta Explorer system using a 7 mL Source 15Q anion exchange column (Amersham-Pharmacia Biotech). The protein was eluted in a buffer containing 50 mM HEPES, pH 6.5 with a 0-0.33 M NaCl gradient over 10 column volumes at a flow rate of 6 ml/min.
- 10 Fractions containing protein were pooled, and benzamidine was added to a final concentration of 10 mM. Protein purity was examined by SDS-PAGE and protein concentration determined by measurement of OD₂₈₀ and use of a theoretical extinction coefficient of 2.012 mg/OD₂₈₀.

EXAMPLE 8

15 Assays

Amidolytic Assay for Determining Inhibition of Serine Protease Activity of Matriptase or MTSP1

- The ability of test compounds to act as inhibitors of rMAP catalytic activity was assessed by determining the inhibitor-induced inhibition of
- 20 amidolytic activity by the MAP, as measured by IC₅₀ values. The assay buffer was HBSA (10 mM Hepes, 150mM sodium chloride, pH 7.4, 0.1% bovine serum albumin). All reagents were from Sigma Chemical Co. (St. Louis, MO), unless otherwise indicated.

- Two IC₅₀ assays (a) one at either 30-minutes or 60-minutes (a 30-minute or a 60-minute preincubation of test compound and enzyme) and (b) one at 0-minutes (no preincubation of test compound and enzyme) were conducted.
- For the IC₅₀ assay at either 30-minutes or 60-minutes, the following reagents were combined in appropriate wells of a Corning microtiter plate: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering a broad
- 30 concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the rMAP (Corvas International) diluted in buffer, yielding a final enzyme concentration of 250 pM as determined by active

-168-

site filtration. Following either a 30-minute or a 60-minute incubation at ambient temperature, the assay was initiated by the addition of 50 microliters of the substrate S-2765 (N- α -Benzyloxycarbonyl-D-arginyl-L-glycyl-L-arginine-p-nitroaniline dihydrochloride; DiaPharma Group, Inc.; Franklin, OH) to each well, yielding a final assay volume of 200 microliters and a final substrate concentration of 100 μ M (about 4-times K_m). Before addition to the assay mixture, S-2765 was reconstituted in deionized water and diluted in HBSA. For the IC_{50} assay at 0 minutes; the same reagents were combined: 50 microliters of HBSA, 50 microliters of the test compound, diluted (covering the identical concentration range) in HBSA (or HBSA alone for uninhibited velocity measurement), and 50 microliters of the substrate S-2765. The assay was initiated by the addition of 50 microliters of rMAP. The final concentrations of all components were identical in both IC_{50} assays (at 30- or 60- and 0-minute).

The initial velocity of chromogenic substrate hydrolysis was measured in both assays by the change of absorbance at 405 nM using a Thermo Max[®] Kinetic Microplate Reader (Molecular Devices) over a 5 minute period, in which less than 5% of the added substrate was used. The concentration of added inhibitor, which caused a 50% decrease in the initial rate of hydrolysis was defined as the respective IC_{50} value in each of the two assays (30- or 60-minutes and 0-minute).

***In vitro* enzyme assays for specificity determination**

The ability of compounds to act as a selective inhibitor of matriptase activity was assessed by determining the concentration of test compound that inhibits the activity of matriptase by 50%, (IC_{50}) as described in the above Example, and comparing IC_{50} value for matriptase to that determined for all or some of the following serine proteases: thrombin, recombinant tissue plasminogen activator (rt-PA), plasmin, activated protein C, chymotrypsin, factor Xa and trypsin.

The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin). The assay for IC_{50} determinations was conducted by combining in appropriate wells of a Corning microtiter plate, 50 microliters of HBSA, 50 microliters of the test compound at

-169-

- a specified concentration (covering a broad concentration range) diluted in HBSA (or HBSA alone for V_0 (uninhibited velocity) measurement), and 50 microliters of the enzyme diluted in HBSA. Following a 30 minute incubation at ambient temperature, 50 microliters of the substrate at the concentrations specified
- 5 below were added to the wells, yielding a final total volume of 200 microliters. The initial velocity of chromogenic substrate hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max® Kinetic Microplate Reader over a 5 minute period in which less than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial
- 10 rate of hydrolysis was defined as the IC_{50} value.

Thrombin (fIIa) Assay

- Enzyme activity was determined using the chromogenic substrate, Pefachrome t-PA (CH_3SO_2 -D-hexahydrotyrosine-glycyl-L-Arginine-p-nitroaniline, obtained from Pentapharm Ltd.). The substrate was reconstituted in deionized
- 15 water prior to use. Purified human α -thrombin was obtained from Enzyme Research Laboratories, Inc. The buffer used for all assays was HBSA (10 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% bovine serum albumin).
- IC_{50} determinations were conducted where HBSA (50 μ L), α -thrombin (50 μ l) (the final enzyme concentration is 0.5 nM) and inhibitor (50 μ l) (covering a
- 20 broad concentration range), were combined in appropriate wells and incubated for 30 minutes at room temperature prior to the addition of substrate Pefachrome-t-PA (50 μ l) (the final substrate concentration is 250 μ M, about 5 times K_m). The initial velocity of Pefachrome t-PA hydrolysis was measured by the change in absorbance at 405 nm using a Thermo Max® Kinetic Microplate
- 25 Reader over a 5 minute period in which less than 5% of the added substrate was used. The concentration of added inhibitor which caused a 50% decrease in the initial rate of hydrolysis was defined as the IC_{50} value.

Factor Xa

- Factor Xa catalytic activity was determined using the chromogenic
- 30 substrate S-2765 (N-benzyloxycarbonyl-D-arginine-L-glycine-L-arginine-p-nitroaniline), obtained from DiaPharma Group (Franklin, OH). All substrates were reconstituted in deionized water prior to use. The final concentration of S-2765

-170-

was 250 μ M (about 5-times K_m). Purified human Factor X was obtained from Enzyme Research Laboratories, Inc. (South Bend, IN) and Factor Xa (FXa) was activated and prepared from it as described [Bock, P.E., Craig, P.A., Olson, S.T., and Singh, P. *Arch. Biochem. Biophys.* 273:375-388 (1989)]. The enzyme was
5 diluted into HBSA prior to assay in which the final concentration was 0.25 nM.
Recombinant tissue plasminogen activator (rt-PA) Assay

rt-PA catalytic activity was determined using the substrate, Pefachrome
t-PA (CH_3SO_2 -D-hexahydrotyrosine-glycyl-L-arginine-p-nitroaniline, obtained from
Pentapharm Ltd.). The substrate was made up in deionized water followed by
10 dilution in HBSA prior to the assay in which the final concentration was 500
micromolar (about 3-times K_m). Human rt-PA (Activase®) was obtained from
Genentech Inc. The enzyme was reconstituted in deionized water and diluted
into HBSA prior to the assay in which the final concentration was 1.0 nM.

Plasmin Assay

15 Plasmin catalytic activity was determined using the chromogenic
substrate, S-2366 [L-pyroglutamyl-L-prolyl-L-arginine-p-nitroaniline
hydrochloride], which was obtained from DiaPharma group. The substrate was
made up in deionized water followed by dilution in HBSA prior to the assay in
which the final concentration was 300 micromolar (about 2.5-times K_m).
20 Purified human plasmin was obtained from Enzyme Research Laboratories, Inc.
The enzyme was diluted into HBSA prior to assay in which the final
concentration was 1.0 nM.

Activated Protein C (aPC) Assay

aPC catalytic activity was determined using the chromogenic substrate,
25 Pefachrome PC (delta-carbobenzloxy-D-lysine-L-prolyl-L-arginine-p-nitroaniline
dihydrochloride), obtained from Pentapharm Ltd.). The substrate was made up
in deionized water followed by dilution in HBSA prior to the assay in which the
final concentration was 400 micromolar (about 3-times K_m). Purified human
aPC was obtained from Hematologic Technologies, Inc. The enzyme was diluted
30 into HBSA prior to assay in which the final concentration was 1.0 nM.

-171-

Chymotrypsin Assay

Chymotrypsin catalytic activity was determined using the chromogenic substrate, S-2586 (methoxy-succinyl-L-arginine-L-prolyl-L-tyrosyl-p-nitroanilide), which was obtained from DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 100 micromolar (about 9-times K_m). Purified (3X-crystallized; CDI) bovine pancreatic alpha-chymotrypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

10 Trypsin Assay

Trypsin catalytic activity was determined using the chromogenic substrate, S-2222 (benzoyl-L-isoleucine-L-glutamic acid-[gamma-methyl ester]-L-arginine-p-nitroanilide), which was obtained from DiaPharma Group. The substrate was made up in deionized water followed by dilution in HBSA prior to the assay in which the final concentration was 250 micromolar (about 4-times K_m). Purified (3X-crystallized; TRL3) bovine pancreatic trypsin was obtained from Worthington Biochemical Corp. The enzyme was reconstituted in deionized water and diluted into HBSA prior to assay in which the final concentration was 0.5 nM.

20

Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the appended claims.

-172-

WHAT IS CLAIMED IS:

1. A substantially purified single chain polypeptide, comprising the protease domain of a type-II membrane-type serine protease (MTSP) or a catalytically active portion thereof, wherein:
 - 5 the MTSP portion of the protein consists essentially of the protease domain of the MTSP or a catalytically active portion thereof.
 2. The substantially purified polypeptide of claim 1, wherein the MTSP is not expressed on endothelial cells.
 3. The substantially purified polypeptide of claim 1, wherein the
10 MTSP is not expressed on normal endothelial cells *in vivo*.
 4. The substantially purified polypeptide of claim 1, wherein the MTSP is a human protein.
 5. The substantially purified polypeptide of claim 1 that consists essentially of the protease domain of an MTSP or a catalytically active portion of
15 the protease domain.
 6. The substantially purified polypeptide of claim 1, wherein the expression and/or activity of the MTSP in tumor cells differs from its level of expression and/or activity in non-tumor cells.
 7. The substantially purified polypeptide of claim 1, wherein the
20 MTSP is detectable in a body fluid at a level that differs from its level in body fluids in a subject not having a tumor.
 8. The substantially purified polypeptide of claim 1, wherein:
the MTSP is present in a tumor; and
a substrate or cofactor for the MTSP is expressed at levels that differ
25 from a non-tumor cell in the same tissue.
 9. The substantially purified polypeptide of claim 1, wherein:
the MTSP exhibits altered substrate specificity in the tumor compared to its specificity in a non-tumor cell in the same tissue.
 10. The substantially purified polypeptide of claim 1, wherein the
30 MTSP has an N-terminus that comprises IVNG, ILGG, VGLL or ILGG.
 11. The substantially purified polypeptide of claim 1, wherein the MTSP is selected from among MTSP1, MTSP3, MTSP4 and MTSP6.

RECTIFIED SHEET (RULE 91)

-173-

12. The substantially purified polypeptide of claim 1, wherein the protease domain comprises the sequence of amino acids set forth as amino acids 615-855 of SEQ ID No. 2, as amino acids 205-437 of SEQ ID NO. 4, as the amino acids in SEQ ID No. 6, or as amino acids 217-443 in SEQ ID No. 12.

5 13. The substantially purified polypeptide of claim 1 that has at least about 40%, 60%, 80%, 90% or 95% sequence identity with a protease domain that comprises the sequence of amino acids set forth as amino acids 615-855 of SEQ ID No. 2, as amino acids 205-437 of SEQ ID NO. 4, as the amino acids in SEQ ID No. 6, or as amino acids 217-443 in SEQ ID No. 12.

10 14. A polypeptide of claim 1, wherein the protease domain portion is encoded by a nucleic acid molecule that hybridizes under conditions of high stringency along its full length to a nucleic acid molecule comprising a sequence of nucleotides set forth in SEQ ID No: 1, 3, 5, 7, 9 or 11 or to a molecule that encodes the protein set forth in SEQ ID No: 2, 4, 6, 8, 10 or 12 or at least one
15 domain thereof.

15. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 1.

16. A mutein of the polypeptide of claim 1, wherein:
up to about 90% of the amino acids are replaced with another amino
20 acid;

and the resulting polypeptide is a single chain and has catalytic activity at least 10% of the unmutated polypeptide.

17. The mutein of claim 16, wherein up to about 95% of the amino acids are replaced.

25 18. The mutein of claim 16, wherein the resulting polypeptide is a single chain and has catalytic activity at least 50% of the unmutated polypeptide.

19. A mutein of the polypeptide of claim 1, wherein a free Cys in the protease domain is replaced with another amino acid, whereby the resulting
30 polypeptide exhibits proteolytic activity.

20. A mutein of the polypeptide of claim 1, wherein a free Cys in the protease domain is replaced with a serine.

-174-

21. A vector comprising the nucleic acid molecule of claim 15.
22. The vector of claim 21 that is an expression vector.
23. The vector of claim 21 that includes a sequence of nucleotides that directs secretion of any protein encoded by a sequence of nucleotides
5 operatively linked thereto.
24. The vector of claim 21 that is a *Pichia* vector or an *E. coli* vector.
25. A cell, comprising the vector of claim 21.
26. The cell of claim 25 that is a prokaryotic cell.
27. The cell of claim 25 that is a eukaryotic cell.
- 10 28. The cell of claim 25 that is selected from among a bacterial cell, a yeast cell, a plant cell, an insect cell and an animal cell.
29. The cell of claim 25 that is a mammalian cell.
30. A method for producing a polypeptide that contains a protease domain of an MTSP, comprising:
15 culturing the cell of claim 25 under conditions whereby the encoded protein is expressed by the cell; and
recovering the expressed protein.
31. The method of claim 30, wherein the cell is a pichia cell and the protein is secreted into the culture medium.
- 20 32. An antisense nucleic acid molecule that comprises at least 14 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the protease domain of an MTSP of claim 1; or
comprises at least 16 contiguous nucleotides or modified nucleotides that
25 are complementary to a contiguous sequence of nucleotides in the protease domain of an MTSP of claim 1.
33. An antibody that specifically binds to the single chain form of a protease domain of the polypeptide of claim 1, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a
30 polyclonal antibody or a monoclonal antibody.

RECTIFIED SHEET (RULE 91)

-175-

34. The polypeptide of claim 1, wherein the MTSP is selected from among corin, MTSP1, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4.
35. A conjugate, comprising:
- 5 a) a protein of claim 1, and
- b) a targeting agent linked to the protein directly or via a linker.
142. The conjugate of claim 35, wherein the targeting agent permits
- 10 i) affinity isolation or purification of the conjugate;
- ii) attachment of the conjugate to a surface;
- iii) detection of the conjugate; or
- iv) targeted delivery to a selected tissue or cell.
36. A combination, comprising:
- a) an inhibitor of the catalytic activity of the polypeptide of claim 1;
- and
- 15 b) another treatment or agent selected from anti-tumor and anti-angiogenic treatments or agents.
37. The combination of claim 36, wherein the inhibitor and the anti-tumor and/or anti-angiogenic agent are formulated in a single pharmaceutical composition or each is formulated in separate pharmaceutical compositions.
- 20 38. The combination of claim 36, wherein the inhibitor is selected from antibodies and antisense oligonucleotides.
39. A solid support comprising two or more polypeptides of claim 1 linked thereto either directly or via a linker.
40. The support of claim 39, wherein the polypeptides comprise an
- 25 array.
41. The support of claim 39, wherein the polypeptides comprise a plurality of different MTSP protease domains.
42. A method for identifying compounds that modulate the protease activity of an MTSP, comprising:
- 30 contacting a polypeptide of claim 1 with a substrate proteolytically cleaved by the MTSP, and, either simultaneously, before or after, adding a test compound or plurality thereof;

RECTIFIED SHEET (RULE 91)

-176-

measuring the amount of substrate cleaved in the presence of the test compound; and

selecting compounds that change the amount cleaved compared to a control, whereby compounds that modulate the activity of the MTSP are
5 identified.

43. The method of claim 42, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

44. The method of claim 42, wherein a plurality of the test substances
10 are screened simultaneously.

45. The method of claim 42, wherein the change in the amount cleaved is assessed by comparing the amount cleaved in the presence of the test compound with the amount in the absence of the test compound.

46. The method of claim 42, wherein a plurality of the test substances
15 are screened for simultaneously.

47. The method of claim 42, wherein a plurality of the polypeptides are linked to a solid support, either directly or via a linker.

48. The method of claim 42, wherein the polypeptides comprise an array.

49. The method of claim 42, wherein the polypeptides comprise a
20 plurality of different MTSP proteases.

50. A method of identifying a compound that specifically binds to a single chain protease domain of an MTSP, comprising:

contacting a polypeptide of claim 1 with a test compound or plurality
25 thereof under conditions conducive to binding thereof; and
identifying compounds that specifically bind to the MTSP single chain protease domain or compounds that inhibit binding of a compound known to bind to the MTSP single chain protease domain, wherein the known compound is contacted with the polypeptide before, simultaneously with or after the test
30 compound.

51. The method of claims 50, wherein the polypeptide is linked either directly or indirectly via a linker to a solid support.

-177-

52. The method of claim 50, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

53. The method of claim 50, wherein a plurality of the test substances
5 are screened for simultaneously.

54. The method of claim 51, wherein a plurality of the polypeptides are linked to a solid support.

55. A substantially purified membrane-type serine protease 3 (MTSP3).

56. The MTSP3 of claim 43 that is selected from the group consisting
10 of:

a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 3;

a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set
15 forth in SEQ ID No. 3;

a polypeptide that comprises the sequence of amino acids set forth as amino acids 205-437 of SEQ ID No. 4;

a polypeptide that comprises a sequence of amino acids having at least about 90% sequence identity with the sequence of amino acids set forth in
20 SEQ ID No. 4; and

a polypeptide encoded by a splice variant of the sequence of nucleotides set forth in SEQ ID No. 3.

57. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 56.

25 58. A vector comprising the nucleic acid molecule of claim 57.

59. A cell, comprising the vector of claim 58.

60. The cell of claim 59 that expresses the MTSP3 on its surface.

61. The cell of claim 59 that is a prokaryotic cell.

62. The cells of claim 59 that is a eukaryotic cell.

30 63. The cell of claim 59 that is selected from among a bacterial cell, a yeast cell, a yeast cell, a plant cell, an insect cell and an animal cell.

64. The cell of claim 59 that is a mammalian cell.

-178-

65. The cell of claim 59, wherein the MTSP3 comprises the sequence of amino acids set forth in SEQ ID No. 4; or a sequence of amino acids encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 3; or a protein having at
5 least about 90% sequence identity with the sequence of amino acids set forth in SEQ ID No. 4 and retaining protease activity.

66. A method for producing an MTSP3, comprising:
culturing the cell of claim 59 under conditions whereby the encoded protein is expressed by the cell; and
10 recovering the expressed protein.

67. An antisense nucleic acid molecule that comprises at least 14 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the nucleic acid molecule of claim 57; or
comprises at least 16 contiguous nucleotides or modified nucleotides that
15 are complementary to a contiguous sequence of nucleotides in the in the nucleic acid molecule of claim 57.

68. An antibody that specifically binds to the MTSP of claim 57, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

20 69. A method for treating or preventing a neoplastic disease, in a mammal, comprising administering to a mammal an effective amount of an inhibitor of an MTSP3 of claim 55.

70. The method of claim 69, wherein the inhibitor is an antibody that specifically binds to the MTSP3, or a fragment or derivative of the antibody
25 containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

71. A substantially purified membrane-type serine protease 4 (MTSP4).

72. The substantially purified MTSP4 that is an MTSP4-L or an MTSP4-S.

30 73. The MTSP4 of claim 71 that is selected from the group consisting of:

-179-

a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 7 or 9;

a polypeptide encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 7 or 9;

a polypeptide that comprises the sequence of amino acids set forth in SEQ ID No. 6, 8 or 10; and

a polypeptide encoded by a splice variant of the sequence of nucleotides set forth in SEQ ID No. 7 or 9.

10

74. The MTSP4 of claim 73 that is an MTSP4-L or an MTSP4-S.

75. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 73.

76. A vector comprising the nucleic acid molecule of claim 74.

15

77. A cell, comprising the vector of claim 76.

78. The cell of claim 77 that expresses the MTSP4 on its surface.

79. The cell of claim 77 that is a prokaryotic cell.

80. The cells of claim 77 that is a eukaryotic cell.

81. The cell of claim 77 that is selected from among a bacterial cell, a yeast cell, a yeast cell, a plant cell, an insect cell and an animal cell.

82. The cell of claim 77 that is a mammalian cell.

83. The cell of claim 77, wherein the MTSP4 comprises the sequence of amino acids set forth in SEQ ID No. 6, 8 or 10; or a sequence of amino acids encoded by a sequence of nucleotides that hybridizes under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 7 or 9; or a protein having at least about 90% sequence identity with the sequence of amino acids set forth in SEQ ID No. 8 or 10 and retaining protease activity.

84. A method for producing an MTSP4, comprising:
culturing the cell of claim 77 under conditions whereby the encoded protein is expressed by the cell; and
recovering the expressed protein.

30

-180-

85. An antisense nucleic acid molecule that comprises at least 14 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the nucleic acid molecule of claim 75 or comprises at least 16 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the nucleic acid molecule of claim 75.

86. An antibody that specifically binds to the MTSP of claim 72, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

10 87. An antibody that specifically binds to the MTSP of claim 73, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

88. A method for treating or preventing a neoplastic disease, in a mammal, comprising administering to a mammal an effective amount of an inhibitor of an MTSP4 of claim 71.

89. The method of claim 88, wherein the inhibitor is an antibody that specifically binds to an MTSP4, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

20 90. A substantially purified membrane-type serine protease 6 (MTSP6) selected from the group consisting of:

a polypeptide encoded by the sequence of nucleotides set forth in SEQ ID No. 11;

25 a polypeptide encoded by a sequence of nucleotides that hybridizes along the full length thereof under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 11;

a polypeptide that comprises the sequence of amino acids set forth as amino acids 217-443 of SEQ ID No. 12;

30 a polypeptide encoded by a splice variant of the sequence of nucleotides set forth in SEQ ID No. 11.

91. A nucleic acid molecule, comprising a sequence of nucleotides that encodes the polypeptide of claim 90.

-181-

92. A vector comprising the nucleic acid molecule of claim 91.
93. A cell, comprising the vector of claim 92.
94. The cell of claim 93 that expresses the MTSP6 on its surface.
95. The cell of claim 93 that is a prokaryotic cell.
- 5 96. The cells of claim 93 that is a eukaryotic cell.
97. The cell of claim 93 that is selected from among a bacterial cell, a yeast cell, a yeast cell, a plant cell, an insect cell and an animal cell.
98. The cell of claim 93 that is a mammalian cell.
99. The cell of claim 93, wherein the MTSP6 comprises the sequence
- 10 of amino acids set forth in SEQ ID No. 12; or a sequence of amino acids encoded by a sequence of nucleotides that hybridizes along the full length thereof under conditions of high stringency to the sequence of nucleotides set forth in SEQ ID No. 11; or a protein having at least about 95% sequence identity with the sequence of amino acids set forth in SEQ ID No. 12 and retaining
- 15 protease activity.
100. A method for producing an MTSP6, comprising:
culturing the cell of claim 93 under conditions whereby the encoded protein is expressed by the cell; and
recovering the expressed protein.
- 20 101. An antisense nucleic acid molecule that comprises at least 14 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the nucleic acid molecule of claim 91; or comprises at least 16 contiguous nucleotides or modified nucleotides that are complementary to a contiguous sequence of nucleotides in the in the nucleic
- 25 acid molecule of claim 91.
102. An antibody that specifically binds to the MTSP of claim 90, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.
- 30 103. A method for treating or preventing a neoplastic disease, in a mammal, comprising administering to a mammal an effective amount of an inhibitor of an MTSP6 of claim 90.

-182-

104. The method of claim 103, wherein the inhibitor is an antibody that specifically binds to the MTSP6, or a fragment or derivative of the antibody containing a binding domain thereof, wherein the antibody is a polyclonal antibody or a monoclonal antibody.

5 105. A recombinant non-human animal, wherein an endogenous gene of an MTSP has been deleted or inactivated by homologous recombination or insertional mutagenesis of the animal or an ancestor thereof.

106. A recombinant non-human animal of claim 105, wherein the MTSP is an MTSP1, MTSP3, MTSP4 or MTSP6.

10 107. A conjugate, comprising:

- a) an MTSP3 or MTSP4 or an MTSP6 of claim 90; and
- b) a targeting agent linked to the protein directly or via a linker.

108. The conjugate of claim 106, wherein the targeting agent permits

- i) affinity isolation or purification of the conjugate;
- 15 ii) attachment of the conjugate to a surface;
- iii) detection of the conjugate; or
- iv) targeted delivery to a selected tissue or cell.

109. A combination, comprising:

- a) an inhibitor of the catalytic activity of an MTSP3 or MTSP4 or
20 MTSP6 of claim 90; and
- b) another treatment or agent selected from anti-tumor and anti-angiogenic treatments or agents.

110. The combination of claim 109, wherein the inhibitor and the anti-tumor and/or anti-angiogenic agent are formulated in a single pharmaceutical
25 composition or each is formulated in separate pharmaceutical compositions.

111. The combination of claim 109, wherein the inhibitor is selected from antibodies and antisense oligonucleotides.

112. A solid support comprising two or more MTSP3 polypeptides and/or MTSP4 polypeptides and/or MTSP6 polypeptides of claim 90 linked
30 thereto either directly or via a linker.

113. The support of claim 112, wherein the polypeptides comprise an array.

-183-

114. A method for identifying compounds that modulate the protease activity of an MTSP selected from MTSP3, MTSP4 or MTSP6 of claim 90, comprising:

5 contacting the MTSP with a substrate proteolytically cleaved by the MTSP, and, either simultaneously, before or after, adding a test compound or plurality thereof;

 measuring the amount of substrate cleaved in the presence of the test compound; and

10 selecting compounds that change the amount cleaved compared to a control, whereby compounds that modulate the activity of the MTSP are identified.

115. The method of claim 114, wherein the test compounds are small molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

15 117. The method of claim 114, wherein the change in the amount cleaved is assessed by comparing the amount cleaved in the presence of the test compound with the amount in the absence of the test compound.

118. The method of claim 114, wherein a plurality of the test substances are screened for simultaneously.

20 119. The method of claim 118, wherein a plurality of the polypeptides are linked to a solid support.

120. A method of identifying a compound that specifically binds to a MTSP selected from MTSP3, MTSP4 and the MTSP6 of claim 90, comprising:

25 contacting the MTSP polypeptide with a test compound or plurality thereof under conditions conducive to binding thereof; and identifying compounds that specifically binds to the MTSP.

121. A method of identifying a compound that specifically binds to a MTSP selected from MTSP3, MTSP4 and the MTSP6 of claim 90, comprising:

30 contacting the MTSP polypeptide with a test compound or plurality thereof under conditions conducive to binding thereof; and

-184-

identifying compounds that specifically binds to the MTSP.

122. The method of any claims 121, wherein the polypeptide is linked either directly or indirectly via a linker to a solid support.

123. The method of claim 121, wherein the test compounds are small
5 molecules, peptides, peptidomimetics, natural products, antibodies or fragments thereof.

124. The method of claim 121, wherein a plurality of the test substances are screened for simultaneously.

125. The method of claim 124, wherein a plurality of the polypeptides,
10 are linked to a solid support.

126. An MTSP6 polypeptide, comprising amino acids set forth as amino acids 46-55 in SEQ ID No. 12 and/or amino acids 368-394 of SEQ ID No. 12, and that is encoded by a sequence of nucleic acids that hybridizes under moderate stringency to nucleic acid encoding the polypeptide set forth in SEQ ID
15 No. 12.

127. The polypeptide of claim 126 that comprises the amino acids set forth as amino acids 46-55 in SEQ ID No. 12 and/or amino acids 368-394 of SEQ ID No. 12, and that is encoded by a sequence of nucleic acids that hybridizes under high stringency along its full length or full length of the protease
20 domain to nucleic acid encoding the polypeptide set forth in SEQ ID No. 12.

128. The polypeptide of claim 126, comprising the sequence of amino acids set forth in SEQ ID No. 12.

129. An isolated nucleic acid molecule, comprising a sequence of nucleic acids that encodes the polypeptide of claim 126.

25 130. A method for treating tumors, comprising administering a prodrug that is specifically cleaved by an MTSP, whereby, upon contact with a cell that expresses MTSP activity, the prodrug is converted into an active drug.

131. The method of claim 130, wherein the MTSP is selected from among an MTSP3, MTSP4 and an MTSP6 of claim 90.

30 132. The method of claim 130, wherein the MTSP is selected from among corin, enterpeptidase, human airway trypsin-like protease (HAT), MTSP1, TMPRSS2, and TMPRSS4.

-185-

133. A method of detecting neoplastic disease, comprising: detecting an MTSP3, MTSP4 or MTSP6 of claim 90 in a biological sample, wherein the amount detected differs from the amount in a subject who does not have neoplastic disease.

5 134. The method of claim 133, wherein the biological sample is selected from the group consisting of blood, urine, saliva, tears, interstitial fluid, cerebrospinal fluid, ascites fluid, tumor tissue biopsy and circulating tumor cells.

135. The method of claim 133, wherein the extracellular domain of the MTSP3, MTSP4 or MTSP6 is in the sample.

10 136. A modulator of the activity of a MTSP, which modulator is identified by the method of any of claims 42 and 114.

137. Use of the polypeptide of claim 1 for identifying compounds that modulate the activity of an MTSP.

15 138. The use of claim 137, wherein the compounds inhibit the proteolytic activity thereof.

139 The use of claim 137, wherein the MTSP is selected from among MTSP1, MTSP3, MTSP4, MTSP6, corin, enterpeptidase, human airway trypsin-like protease (HAT), TMPRSS2, and TMPRSS4.

20 140. Use of a polypeptide of any of claims 55, 71 and 90 for treatment of a neoplastic disease.

141. Use of a polypeptide of claim 55, 71 and 90 for formulation of a medicament of treatment of neoplastic disease.




FIG. 1

Splice variant and domain organization of MTSP4




FIG. 2

3/4




FIG. 3

4/4

▽

MTSP3	194	LACGKS-----LKT	PRVVGEEASVDSWPWQVSIQYDKQHVC	CGGSILD	236
			205		
MTSP4-S	396	PQCDGRPD	CRDGSDEEHCECGLQGPSSRIVGGAVSSEGEWPWQASLQVRGRHICGGALIA		455
			424		
MTSP4-L	540	PQCDGRPD	CRDGSDEEHCECGLQGPSSRIVGGAVSSEGEWPWQASLQVRGRHICGGALIA		599
			568		
MTSP6	205	TACGHR-----RGYSSRIVGGNMSLLSQWPWQASLQFQGYHLC	CGGSVIT		248
			217 +		
MTSP3	237	PHWVLTA	AHCFRKHTDVFN--WKVRAGSDKLGS----	FPSLAVAKIIIIIEFNPMYPKDND	290
MTSP4-S	456	DRWVITA	AHCFQEDSMAS	TVLWTVFLGKVWQNSRWPGEVSFKVSRLLLHPYHEEDSHDYD	515
MTSP4-L	600	DRWVITA	AHCFQEDSMAS	TVLWTVFLGKVWQNSRWPGEVSFKVSRLLLHPYHEEDSHDYD	718
MTSP6	249	PLWIITA	AHCVYDLYLPKS--WTIQVGLVSLLD--NPAPSHLVEKIVYH	SKYKPKRLGND	304
MTSP3	291	IALMKLQ	FPLTFSGTVRPICLPFFDEELTPATPLWIIGWGFTKQNGGKMSDILLQASVQV		350
			*		
MTSP4-S	516	VALLQLD	HPVVRSAAVRPVCLPARSHFFEPGLHCWITGWGALRE-GGPISNALQKVDVQL		574
			*		
MTSP4-L	660	VALLQLD	HPVVRSAAVRPVCLPARSHFFEPGLHCWITGWGALRE-GGPISNALQKVDVQL		718
			*		
MTSP6	305	IALMKLAG	PLTFNEMIQPVCLPNSEENFPDGKVCWTSGWGATED-GGDASPVLNHA	AVPL	363
			*		
MTSP3	351	IDSTRCN	ADDA	YQGEVTEKMMCAGIPEGGVDTCQGDSGGPLMYQSDQ--WHVVGIVSWGY	408
MTSP4-S	575	IPQDLCS--EVYRYQVTPRMLCAGYRK	GKKDACQGDSGGPLVCKALSGRWFLAGLVSWGL		632
MTSP4-L	719	IPQDLCS--EVYRYQVTPRMLCAGYRK	GKKDACQGDSGGPLVCKALSGRWFLAGLVSWGL		776
MTSP6	364	ISNKICN	HRDVYGGIISPSMLCAGYLTGGVDSCQGDSGPLVCQERR-LWKVLVGATSF	GI	442
MTSP3	409	GCGGPST	PGVYTKVSA	YLNWIYNVWKAEL	437
MTSP4-S	633	GCGRPNY	FGVYTRITGVISW	IQQVVT	658
MTSP4-L	777	GCGRPNY	FGVYTRITGVISW	IQQVVT	802
MTSP6	423	GCAEVNK	PGVYTRVTSFLDWIHEQMERDLKT		453

▽ cleavage site
+ potential glycosylation site
* unpaired cysteine

FIG. 4

SUBSTITUTE SHEET (RULE 26)

ggg ggc ccg aag gac ttc ggc gcg gga ctc aag tac aac tcc cgg cac 100
Gly Gly Pro Lys Asp Phe Gly Ala Gly Leu Lys Tyr Asn Ser Arg His
15 20 25

gag	aaa	gtg	aat	ggc	ttg	gag	gaa	ggc	gtg	gag	ttc	ctg	cca	gtc	aac	148
Glu	Lys	Val	Asn	Gly	Leu	Glu	Glu	Gly	Val	Glu	Phe	Leu	Pro	Val	Asn	
			30					35					40			
aac	gtc	aag	aag	gtg	gaa	aag	cat	ggc	ccg	ggg	cgc	tgg	gtg	gtg	ctg	196
Asn	Val	Lys	Lys	Val	Glu	Lys	His	Gly	Pro	Gly	Arg	Trp	Val	Val	Leu	
		45					50					55				
gca	gcc	gtg	ctg	atc	ggc	ctc	ctc	ttg	gtc	ttg	ctg	ggg	atc	ggc	ttc	244
Ala	Ala	Val	Leu	Ile	Gly	Leu	Leu	Val	Leu	Leu	Gly	Ile	Gly	Phe		
	60					65					70					
ctg	gtg	tgg	cat	ttg	cag	tac	cgg	gac	gtg	cgt	gtc	cag	aag	gtc	ttc	292
Leu	Val	Trp	His	Leu	Gln	Tyr	Arg	Asp	Val	Arg	Val	Gln	Lys	Val	Phe	
	75				80					85					90	
aat	ggc	tac	atg	agg	atc	aca	aat	gag	aat	ttt	gtg	gat	gcc	tac	gag	340
Asn	Gly	Tyr	Met	Arg	Ile	Thr	Asn	Glu	Asn	Phe	Val	Asp	Ala	Tyr	Glu	
				95					100					105		
aac	tcc	aac	tcc	act	gag	ttt	gta	agc	ctg	gcc	agc	aag	gtg	aag	gac	388
Asn	Ser	Asn	Ser	Thr	Glu	Phe	Val	Ser	Leu	Ala	Ser	Lys	Val	Lys	Asp	
			110					115					120			
gcg	ctg	aag	ctg	ctg	tac	agc	gga	gtc	cca	ttc	ctg	ggc	ccc	tac	cac	436
Ala	Leu	Lys	Leu	Leu	Tyr	Ser	Gly	Val	Pro	Phe	Leu	Gly	Pro	Tyr	His	
		125					130					135				
aag	gag	tgc	gct	gtg	acg	gcc	ttc	agc	gag	ggc	agc	gtc	atc	gcc	tac	484
Lys	Glu	Ser	Ala	Val	Thr	Ala	Phe	Ser	Glu	Gly	Ser	Val	Ile	Ala	Tyr	
	140					145					150					
tac	tgg	tct	gag	ttc	agc	atc	cgc	cag	cac	ctg	gtg	gag	gag	gcc	gag	532
Tyr	Trp	Ser	Glu	Phe	Ser	Ile	Pro	Gln	His	Leu	Val	Glu	Glu	Ala	Glu	
	155				160					165					170	
cgc	gtc	atg	gcc	gag	gag	cgc	gta	gtc	atg	ctg	ccc	ccg	cgg	gcg	cgc	580
Arg	Val	Met	Ala	Glu	Glu	Arg	Val	Val	Met	Leu	Pro	Pro	Arg	Ala	Arg	
			175						180					185		
tcc	ctg	aag	tcc	ttt	gtg	gtc	acc	tca	gtg	gtg	gct	ttc	ccc	acg	gac	628
Ser	Leu	Lys	Ser	Phe	Val	Val	Thr	Ser	Val	Val	Ala	Phe	Pro	Thr	Asp	
			190					195					200			
tcc	aaa	aca	gta	cag	agg	acc	cag	gac	aac	agc	tgc	agc	ttt	ggc	ctg	676
Ser	Lys	Thr	Val	Gln	Arg	Thr	Gln	Asp	Asn	Ser	Cys	Ser	Phe	Gly	Leu	
		205					210					215				
cac	gcc	cgc	ggt	gtg	gag	ctg	atg	cgc	ttc	acc	acg	ccc	ggc	ttc	cct	724
His	Ala	Arg	Gly	Val	Glu	Leu	Met	Arg	Phe	Thr	Thr	Pro	Gly	Phe	Pro	
	220					225					230					
gac	agc	ccc	tac	ccc	gct	cat	gcc	cgc	tgc	cag	tgg	gcc	ctg	cgg	ggg	772
Asp	Ser	Pro	Tyr	Pro	Ala	His	Ala	Arg	Cys	Gln	Trp	Ala	Leu	Arg	Gly	
	235				240					245					250	
gac	gcc	gac	tca	gtg	ctg	agc	ctc	acc	ttc	cgc	agc	ttt	gac	ctt	gcg	820
Asp	Ala	Asp	Ser	Val	Leu	Ser	Leu	Thr	Phe	Arg	Ser	Phe	Asp	Leu	Ala	
				255					260					265		
tcc	tgc	gac	gag	cgc	ggc	agc	gac	ctg	gtg	acg	gtg	tac	aac	acc	ctg	868
Ser	Cys	Asp	Glu	Arg	Gly	Ser	Asp	Leu	Val	Thr	Val	Tyr	Asn	Thr	Leu	

270						275						280						
agc	ccc	atg	gag	ccc	cac	gcc	ctg	gtg	cag	ttg	tgt	ggc	acc	tac	cct	916		
Ser	Pro	Met	Glu	Pro	His	Ala	Leu	Val	Gln	Leu	Cys	Gly	Thr	Tyr	Pro			
		285					290					295						
ccc	tcc	tac	aac	ctg	acc	ttc	cac	tcc	tcc	cag	aac	gtc	ctg	ctc	atc	964		
Pro	Ser	Tyr	Asn	Leu	Thr	Phe	His	Ser	Ser	Gln	Asn	Val	Leu	Leu	Ile			
	300					305					310							
aca	ctg	ata	acc	aac	act	gag	cgg	cgg	cat	ccc	ggc	ttt	gag	gcc	acc	1012		
Thr	Leu	Ile	Thr	Asn	Thr	Glu	Arg	Arg	His	Pro	Gly	Phe	Glu	Ala	Thr			
	315				320					325					330			
ttc	ttc	cag	ctg	cct	agg	atg	agc	agc	tgt	gga	ggc	cgc	tta	cgt	aaa	1060		
Phe	Phe	Gln	Leu	Pro	Arg	Met	Ser	Ser	Cys	Gly	Gly	Arg	Leu	Arg	Lys			
				335					340					345				
gcc	cag	ggg	aca	ttc	aac	agc	ccc	tac	tac	cca	ggc	cac	tac	cca	ccc	1108		
Ala	Gln	Gly	Thr	Phe	Asn	Ser	Pro	Tyr	Tyr	Pro	Gly	His	Tyr	Pro	Pro			
			350					355					360					
aac	att	gac	tgc	aca	tgg	aac	att	gag	gtg	ccc	aac	aac	cag	cat	gtg	1156		
Asn	Ile	Asp	Cys	Thr	Trp	Asn	Ile	Glu	Val	Pro	Asn	Asn	Gln	His	Val			
		365					370					375						
aag	gtg	agc	ttc	aaa	ttc	ttc	tac	ctg	ctg	gag	ccc	ggc	gtg	cct	gcg	1204		
Lys	Val	Ser	Phe	Lys	Phe	Phe	Tyr	Leu	Leu	Glu	Pro	Gly	Val	Pro	Ala			
	380				385					390								
ggc	acc	tgc	ccc	aag	gac	tac	gtg	gag	atc	aat	ggg	gag	aaa	tac	tgc	1252		
Gly	Thr	Cys	Pro	Lys	Asp	Tyr	Val	Glu	Ile	Asn	Gly	Glu	Lys	Tyr	Cys			
	395			400						405					410			
gga	gag	agg	tcc	cag	ttc	gtc	gtc	acc	agc	aac	agc	aac	aag	atc	aca	1300		
Gly	Glu	Arg	Ser	Gln	Phe	Val	Val	Thr	Ser	Asn	Ser	Asn	Lys	Ile	Thr			
				415				420						425				
gtt	cgc	ttc	cac	tca	gat	cag	tcc	tac	acc	gac	acc	ggc	ttc	tta	gct	1348		
Val	Arg	Phe	His	Ser	Asp	Gln	Ser	Tyr	Thr	Asp	Thr	Gly	Phe	Leu	Ala			
			430					435					440					
gaa	tac	ctc	tcc	tac	gac	tcc	agt	gac	cca	tgc	ccg	ggg	cag	ttc	acg	1396		
Glu	Tyr	Leu	Ser	Tyr	Asp	Ser	Ser	Asp	Pro	Cys	Pro	Gly	Gln	Phe	Thr			
		445				450						455						
tgc	cgc	acg	ggg	cgg	tgt	atc	cgg	aag	gag	ctg	cgc	tgt	gat	ggc	tgg	1444		
Cys	Arg	Thr	Gly	Arg	Cys	Ile	Arg	Lys	Glu	Leu	Arg	Cys	Asp	Gly	Trp			
	460				465						470							
gcc	gac	tgc	acc	gac	cac	agc	gat	gag	ctc	aac	tgc	agt	tgc	gac	gcc	1492		
Ala	Asp	Cys	Thr	Asp	His	Ser	Asp	Glu	Leu	Asn	Cys	Ser	Cys	Asp	Ala			
				480						485					490			
ggc	cac	cag	ttc	acg	tgc	aag	aac	aag	ttc	tgc	aag	ccc	ctc	ttc	tgg	1540		
Gly	His	Gln	Phe	Thr	Cys	Lys	Asn	Lys	Phe	Cys	Lys	Pro	Leu	Phe	Trp			
			495					500						505				
gtc	tgc	gac	agt	gtg	aac	gac	tgc	gga	gac	aac	agc	gac	gag	cag	ggg	1588		
Val	Cys	Asp	Ser	Val	Asn	Asp	Cys	Gly	Asp	Asn	Ser	Asp	Glu	Gln	Gly			
			510					515					520					

tgc agt tgt ccg gcc cag acc ttc agg tgt tcc aat ggg aag tgc ctc Cys Ser Cys Pro Ala Gln Thr Phe Arg Cys Ser Asn Gly Lys Cys Leu 525 530 535	1636
tcg aaa agc cag cag tgc aat ggg aag gac gac tgt ggg gac ggg tcc Ser Lys Ser Gln Gln Cys Asn Gly Lys Asp Asp Cys Gly Asp Gly Ser 540 545 550	1684
gac gag gcc tcc tgc ccc aag gtg aac gtc gtc act tgt acc aaa cac Asp Glu Ala Ser Cys Pro Lys Val Asn Val Val Thr Cys Thr Lys His 555 560 565 570	1732
acc tac cgc tgc ctc aat ggg ctc tgc ttg agc aag ggc aac cct gag Thr Tyr Arg Cys Leu Asn Gly Leu Cys Leu Ser Lys Gly Asn Pro Glu 575 580 585	1780
tgt gac ggg aag gag gac tgt agc gac ggc tca gat gag aag gac tgc Cys Asp Gly Lys Glu Asp Cys Ser Asp Gly Ser Asp Glu Lys Asp Cys 590 595 600	1828
gac tgt ggg ctg cgg tca ttc acg aga cag gct cgt gtt gtt ggg ggc Asp Cys Gly Leu Arg Ser Phe Thr Arg Gln Ala Arg Val Val Gly Gly 605 610 615	1876
acg gat gcg gat gag ggc gag tgg ccc tgg cag gta agc ctg cat gct Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val Ser Leu His Ala 620 625 630	1924
ctg ggc cag ggc cac atc tgc ggt gct tcc ctc atc tct ccc aac tgg Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu Ile Ser Pro Asn Trp 635 640 645 650	1972
ctg gtc tct gcc gca cac tgc tac atc gat gac aga gga ttc agg tac Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp Arg Gly Phe Arg Tyr 655 660 665	2020
tca gac ccc acg cag tgg acg gcc ttc ctg ggc ttg cac gac cag agc Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly Leu His Asp Gln Ser 670 675 680	2068
cag cgc agc gcc cct ggg gtg cag gag cgc agg ctc aag cgc atc atc Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg Leu Lys Arg Ile Ile 685 690 695	2116
tcc cac ccc ttc ttc aat gac ttc acc ttc gac tat gac atc gcg ctg Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp Tyr Asp Ile Ala Leu 700 705 710	2164
ctg gag ctg gag aaa ccg gca gag tac agc tcc atg gtg cgg ccc atc Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser Met Val Arg Pro Ile 715 720 725 730	2212
tgc ctg ccg gac gcc tcc cat gtc ttc cct gcc ggc aag gcc atc tgg Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp 735 740 745	2260
gtc acg ggc tgg gga cac acc cag tat gga ggc act ggc gcg ctg atc Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly Thr Gly Ala Leu Ile 750 755 760	2308
ctg caa aag ggt gag atc cgc gtc atc aac cag acc acc tgc gag aac Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr Thr Cys Glu Asn	2356

765					770					775						
ctc	ctg	ccg	cag	cag	atc	acg	ccg	cgc	atg	atg	tgc	gtg	ggc	ttc	ctc	2404
Leu	Leu	Pro	Gln	Gln	Ile	Thr	Pro	Arg	Met	Met	Cys	Val	Gly	Phe	Leu	
	780					785					790					
agc	ggc	ggc	gtg	gac	tcc	tgc	cag	ggt	gat	tcc	ggg	gga	ccc	ctg	tcc	2452
Ser	Gly	Gly	Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Ser	
795					800					805					810	
agc	gtg	gag	gcg	gat	ggg	cgg	atc	ttc	cag	gcc	ggt	gtg	gtg	agc	tgg	2500
Ser	Val	Glu	Ala	Asp	Gly	Arg	Ile	Phe	Gln	Ala	Gly	Val	Val	Ser	Trp	
				815					820					825		
gga	gac	ggc	tgc	gct	cag	agg	aac	aag	cca	ggc	gtg	tac	aca	agg	ctc	2548
Gly	Asp	Gly	Cys	Ala	Gln	Arg	Asn	Lys	Pro	Gly	Val	Tyr	Thr	Arg	Leu	
			830					835					840			
cct	ctg	ttt	cgg	gac	tgg	atc	aaa	gag	aac	act	ggg	gta	ta	ggggccgggg		2599
Pro	Leu	Phe	Arg	Asp	Trp	Ile	Lys	Glu	Asn	Thr	Gly	Val				
		845					850					855				
ccacccaaat	gtgtacacct	gcgggggccac	ccatcgctcca	ccccagtggtg	cacgacctgca											2659
ggctggagac	tggaccgctg	actgcaccag	cgccccagaga	acatacactg	tgaactcaat											2719
ctccagggct	ccaaatctgc	ctagaaaacc	tctcgcttcc	tcagcctcca	aagtggagct											2779
gggaggtaga	agggggaggac	actggtggtt	ctactgaccc	aactgggggc	aaagggttga											2839
agacacagcc	tccccgccca	gccccaaagt	gggcccaggc	gcgttttgtg	atatctgcct											2899
ccccgtctctg	taaggagcag	cgggaacgga	gcttcggagc	ctcctcagtg	aaggtggtgg											2959
ggctgccgga	tctgggctgt	ggggcccttg	ggccacgctc	ttgaggaagc	ccaggctcgg											3019
aggaccctg	aaaacagacg	ggtctgagac	tgaattgtt	ttaccagctc	ccagggtgga											3079
cttcactgtg	tgtatttgtg	taaatgggta	aaacaattta	tttcttttta	aaaaaaaaaa											3139
aaaaaaaaa																3147
<210>	2															
<211>	855															
<212>	PRT															
<213>	Homo Sapien															
<400>	2															
Met	Gly	Ser	Asp	Arg	Ala	Arg	Lys	Gly	Gly	Gly	Gly	Pro	Lys	Asp	Phe	
1				5					10					15		
Gly	Ala	Gly	Leu	Lys	Tyr	Asn	Ser	Arg	His	Glu	Lys	Val	Asn	Gly	Leu	
			20					25					30			
Glu	Glu	Gly	Val	Glu	Phe	Leu	Pro	Val	Asn	Asn	Val	Lys	Lys	Val	Glu	
		35					40					45				
Lys	His	Gly	Pro	Gly	Arg	Trp	Val	Val	Leu	Ala	Ala	Val	Leu	Ile	Gly	
	50					55					60					
Leu	Leu	Leu	Val	Leu	Leu	Gly	Ile	Gly	Phe	Leu	Val					

Val	Thr	Ser	180	Val	Val	Ala	Phe	Pro	185	Thr	Asp	Ser	Lys	Thr	190	Val	Gln	Arg
Thr	Gln	195	Asn	Ser	Cys	Ser	200	Phe	Gly	Leu	His	Ala	205	Arg	Gly	Val	Glu	
Leu	Met	210	Arg	Phe	Thr	Thr	215	Pro	Gly	Phe	Pro	Asp	220	Ser	Pro	Tyr	Pro	Ala
225	His	Ala	Arg	Cys	Gln	Trp	230	Ala	Leu	Arg	Gly	Asp	235	Ala	Asp	Ser	Val	Leu
Ser	Leu	Thr	245	Phe	Arg	Ser	250	Phe	Asp	Leu	Ala	Ser	255	Cys	Asp	Glu	Arg	Gly
Ser	Asp	Leu	260	Val	Thr	Val	265	Tyr	Asn	Thr	Leu	Ser	270	Pro	Met	Glu	Pro	His
Ala	Leu	Val	275	Gln	Leu	Cys	280	Gly	Thr	Tyr	Pro	Pro	285	Ser	Tyr	Asn	Leu	Thr
Phe	His	Ser	290	Ser	Gln	Asn	295	Val	Leu	Leu	Ile	Thr	300	Leu	Ile	Thr	Asn	Thr
305	Glu	Arg	Arg	His	Pro	Gly	310	Phe	Glu	Ala	Thr	Phe	315	Phe	Gln	Leu	Pro	Arg
Met	Ser	Ser	325	Cys	Gly	Gly	330	Arg	Leu	Arg	Lys	Ala	335	Gln	Gly	Thr	Phe	Asn
Ser	Pro	Tyr	340	Tyr	Pro	Gly	345	His	Tyr	Pro	Pro	Asn	350	Ile	Asp	Cys	Thr	Trp
Asn	Ile	Glu	355	Val	Pro	Asn	360	Asn	His	Val	Lys	Val	365	Ser	Phe	Lys	Phe	
Phe	Tyr	Leu	370	Leu	Glu	Pro	375	Gly	Val	Pro	Ala	Gly	380	Thr	Cys	Pro	Lys	Asp
385	Tyr	Val	Glu	Ile	Asn	Gly	390	Glu	Lys	Tyr	Cys	Gly	395	Glu	Arg	Ser	Gln	Phe
Val	Val	Thr	405	Ser	Asn	Ser	410	Asn	Lys	Ile	Thr	Val	415	Arg	Phe	His	Ser	Asp
Gln	Ser	Tyr	420	Thr	Asp	Thr	425	Gly	Phe	Leu	Ala	Glu	430	Tyr	Leu	Ser	Tyr	Asp
Ser	Ser	Asp	435	Pro	Cys	Pro	440	Gly	Gln	Phe	Thr	Cys	445	Arg	Thr	Gly	Arg	Cys
Ile	Arg	Lys	450	Glu	Leu	Arg	455	Cys	Asp	Gly	Trp	Ala	460	Asp	Cys	Thr	Asp	His
465	Ser	Asp	Glu	Leu	Asn	Cys	470	Ser	Cys	Asp	Ala	Gly	475	His	Gln	Phe	Thr	Cys
Lys	Asn	Lys	485	Phe	Cys	Lys	490	Pro	Leu	Phe	Trp	Val	495	Cys	Asp	Ser	Val	Asn
Asp	Cys	Gly	500	Asp	Asn	Ser	505	Glu	Gln	Gly	Cys	Ser	510	Cys	Pro	Ala	Gln	
Thr	Phe	Arg	515	Cys	Ser	Asn	520	Gly	Lys	Cys	Leu	Ser	525	Lys	Ser	Gln	Gln	Cys
Asn	Gly	Lys	530	Asp	Asp	Cys	535	Gly	Asp	Gly	Ser	Asp	540	Glu	Ala	Ser	Cys	Pro
545	Lys	Val	Asn	Val	Val	Thr	550	Cys	Thr	Lys	His	Thr	555	Tyr	Arg	Cys	Leu	Asn
Gly	Leu	Cys	565	Leu	Ser	Lys	570	Gly	Asn	Pro	Glu	Cys	575	Asp	Gly	Lys	Glu	Asp
Cys	Ser	Asp	580	Gly	Ser	Asp	585	Glu	Lys	Asp	Cys	Asp	590	Cys	Gly	Leu	Arg	Ser
Phe	Thr	Arg	595	Gln	Ala	Arg	600	Val	Val	Gly	Gly	Thr	605	Asp	Ala	Asp	Glu	Gly
Glu	Trp	Pro	610	Trp	Gln	Val	615	Ser	Leu	His	Ala	Leu	620	Gly	Gln	Gly	His	Ile
625	Cys	Gly	Ala	Ser	Leu	Ile	630	Ser	Pro	Asn	Trp	Leu	635	Val	Ser	Ala	Ala	His
Cys	Tyr	Ile	645	Asp	Asp	Arg	650	Gly	Phe	Arg	Tyr	Ser	655	Asp	Pro	Thr	Gln	Trp
Thr	Ala	Phe	660	Leu	Gly	Leu	665	His	Asp	Gln	Ser	Gln	670	Arg	Ser	Ala	Pro	Gly

Val	Gln	Glu	Arg	Arg	Leu	Lys	Arg	Ile	Ile	Ser	His	Pro	Phe	Phe	Asn
690						695					700				
Asp	Phe	Thr	Phe	Asp	Tyr	Asp	Ile	Ala	Leu	Leu	Glu	Leu	Glu	Lys	Pro
705					710					715					720
Ala	Glu	Tyr	Ser	Ser	Met	Val	Arg	Pro	Ile	Cys	Leu	Pro	Asp	Ala	Ser
				725					730					735	
His	Val	Phe	Pro	Ala	Gly	Lys	Ala	Ile	Trp	Val	Thr	Gly	Trp	Gly	His
				740				745					750		
Thr	Gln	Tyr	Gly	Gly	Thr	Gly	Ala	Leu	Ile	Leu	Gln	Lys	Gly	Glu	Ile
		755					760					765			
Arg	Val	Ile	Asn	Gln	Thr	Thr	Cys	Glu	Asn	Leu	Leu	Pro	Gln	Gln	Ile
	770					775					780				
Thr	Pro	Arg	Met	Met	Cys	Val	Gly	Phe	Leu	Ser	Gly	Gly	Val	Asp	Ser
785					790					795					800
Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Ser	Ser	Val	Glu	Ala	Asp	Gly
				805					810					815	
Arg	Ile	Phe	Gln	Ala	Gly	Val	Val	Ser	Trp	Gly	Asp	Gly	Cys	Ala	Gln
			820					825					830		
Arg	Asn	Lys	Pro	Gly	Val	Tyr	Thr	Arg	Leu	Pro	Leu	Phe	Arg	Asp	Trp
		835					840					845			
Ile	Lys	Glu	Asn	Thr	Gly	Val									
	850					855									

<210> 3

<211> 2137

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (261)...(1574)

<223> DNA sequence encoding a transmembrane serine
protease (MTSP3) protein

<400> 3

ccatccta	at	acgactc	act	atagggctcg	agcggccgcc	cgggcaggtc	agagagagggc	60
agcagcttgc	tcagcggaca	aggatgctgg	gcgtgagggga	ccaaggcctg	ccctgcactc			120
gggcctcctc	cagccagtgc	tgaccagggga	cttctgacct	gctggccagc	caggacctgt			180
gtggggagggc	cctcctgctg	ccttgggggtg	acaatctcag	ctccaggcta	cagggagacc			240
gggaggatca	cagagccagc	atg tta cag gat	cct gac agt gat	caa cct ctg				293
		Met Leu Gln Asp	Pro Asp Ser Asp	Gln Pro Leu				
		1	5	10				
aac agc ctc	gat gtc aaa	ccc ctg cgc	aaa ccc cgt	atc ccc atg	gag			341
Asn Ser Leu	Asp Val Lys	Pro Leu Arg	Lys Pro Arg	Ile Pro Met	Glu			
	15	20		25				
acc ttc aga	aag gtg ggg	atc ccc atc	atc ata gca	cta ctg agc	ctg			389
Thr Phe Arg	Lys Val Gly	Ile Pro Ile	Ile Ile Ala	Leu Leu Ser	Leu			
	30	35		40				
gcg agt atc	atc att gtg	gtt gtc ctc	atc aag gtg	att ctg gat	aaa			437
Ala Ser Ile	Ile Ile Val	Val Val Leu	Ile Lys Val	Ile Leu Asp	Lys			
	45	50		55				
tac tac ttc	ctc tgc ggg	cag cct ctc	cac ttc atc	ccg agg aag	cag			485
Tyr Tyr Phe	Leu Cys Gly	Gln Pro Leu	His Phe Ile	Pro Arg Lys	Gln			
	60	65		70	75			
ctg tgt gac	gga gag ctg	gac tgt ccc	ttg ggg gag	gac gag gag	cac			533
Leu Cys Asp	Gly Glu Leu	Asp Cys Pro	Leu Gly Glu	Asp Glu Glu	His			

tgt Cys	gtc Val	aag Lys	agc Ser 95	ttc Phe	ccc Pro	gaa Glu	ggg Gly	cct Pro 100	gca Ala	gtg Val	gca Ala	gtc Val	cgc Arg 105	ctc Leu	tcc Ser	581
aag Lys	gac Asp	cga Arg 110	tcc Ser	aca Thr	ctg Leu	cag Gln	gtg Val 115	ctg Leu	gac Asp	tcg Ser	gcc Ala	aca Thr 120	ggg Gly	aac Asn	tgg Trp	629
ttc Phe	tct Ser 125	gcc Ala	tgt Cys	ttc Phe	gac Asp	aac Asn 130	ttc Phe	aca Thr	gaa Glu	gct Ala	ctc Leu 135	gct Ala	gag Glu	aca Thr	gcc Ala	677
tgt Cys 140	agg Arg	cag Gln	atg Met	ggc Gly	tac Tyr 145	agc Ser	agc Ser	aaa Lys	ccc Pro	acc Thr 150	ttc Phe	aga Arg	gct Ala	gtg Val	gag Glu 155	725
att Ile	ggc Gly	cca Pro	gac Asp	cag Gln 160	gat Asp	ctg Leu	gat Asp	gtt Val	gtt Val 165	gaa Glu	atc Ile	aca Thr	gaa Glu	aac Asn 170	agc Ser	773
cag Gln	gag Glu	ctt Leu 175	cgc Arg	atg Met	cgg Arg	aac Asn	tca Ser	agt Ser 180	ggg Gly	ccc Pro	tgt Cys	ctc Leu	tca Ser 185	ggc Gly	tcc Ser	821
ctg Leu	gtc Val	tcc Ser 190	ctg Leu	cac His	tgt Cys	ctt Leu	gcc Ala 195	tgt Cys	ggg Gly	aag Lys	agc Ser	ctg Leu 200	aag Lys	acc Thr	ccc Pro	869
cgt Arg	gtg Val 205	gtg Val	ggg Gly	ggg Gly	gag Glu	gag Glu 210	gcc Ala	tct Ser	gtg Val	gat Asp	tct Ser 215	tgg Trp	cct Pro	tgg Trp	cag Gln	917
gtc Val 220	agc Ser	atc Ile	cag Gln	tac Tyr	gac Asp 225	ata Ile	cag Gln	cac His	gtc Val 230	tgt Cys	gga Gly	ggg Gly	agc Ser	atc Ile	ctg Leu 235	965
gac Asp	ccc Pro	cac His	tgg Trp	gtc Val 240	ctc Leu	acg Thr	gca Ala	gcc Ala	cac His 245	tgc Cys	ttc Phe	agg Arg	aaa Lys	cat His 250	acc Thr	1013
gat Asp	gtg Val	ttc Phe	aac Asn 255	tgg Trp	aag Lys	gtg Val	cgg Arg	gca Ala 260	ggc Gly	tca Ser	gac Asp	aaa Lys	ctg Leu 265	ggc Gly	agc Ser	1061
ttc Phe	cca Pro	tcc Ser 270	ctg Leu	gct Ala	gtg Val	gcc Ala	aag Lys 275	atc Ile	atc Ile	atc Ile	att Ile	gaa Glu 280	ttc Phe	aac Asn	ccc Pro	1109
atg Met	tac Tyr 285	ccc Pro	aaa Lys	gac Asp	aat Asn	gac Asp 290	atc Ile	gcc Ala	ctc Leu	atg Met	aag Lys 295	ctg Leu	cag Gln	ttc Phe	cca Pro	1157
ctc Leu 300	act Thr	ttc Phe	tca Ser	ggc Gly	aca Thr 305	gtc Val	agg Arg	ctc Leu	atc Ile	tgt Cys 310	ctg Leu	ccc Pro	ttc Phe	ttt Phe	gat Asp 315	1205
gag Glu	gag Glu	ctc Leu	act Thr	cca Pro 320	gcc Ala	acc Thr	cca Pro	ctc Leu	tgg Trp 325	atc Ile	att Ile	gga Gly	tgg Trp	ggc Gly 330	ttt Phe	1253

9/66

```

acg aag cag aat gga ggg aag atg tct gac ata ctg ctg cag gcg tca 1301
Thr Lys Gln Asn Gly Gly Lys Met Ser Asp Ile Leu Leu Gln Ala Ser
335 340 345

gtc cag gtc att gac agc aca cgg tgc aat gca gac gat gcg tac cag 1349
Val Gln Val Ile Asp Ser Thr Arg Cys Asn Ala Asp Asp Ala Tyr Gln
350 355 360

ggg gaa gtc acc gag aag atg atg tgt gca ggc atc ccg gaa ggg ggt 1397
Gly Glu Val Thr Glu Lys Met Met Cys Ala Gly Ile Pro Glu Gly Gly
365 370 375

gtg gac acc tgc cag ggt gac agt ggt ggg ccc ctg atg tac caa tct 1445
Val Asp Thr Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Tyr Gln Ser
380 385 390 395

gac cag tgg cat gtg gtg ggc atc gtt agc tgg ggc tat ggc tgc ggg 1493
Asp Gln Trp His Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Gly
400 405 410

ggc ccg agc acc cca gga gta tac acc aag gtc tca gcc tat ctc aac 1541
Gly Pro Ser Thr Pro Gly Val Tyr Thr Lys Val Ser Ala Tyr Leu Asn
415 420 425

tgg atc tac aat gtc tgg aag gct gag ctg taa tgctgctgcc cctttgcagt 1594
Trp Ile Tyr Asn Val Trp Lys Ala Glu Leu *
430 435

gctgggagcc gcttccctcc tgccctgccc acctggggat ccccccaggc cagacacaga 1654
gcaagagtc ccttgggtac acccctctgc ccacagcctc agcatttctt ggagcagcaa 1714
agggcctcaa ttctgttaag agaccctcgc agcccagagg cgcccagagg aagtcagcag 1774
ccctagctcg gccacacttg gtgctcccag catcccaggg agagacacag cccactgaac 1834
aaggtctcag ggttattgct aagccaagaa ggaactttcc cactactactg aatggaagca 1894
ggctgtcttg taaaagccca gatcactgtg ggctggagag gagaaggaaa gggctctgcgc 1954
cagccctgtc cgtcttcacc catccccaag cctactagag caagaaacca gttgtaatat 2014
aaaatgcact gccctactgt tggatatgact accgttacct actgttgtca ttgttattac 2074
agctatggcc actattatta aagagctgtg taacaaaaaa aaaaaaaaaa aaaaaaaaaa 2134
aaa 2137

```

<210> 4
 <211> 437
 <212> PRT
 <213> Homo Sapien

```

<400> 4
Met Leu Gln Asp Pro Asp Ser Asp Gln Pro Leu Asn Ser Leu Asp Val
1 5 10 15
Lys Pro Leu Arg Lys Pro Arg Ile Pro Met Glu Thr Phe Arg Lys Val
20 25 30
Gly Ile Pro Ile Ile Ile Ala Leu Leu Ser Leu Ala Ser Ile Ile Ile
35 40 45
Val Val Val Leu Ile Lys Val Ile Leu Asp Lys Tyr Tyr Phe Leu Cys
50 55 60
Gly Gln Pro Leu His Phe Ile Pro Arg Lys Gln Leu Cys Asp Gly Glu
65 70 75 80
Leu Asp Cys Pro Leu Gly Glu Asp Glu Glu His Cys Val Lys Ser Phe
85 90 95
Pro Glu Gly Pro Ala Val Ala Val Arg Leu Ser Lys Asp Arg Ser Thr
100 105 110
Leu Gln Val Leu Asp Ser Ala Thr Gly Asn Trp Phe Ser Ala Cys Phe
115 120 125
Asp Asn Phe Thr Glu Ala Leu Ala Glu Thr Ala Cys Arg Gln Met Gly

```

10/66

130 135 140
 Tyr Ser Ser Lys Pro Thr Phe Arg Ala Val Glu Ile Gly Pro Asp Gln
 145 150 155
 Asp Leu Asp Val Val Glu Ile Thr Glu Asn Ser Gln Glu Leu Arg Met
 165 170 175
 Arg Asn Ser Ser Gly Pro Cys Leu Ser Gly Ser Leu Val Ser Leu His
 180 185 190
 Cys Leu Ala Cys Gly Lys Ser Leu Lys Thr Pro Arg Val Val Gly Gly
 195 200 205
 Glu Glu Ala Ser Val Asp Ser Trp Pro Trp Gln Val Ser Ile Gln Tyr
 210 215 220
 Asp Ile Gln His Val Cys Gly Gly Ser Ile Leu Asp Pro His Trp Val
 225 230 235
 Leu Thr Ala Ala His Cys Phe Arg Lys His Thr Asp Val Phe Asn Trp
 245 250 255
 Lys Val Arg Ala Gly Ser Asp Lys Leu Gly Ser Phe Pro Ser Leu Ala
 260 265 270
 Val Ala Lys Ile Ile Ile Ile Glu Phe Asn Pro Met Tyr Pro Lys Asp
 275 280 285
 Asn Asp Ile Ala Leu Met Lys Leu Gln Phe Pro Leu Thr Phe Ser Gly
 290 295 300
 Thr Val Arg Leu Ile Cys Leu Pro Phe Phe Asp Glu Glu Leu Thr Pro
 305 310 315
 Ala Thr Pro Leu Trp Ile Ile Gly Trp Gly Phe Thr Lys Gln Asn Gly
 325 330 335
 Gly Lys Met Ser Asp Ile Leu Leu Gln Ala Ser Val Gln Val Ile Asp
 340 345 350
 Ser Thr Arg Cys Asn Ala Asp Asp Ala Tyr Gln Gly Glu Val Thr Glu
 355 360 365
 Lys Met Met Cys Ala Gly Ile Pro Glu Gly Gly Val Asp Thr Cys Gln
 370 375 380
 Gly Asp Ser Gly Gly Pro Leu Met Tyr Gln Ser Asp Gln Trp His Val
 385 390 395
 Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Gly Gly Pro Ser Thr Pro
 405 410 415
 Gly Val Tyr Thr Lys Val Ser Ala Tyr Leu Asn Trp Ile Tyr Asn Val
 420 425 430
 Trp Lys Ala Glu Leu
 435

<210> 5

<211> 708

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (1)...(708)

<223> MTSP4 protease domain cDNA

<400> 5

att gtt ggt gga gct gtg tcc tcc gag ggt gag tgg cca tgg cag gcc 48
 Ile Val Gly Gly Ala Val Ser Ser Glu Gly Glu Trp Pro Trp Gln Ala
 1 5 10 15
 agc ctc cag gtt cgg ggt cga cac atc tgt ggg ggg gcc ctc atc gct 96
 Ser Leu Gln Val Arg Gly Arg His Ile Cys Gly Gly Ala Leu Ile Ala
 20 25 30
 gac cgc tgg gtg ata aca gct gcc cac tgc ttc cag gag gac agc atg 144
 Asp Arg Trp Val Ile Thr Ala Ala His Cys Phe Gln Glu Asp Ser Met
 35 40 45

11/66

```

gcc tcc acg gtg ctg tgg acc gtg ttc ctg ggc aag gtg tgg cag aac      192
Ala Ser Thr Val Leu Trp Thr Val Phe Leu Gly Lys Val Trp Gln Asn
      50                      55                      60

tcg cgc tgg cct gga gag gtg tcc ttc aag gtg agc cgc ctg ctc ctg      240
Ser Arg Trp Pro Gly Glu Val Ser Phe Lys Val Ser Arg Leu Leu Leu
      65                      70                      75                      80

cac ccg tac cac gaa gag gac agc cat gac tac gac gtg gcg ctg ctg      288
His Pro Tyr His Glu Glu Asp Ser His Asp Tyr Asp Val Ala Leu Leu
                        85                      90                      95

cag ctc gac cac ccg gtg gtg cgc tgc gcc gcc gtg cgc ccc gtc tgc      336
Gln Leu Asp His Pro Val Val Arg Ser Ala Ala Val Arg Pro Val Cys
                        100                      105                      110

ctg ccc gcg cgc tcc cac ttc ttc gag ccc ggc ctg cac tgc tgg att      384
Leu Pro Ala Arg Ser His Phe Phe Glu Pro Gly Leu His Cys Trp Ile
      115                      120                      125

acg ggc tgg ggc gcc ttg cgc gag ggc ggc ccc atc agc aac gct ctg      432
Thr Gly Trp Gly Ala Leu Arg Glu Gly Gly Pro Ile Ser Asn Ala Leu
      130                      135                      140

cag aaa gtg gat gtg cag ttg atc cca cag gac ctg tgc agc gag gtc      480
Gln Lys Val Asp Val Gln Leu Ile Pro Gln Asp Leu Cys Ser Glu Val
      145                      150                      155                      160

tat cgc tac cag gtg acg cca cgc atg ctg tgt gcc gcc tac cgc aag      528
Tyr Arg Tyr Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys
                        165                      170                      175

ggc aag aag gat gcc tgt cag ggt gac tca ggt ggt ccg ctg gtg tgc      576
Gly Lys Lys Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys
                        180                      185                      190

aag gca ctc agt ggc cgc tgg ttc ctg gcg ggg ctg gtc agc tgg ggc      624
Lys Ala Leu Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly
      195                      200                      205

ctg ggc tgt ggc cgg cct aac tac ttc ggc gtc tac acc cgc atc aca      672
Leu Gly Cys Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr
      210                      215                      220

ggt gtg atc agc tgg atc cag caa gtg gtg acc tga      708
Gly Val Ile Ser Trp Ile Gln Gln Val Val Thr *
      225                      230                      235

```

<210> 6
 <211> 235
 <212> PRT
 <213> Homo Sapien

```

<400> 6
Ile Val Gly Gly Ala Val Ser Ser Glu Gly Glu Trp Pro Trp Gln Ala
 1      5      10      15
Ser Leu Gln Val Arg Gly Arg His Ile Cys Gly Gly Ala Leu Ile Ala
      20      25      30
Asp Arg Trp Val Ile Thr Ala Ala His Cys Phe Gln Glu Asp Ser Met
      35      40      45

```

12/66

Ala Ser Thr Val Leu Trp Thr Val Phe Leu Gly Lys Val Trp Gln Asn
 50 55 60
 Ser Arg Trp Pro Gly Glu Val Ser Phe Lys Val Ser Arg Leu Leu Leu
 65 70 75 80
 His Pro Tyr His Glu Glu Asp Ser His Asp Tyr Asp Val Ala Leu Leu
 85 90 95
 Gln Leu Asp His Pro Val Val Arg Ser Ala Ala Val Arg Pro Val Cys
 100 105 110
 Leu Pro Ala Arg Ser His Phe Phe Glu Pro Gly Leu His Cys Trp Ile
 115 120 125
 Thr Gly Trp Gly Ala Leu Arg Glu Gly Gly Pro Ile Ser Asn Ala Leu
 130 135 140
 Gln Lys Val Asp Val Gln Leu Ile Pro Gln Asp Leu Cys Ser Glu Val
 145 150 155 160
 Tyr Arg Tyr Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys
 165 170 175
 Gly Lys Lys Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys
 180 185 190
 Lys Ala Leu Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly
 195 200 205
 Leu Gly Cys Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr
 210 215 220
 Gly Val Ile Ser Trp Ile Gln Gln Val Val Thr
 225 230 235

<210> 7

<211> 3104

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (33)...(2441)

<223> cDNA encoding :MTSP4-L (long form) splice variant

<400> 7

tcacgcggcca gaggggtgatc agtgagcaga ag atg ccc gtg gcc gag gcc ccc 53
 Met Pro Val Ala Glu Ala Pro
 1 5

cag gtg gct ggc ggg cag ggg gac gga ggt gat ggc gag gaa gcg gag 101
 Gln Val Ala Gly Gly Gln Gly Asp Gly Gly Asp Gly Glu Glu Ala Glu
 10 15 20

ccg gag ggg atg ttc aag gcc tgt gag gac tcc aag aga aaa gcc cgg 149
 Pro Glu Gly Met Phe Lys Ala Cys Glu Asp Ser Lys Arg Lys Ala Arg
 25 30 35

ggc tac ctc cgc ctg gtg ccc ctg ttt gtg ctg ctg gcc ctg ctc gtg 197
 Gly Tyr Leu Arg Leu Val Pro Leu Phe Val Leu Leu Ala Leu Leu Val
 40 45 50 55

ctg gct tcg gcg ggg gtg cta ctc tgg tat ttc cta ggg tac aag gcg 245
 Leu Ala Ser Ala Gly Val Leu Leu Trp Tyr Phe Leu Gly Tyr Lys Ala
 60 65 70

gag gtg atg gtc agc cag gtg tac tca ggc agt ctg cgt gta ctc aat 293
 Glu Val Met Val Ser Gln Val Tyr Ser Gly Ser Leu Arg Val Leu Asn
 75 80 85

cgc cac ttc tcc cag gat ctt acc cgc cgg gaa tct agt gcc ttc cgc 341
 Arg His Phe Ser Gln Asp Leu Thr Arg Arg Glu Ser Ser Ala Phe Arg

13/66

90	95	100	
agt gaa acc gcc aaa gcc cag aag atg ctc aag gag ctc atc acc agc Ser Glu Thr Ala Lys Ala Gln Lys Met Leu Lys Glu Leu Ile Thr Ser 105 110 115			389
acc cgc ctg gga act tac tac aac tcc agc tcc gtc tat tcc ttt ggg Thr Arg Leu Gly Thr Tyr Tyr Asn Ser Ser Ser Val Tyr Ser Phe Gly 120 125 130 135			437
gag gga ccc ctc acc tgc ttc ttc tgg ttc att ctc caa atc ccc gag Glu Gly Pro Leu Thr Cys Phe Phe Trp Phe Ile Leu Gln Ile Pro Glu 140 145 150			485
cac cgc cgg ctg atg ctg agc ccc gag gtg gtg cag gca ctg ctg gtg His Arg Arg Leu Met Leu Ser Pro Glu Val Val Gln Ala Leu Leu Val 155 160 165			533
gag gag ctg ctg tcc aca gtc aac agc tcg gct gcc gtc ccc tac agg Glu Glu Leu Leu Ser Thr Val Asn Ser Ser Ala Val Pro Tyr Arg 170 175 180			581
gcc gag tac gaa gtg gac ccc gag ggc cta gtg atc ctg gaa gcc agt Ala Glu Tyr Glu Val Asp Pro Glu Gly Leu Val Ile Leu Glu Ala Ser 185 190 195			629
gtg aaa gac ata gct gca ttg aat tcc acg ctg ggt tgt tac cgc tac Val Lys Asp Ile Ala Ala Leu Asn Ser Thr Leu Gly Cys Tyr Arg Tyr 200 205 210 215			677
agc tac gtg ggc cag ggc cag gtc ctc cgg ctg aag ggg cct gac cac Ser Tyr Val Gly Gln Gly Gln Val Leu Arg Leu Lys Gly Pro Asp His 220 225 230			725
ctg gcc tcc agc tgc ctg tgg cac ctg cag ggc ccc aag gac ctc atg Leu Ala Ser Ser Cys Leu Trp His Leu Gln Gly Pro Lys Asp Leu Met 235 240 245			773
ctc aaa ctc cgg ctg gag tgg acg ctg gca gag tgc cgg gac cga ctg Leu Lys Leu Arg Leu Glu Trp Thr Leu Ala Glu Cys Arg Asp Arg Leu 250 255 260			821
gcc atg tat gac gtg gcc ggg ccc ctg gag aag agg ctc atc acc tcg Ala Met Tyr Asp Val Ala Gly Pro Leu Glu Lys Arg Leu Ile Thr Ser 265 270 275			869
gtg tac ggc tgc agc cgc cag gag ccc gtg gtg gag gtt ctg gcg tcg Val Tyr Gly Cys Ser Arg Gln Glu Pro Val Val Glu Val Leu Ala Ser 280 285 290 295			917
ggg gcc atc atg gcg gtc gtc tgg aag aag ggc ctg cac agc tac tac Gly Ala Ile Met Ala Val Val Trp Lys Lys Gly Leu His Ser Tyr Tyr 300 305 310			965
gac ccc ttc gtg ctc tcc gtg cag cgg gtg gtc ttc cag gcc tgt gaa Asp Pro Phe Val Leu Ser Val Gln Pro Val Val Phe Gln Ala Cys Glu 315 320 325			1013
gtg aac ctg acg ctg gac aac agg ctc gac tcc cag ggc gtc ctc agc Val Asn Leu Thr Leu Asp Asn Arg Leu Asp Ser Gln Gly Val Leu Ser 330 335 340			1061

acc ccg tac ttc ccc agc tac tac tcg ccc caa acc cac tgc tcc tgg Thr Pro Tyr Phe Pro Ser Tyr Tyr Ser Pro Gln Thr His Cys Ser Trp 345 350 355	1109
cac ctc acg gtg ccc tct ctg gac tac ggc ttg gcc ctc tgg ttt gat His Leu Thr Val Pro Ser Leu Asp Tyr Gly Leu Ala Leu Trp Phe Asp 360 365 370 375	1157
gcc tat gca ctg agg agg cag aag tat gat ttg ccg tgc acc cag ggc Ala Tyr Ala Leu Arg Arg Gln Lys Tyr Asp Leu Pro Cys Thr Gln Gly 380 385 390	1205
cag tgg acg atc cag aac agg agg ctg tgt ggc ttg cgc atc ctg cag Gln Trp Thr Ile Gln Asn Arg Arg Leu Cys Gly Leu Arg Ile Leu Gln 395 400 405	1253
ccc tac gcc gag agg atc ccc gtg gtg gcc acg gcc ggg atc acc atc Pro Tyr Ala Glu Arg Ile Pro Val Val Ala Thr Ala Gly Ile Thr Ile 410 415 420	1301
aac ttc acc tcc cag atc tcc ctc acc ggg ccc ggt gtg cgg gtg cac Asn Phe Thr Ser Gln Ile Ser Leu Thr Gly Pro Gly Val Arg Val His 425 430 435	1349
tat ggc ttg tac aac cag tcg gac ccc tgc cct gga gag ttc ctc tgt Tyr Gly Leu Tyr Asn Gln Ser Asp Pro Cys Pro Gly Glu Phe Leu Cys 440 445 450 455	1397
tct gtg aat gga ctc tgt gtc cct gcc tgt gat ggg gtc aag gac tgc Ser Val Asn Gly Leu Cys Val Pro Ala Cys Asp Gly Val Lys Asp Cys 460 465 470	1445
ccc aac gcc ctg gat gag aga aac tgc gtt tgc aga gcc aca ttc cag Pro Asn Gly Leu Asp Glu Arg Asn Cys Val Cys Arg Ala Thr Phe Gln 475 480 485	1493
tgc aaa gag gac agc aca tgc atc tca ctg ccc aag gtc tgt gat ggg Cys Lys Glu Asp Ser Thr Cys Ile Ser Leu Pro Lys Val Cys Asp Gly 490 495 500	1541
cag cct gat tgt ctc aac ggc agc gac gaa gag cag tgc cag gaa ggg Gln Pro Asp Cys Leu Asn Gly Ser Asp Glu Glu Gln Cys Gln Glu Gly 505 510 515	1589
gtg cca tgt ggg aca ttc acc ttc cag tgt gag gac cgg agc tgc gtg Val Pro Cys Gly Thr Phe Thr Phe Gln Cys Glu Asp Arg Ser Cys Val 520 525 530 535	1637
aag aag ccc aac ccg cag tgt gat ggg cgg ccc gac tgc agg gac ggc Lys Lys Pro Asn Pro Gln Cys Asp Gly Arg Pro Asp Cys Arg Asp Gly 540 545 550	1685
tcg gat gag gag cac tgt gaa tgt ggc ctc cag ggc ccc tcc agc cgc Ser Asp Glu Glu His Cys Glu Cys Gly Leu Gln Gly Pro Ser Ser Arg 555 560 565	1733
att gtt ggt gga gct gtg tcc tcc gag ggt gag tgg cca tgg cag gcc Ile Val Gly Gly Ala Val Ser Ser Glu Gly Glu Trp Pro Trp Gln Ala 570 575 580	1781
agc ctc cag gtt cgg ggt cga cac atc tgt ggg ggg gcc ctc atc gct Ser Leu Gln Val Arg Gly Arg His Ile Cys Gly Gly Ala Leu Ile Ala 585 590 595	1829

15/66

585	590	595	
gac cgc tgg gtg ata aca gct gcc cac tgc ttc cag gag gac agc atg Asp Arg Trp Val Ile Thr Ala Ala His Cys Phe Gln Glu Asp Ser Met 600 605 610 615			1877
gcc tcc acg gtg ctg tgg acc gtg ttc ctg ggc aag gtg tgg cag aac Ala Ser Thr Val Leu Trp Thr Val Phe Leu Gly Lys Val Trp Gln Asn 620 625 630			1925
tcg cgc tgg cct gga gag gtg tcc ttc aag gtg agc cgc ctg ctc ctg Ser Arg Trp Pro Gly Glu Val Ser Phe Lys Val Ser Arg Leu Leu Leu 635 640 645			1973
cac ccg tac cac gaa gag gac agc cat gac tac gac gtg gcg ctg ctg His Pro Tyr His Glu Glu Asp Ser His Asp Tyr Asp Val Ala Leu Leu 650 655 660			2021
cag ctc gac cac ccg gtg gtg cgc tcg gcc gcc gtg cgc ccc gtc tgc Gln Leu Asp His Pro Val Val Arg Ser Ala Ala Val Arg Pro Val Cys 665 670 675			2069
ctg ccc gcg cgc tcc cac ttc ttc gag ccc ggc ctg cac tgc tgg att Leu Pro Ala Arg Ser His Phe Phe Glu Pro Gly Leu His Cys Trp Ile 680 685 690 695			2117
acg ggc tgg ggc gcc ttg cgc gag ggc ggc ccc atc agc aac gct ctg Thr Gly Trp Gly Ala Leu Arg Glu Gly Gly Pro Ile Ser Asn Ala Leu 700 705 710			2165
cag aaa gtg gat gtg cag ttg atc cca cag gac ctg tgc agc gag gtc Gln Lys Val Asp Val Gln Leu Ile Pro Gln Asp Leu Cys Ser Glu Val 715 720 725			2213
tat cgc tac cag gtg acg cca cgc atg ctg tgt gcc ggc tac cgc aag Tyr Arg Tyr Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys 730 735 740			2261
ggc aag aag gat gcc tgt cag ggt gac tca ggt ggt ccg ctg gtg tgc Gly Lys Lys Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys 745 750 755			2309
aag gca ctc agt ggc cgc tgg ttc ctg gcg ggg ctg gtc agc tgg ggc Lys Ala Leu Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly 760 765 770 775			2357
ctg ggc tgt ggc cgg cct aac tac ttc ggc gtc tac acc cgc atc aca Leu Gly Cys Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr 780 785 790			2405
ggg gtg atc agc tgg atc cag caa gtg gtg acc tga ggaactgcc Gly Val Ile Ser Trp Ile Gln Gln Val Val Thr *			2451
ccctgcaaag cagggccac ctctggact cagagagccc agggcaactg ccaagcaggg ggacaagtat tctggcgggg ggtgggggag agagcaggcc ctgtggtggc aggaggggca tcttggttcg tccctgatgt ctgtccagta tggcaggagg atgagaagtg ccagcagttg ggggtaaca cgtcccttga ggacccaggc ccacacccag cccttttgcc tcccaattct ctctcctccg tcccttct ccaactgtgc ctaatgcaag gcagtggctc agcagcaaga atgctggttc tacatcccga ggagtgtctg aggtgcgccc cactctgtac agaggctgtt tgggcagcct tgccctccaga gagcagattc cagcttcgga agccctggg ctaacttggg atctgggaat ggaaggtgct cccatcggag gggaccctca gagccctgga gactgccagg			2511 2571 2631 2691 2751 2811 2871 2931

```

tgggcctgct gccactgtaa gccaaaaggt ggggaagtcc tgactccagg gtccttgccc 2991
caccctgcc tgccacctgg gccctcacag cccagaccct cactgggagg tgagctcagc 3051
tgccctttgg aataaagctg cctgatgcaa aaaaaaaaaa aaaaaaaaaa aaa 3104

```

<210> 8
 <211> 802
 <212> PRT
 <213> Homo Sapien

<400> 8
 Met Pro Val Ala Glu Ala Pro Gln Val Ala Gly Gly Gln Gly Asp Gly
 1 5 10 15
 Gly Asp Gly Glu Ala Glu Pro Glu Gly Met Phe Lys Ala Cys Glu
 20 25 30
 Asp Ser Lys Arg Lys Ala Arg Gly Tyr Leu Arg Leu Val Pro Leu Phe
 35 40 45
 Val Leu Leu Ala Leu Leu Val Leu Ala Ser Ala Gly Val Leu Leu Trp
 50 55 60
 Tyr Phe Leu Gly Tyr Lys Ala Glu Val Met Val Ser Gln Val Tyr Ser
 65 70 75 80
 Gly Ser Leu Arg Val Leu Asn Arg His Phe Ser Gln Asp Leu Thr Arg
 85 90 95
 Arg Glu Ser Ser Ala Phe Arg Ser Glu Thr Ala Lys Ala Gln Lys Met
 100 105 110
 Leu Lys Glu Leu Ile Thr Ser Thr Arg Leu Gly Thr Tyr Tyr Asn Ser
 115 120 125
 Ser Ser Val Tyr Ser Phe Gly Glu Gly Pro Leu Thr Cys Phe Phe Trp
 130 135 140
 Phe Ile Leu Gln Ile Pro Glu His Arg Arg Leu Met Leu Ser Pro Glu
 145 150 155 160
 Val Val Gln Ala Leu Leu Val Glu Glu Leu Leu Ser Thr Val Asn Ser
 165 170 175
 Ser Ala Ala Val Pro Tyr Arg Ala Glu Tyr Glu Val Asp Pro Glu Gly
 180 185 190
 Leu Val Ile Leu Glu Ala Ser Val Lys Asp Ile Ala Ala Leu Asn Ser
 195 200 205
 Thr Leu Gly Cys Tyr Arg Tyr Ser Tyr Val Gly Gln Gly Gln Val Leu
 210 215 220
 Arg Leu Lys Gly Pro Asp His Leu Ala Ser Ser Cys Leu Trp His Leu
 225 230 235 240
 Gln Gly Pro Lys Asp Leu Met Leu Lys Leu Arg Leu Glu Trp Thr Leu
 245 250 255
 Ala Glu Cys Arg Asp Arg Leu Ala Met Tyr Asp Val Ala Gly Pro Leu
 260 265 270
 Glu Lys Arg Leu Ile Thr Ser Val Tyr Gly Cys Ser Arg Gln Glu Pro
 275 280 285
 Val Val Glu Val Leu Ala Ser Gly Ala Ile Met Ala Val Val Trp Lys
 290 295 300
 Lys Gly Leu His Ser Tyr Tyr Asp Pro Phe Val Leu Ser Val Gln Pro
 305 310 315 320
 Val Val Phe Gln Ala Cys Glu Val Asn Leu Thr Leu Asp Asn Arg Leu
 325 330 335
 Asp Ser Gln Gly Val Leu Ser Thr Pro Tyr Phe Pro Ser Tyr Tyr Ser
 340 345 350
 Pro Gln Thr His Cys Ser Trp His Leu Thr Val Pro Ser Leu Asp Tyr
 355 360 365
 Gly Leu Ala Leu Trp Phe Asp Ala Tyr Ala Leu Arg Arg Gln Lys Tyr
 370 375 380
 Asp Leu Pro Cys Thr Gln Gly Gln Trp Thr Ile Gln Asn Arg Arg Leu
 385 390 395 400
 Cys Gly Leu Arg Ile Leu Gln Pro Tyr Ala Glu Arg Ile Pro Val Val
 405 410 415

Ala Thr Ala Gly Ile Thr Ile Asn Phe Thr Ser Gln Ile Ser Leu Thr
 Gly Pro Gly Val Arg Val His Tyr Gly Leu Tyr Asn Gln Ser Asp Pro
 Cys Pro Gly Glu Phe Leu Cys Ser Val Asn Gly Leu Cys Val Pro Ala
 Cys Asp Gly Val Lys Asp Cys Pro Asn Gly Leu Asp Glu Arg Asn Cys
 Val Cys Arg Ala Thr Phe Gln Cys Lys Glu Asp Ser Thr Cys Ile Ser
 Leu Pro Lys Val Cys Asp Gly Gln Pro Asp Cys Leu Asn Gly Ser Asp
 Glu Glu Gln Cys Gln Glu Gly Val Pro Cys Gly Thr Phe Thr Phe Gln
 Cys Glu Asp Arg Ser Cys Val Lys Lys Pro Asn Pro Gln Cys Asp Gly
 Arg Pro Asp Cys Arg Asp Gly Ser Asp Glu Glu His Cys Glu Cys Gly
 Leu Gln Gly Pro Ser Ser Arg Ile Val Gly Gly Ala Val Ser Ser Glu
 Gly Glu Trp Pro Trp Gln Ala Ser Leu Gln Val Arg Gly Arg His Ile
 Cys Gly Gly Ala Leu Ile Ala Asp Arg Trp Val Ile Thr Ala Ala His
 Cys Phe Gln Glu Asp Ser Met Ala Ser Thr Val Leu Trp Thr Val Phe
 Leu Gly Lys Val Trp Gln Asn Ser Arg Trp Pro Gly Glu Val Ser Phe
 Lys Val Ser Arg Leu Leu Leu His Pro Tyr His Glu Glu Asp Ser His
 Asp Tyr Asp Val Ala Leu Leu Gln Leu Asp His Pro Val Val Arg Ser
 Ala Ala Val Arg Pro Val Cys Leu Pro Ala Arg Ser His Phe Phe Glu
 Pro Gly Leu His Cys Trp Ile Thr Gly Trp Gly Ala Leu Arg Glu Gly
 Gly Pro Ile Ser Asn Ala Leu Gln Lys Val Asp Val Gln Leu Ile Pro
 Gln Asp Leu Cys Ser Glu Val Tyr Arg Tyr Gln Val Thr Pro Arg Met
 Leu Cys Ala Gly Tyr Arg Lys Gly Lys Lys Asp Ala Cys Gln Gly Asp
 Ser Gly Gly Pro Leu Val Cys Lys Ala Leu Ser Gly Arg Trp Phe Leu
 Ala Gly Leu Val Ser Trp Gly Leu Gly Cys Gly Arg Pro Asn Tyr Phe
 Gly Val Tyr Thr Arg Ile Thr Gly Val Ile Ser Trp Ile Gln Gln Val
 Val Thr

<210> 9

<211> 2672

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (33)...(2009)

<223> cDNA encoding: MTSP4-S (short form) splice variant

<400> 9

tcatacggcca gagggatgac agtgagcaga ag	atg ccc gtg gcc gag gcc ccc	53
Met Pro Val Ala Glu Ala Pro		
1 5		
cag gtg gct ggc ggg cag ggg gac gga ggt gat ggc gag gaa gcg gag	101	
Gln Val Ala Gly Gly Gln Gly Asp Gly Gly Asp Gly Glu Glu Ala Glu		
10 15 20		
ccg gag ggg atg ttc aag gcc tgt gag gac tcc aag aga aaa gcc cgg	149	
Pro Glu Gly Met Phe Lys Ala Cys Glu Asp Ser Lys Arg Lys Ala Arg		
25 30 35		
ggc tac ctc cgc ctg gtg ccc ctg ttt gtg ctg ctg gcc ctg ctc gtg	197	
Gly Tyr Leu Arg Leu Val Pro Leu Phe Val Leu Ala Leu Leu Val		
40 45 50 55		
ctg gct tcg gcg ggg gtg cta ctc tgg tat ttc cta ggg tac aag gcg	245	
Leu Ala Ser Ala Gly Val Leu Leu Trp Tyr Phe Leu Gly Tyr Lys Ala		
60 65 70		
gag gtg atg gtc agc cag gtg tac tca ggc agt ctg cgt gta ctc aat	293	
Glu Val Met Val Ser Gln Val Tyr Ser Gly Ser Leu Arg Val Leu Asn		
75 80 85		
cgc cac ttc tcc cag gat ctt acc cgc cgg gaa tct agt gcc ttc cgc	341	
Arg His Phe Ser Gln Asp Leu Thr Arg Arg Glu Ser Ser Ala Phe Arg		
90 95 100		
agt gaa acc gcc aaa gcc cag aag atg ctc aag gag ctc atc acc agc	389	
Ser Glu Thr Ala Lys Ala Gln Lys Met Leu Lys Glu Leu Ile Thr Ser		
105 110 115		
acc cgc ctg gga act tac tac aac tcc agc tcc gtc tat tcc ttt ggg	437	
Thr Arg Leu Gly Thr Tyr Tyr Asn Ser Ser Ser Val Tyr Ser Phe Gly		
120 125 130 135		
gtg tac ggc tgc agc cgc cag gag ccc gtg gtg gag gtt ctg gcg tcg	485	
Val Tyr Gly Cys Ser Arg Gln Glu Pro Val Val Glu Val Leu Ala Ser		
140 145 150		
ggg gcc atc atg gcg gtc gtc tgg aag aag ggc ctg cac agc tac tac	533	
Gly Ala Ile Met Ala Val Val Trp Lys Lys Gly Leu His Ser Tyr Tyr		
155 160 165		
gac ccc ttc gtg ctc tcc gtg cag ccg gtg gtc ttc cag gcc tgt gaa	581	
Asp Pro Phe Val Leu Ser Val Gln Pro Val Val Phe Gln Ala Cys Glu		
170 175 180		
gtg aac ctg acg ctg gac aac agg ctc gac tcc cag ggc gtc ctc agc	629	
Val Asn Leu Thr Leu Asp Asn Arg Leu Asp Ser Gln Gly Val Leu Ser		
185 190 195		
acc ccg tac ttc ccc agc tac tac tcg ccc caa acc cac tgc tcc tgg	677	
Thr Pro Tyr Phe Pro Ser Tyr Tyr Ser Pro Gln Thr His Cys Ser Trp		
200 205 210 215		
cac ctc acg gtg ccc tct ctg gac tac ggc ttg gcc ctc tgg ttt gat	725	
His Leu Thr Val Pro Ser Leu Asp Tyr Gly Leu Ala Leu Trp Phe Asp		
220 225 230		
gcc tat gca ctg agg agg cag aag tat gat ttg ccg tgc acc cag ggc	773	
Ala Tyr Ala Leu Arg Arg Gln Lys Tyr Asp Leu Pro Cys Thr Gln Gly		

235				240				245								
cag Gln	tgg Trp	acg Thr 250	atc Ile	cag Gln	aac Asn	agg Arg	agg Arg 255	ctg Leu	tgt Cys	ggc Gly	ttg Leu	cgc Arg 260	atc Ile	ctg Leu	cag Gln	821
ccc Pro	tac Tyr 265	gcc Ala	gag Glu	agg Arg	atc Ile	ccc Pro 270	gtg Val	gtg Val	gcc Ala	acg Thr	gcc Ala 275	ggg Gly	atc Ile	acc Thr	atc Ile	869
aac Asn 280	ttc Phe	acc Thr	tcc Ser	cag Gln	atc Ile 285	tcc Ser	ctc Leu	acc Thr	ggg Gly	ccc Pro 290	ggg Gly	gtg Val	cgg Arg	gtg Val	cac His 295	917
tat Tyr	ggc Gly	ttg Leu	tac Tyr	aac Asn 300	cag Gln	tcg Ser	gac Asp	ccc Pro	tgc Cys 305	cct Pro	gga Gly	gag Glu	ttc Phe	ctc Leu 310	tgt Cys	965
tct Ser	gtg Val	aat Asn	gga Gly 315	ctc Leu	tgt Cys	gtc Val	cct Pro	gcc Ala 320	tgt Cys	gat Asp	ggg Gly	gtc Val	aag Lys 325	gac Asp	tgc Cys	1013
ccc Pro	aac Asn	ggc Gly 330	ctg Leu	gat Asp	gag Glu	aga Arg	aac Asn 335	tgc Cys	gtt Val	tgc Cys	aga Arg	gcc Ala 340	aca Thr	ttc Phe	cag Gln	1061
tgc Cys	aaa Lys 345	gag Glu	gac Asp	agc Ser	aca Thr	tgc Cys 350	atc Ile	tca Ser	ctg Leu	ccc Pro	aag Lys 355	gtc Val	tgt Cys	gat Asp	ggg Gly	1109
cag Gln 360	cct Pro	gat Asp	tgt Cys	ctc Leu	aac Asn 365	ggc Gly	agc Ser	gac Asp	gaa Glu	gag Glu 370	cag Gln	tgc Cys	cag Gln	gaa Glu	ggg Gly 375	1157
gtg Val	cca Pro	tgt Cys	ggg Gly	aca Thr 380	ttc Phe	acc Thr	ttc Phe	cag Gln	tgt Cys 385	gag Glu	gac Asp	cgg Arg	agc Ser	tgc Cys 390	gtg Val	1205
aag Lys	aag Lys	ccc Pro	aac Asn 395	ccg Pro	cag Gln	tgt Cys	gat Asp	ggg Gly 400	cgg Arg	ccc Pro	gac Asp	tgc Cys	agg Arg 405	gac Asp	ggc Gly	1253
tcg Ser	gat Asp	gag Glu 410	gag Glu	cac His	tgt Cys	gaa Glu	tgt Cys 415	ggc Gly	ctc Leu	cag Gln	ggc Gly	ccc Pro 420	tcc Ser	agc Ser	cgc Arg	1301
att Ile	gtt Val 425	ggg Gly	gga Gly	gct Ala	gtg Val 430	tcc Ser	tcc Ser	gag Glu	ggg Gly	gag Glu	tgg Trp 435	cca Pro	tgg Trp	cag Gln	gcc Ala	1349
agc Ser 440	ctc Leu	cag Gln	gtt Val	cgg Arg	ggg Gly 445	cga Arg	cac His	atc Ile	tgt Cys	ggg Gly 450	ggg Gly	gcc Ala	ctc Leu	atc Ile	gct Ala 455	1397
gac Asp	cgc Arg	tgg Trp	gtg Val	ata Ile 460	aca Thr	gct Ala	gcc Ala	cac His	tgc Cys 465	ttc Phe	cag Gln	gag Glu	gac Asp 470	agc Ser	atg Met	1445
gcc Ala	tcc Ser	acg Thr 475	gtg Val	ctg Leu	tgg Trp	acc Thr	gtg Val	ttc Phe 480	ctg Leu	ggc Gly	aag Lys	gtg Val	tgg Trp 485	cag Gln	aac Asn	1493

20/66

tcg cgc tgg cct gga gag gtg tcc ttc aag gtg agc cgc ctg ctc ctg Ser Arg Trp Pro Gly Glu Val Ser Phe Lys Val Ser Arg Leu Leu Leu 490 495 500	1541
cac ccg tac cac gaa gag gac agc cat gac tac gac gtg gcg ctg ctg His Pro Tyr His Glu Glu Asp Ser His Asp Tyr Asp Val Ala Leu Leu 505 510 515	1589
cag ctc gac cac ccg gtg gtg cgc tgc gcc gcc gtg cgc ccc gtc tgc Gln Leu Asp His Pro Val Val Arg Ser Ala Ala Val Arg Pro Val Cys 520 525 530 535	1637
ctg ccc gcg cgc tcc cac ttc ttc gag ccc gcc ctg cac tgc tgg att Leu Pro Ala Arg Ser His Phe Phe Glu Pro Gly Leu His Cys Trp Ile 540 545 550	1685
acg gcc tgg gcc gcc ttg cgc gag gcc gcc ccc atc agc aac gct ctg Thr Gly Trp Gly Ala Leu Arg Glu Gly Gly Pro Ile Ser Asn Ala Leu 555 560 565	1733
cag aaa gtg gat gtg cag ttg atc cca cag gac ctg tgc agc gag gtc Gln Lys Val Asp Val Gln Leu Ile Pro Gln Asp Leu Cys Ser Glu Val 570 575 580	1781
tat cgc tac cag gtg acg cca cgc atg ctg tgt gcc gcc tac cgc aag Tyr Arg Tyr Gln Val Thr Pro Arg Met Leu Cys Ala Gly Tyr Arg Lys 585 590 595	1829
ggc aag aag gat gcc tgt cag ggt gac tca ggt ggt ccg ctg gtg tgc Gly Lys Lys Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro Leu Val Cys 600 605 610 615	1877
aag gca ctc agt gcc cgc tgg ttc ctg gcg ggg ctg gtc agc tgg gcc Lys Ala Leu Ser Gly Arg Trp Phe Leu Ala Gly Leu Val Ser Trp Gly 620 625 630	1925
ctg gcc tgt gcc cgg cct aac tac ttc gcc gtc tac acc cgc atc aca Leu Gly Cys Gly Arg Pro Asn Tyr Phe Gly Val Tyr Thr Arg Ile Thr 635 640 645	1973
ggt gtg atc agc tgg atc cag caa gtg gtg acc tga ggaactgccc Gly Val Ile Ser Trp Ile Gln Gln Val Val Thr * 650 655	2019
ccctgcaaag cagggccac ctctggact cagagagccc agggcaactg ccaagcaggg ggacaagtat tctggcgggg ggtgggggag agagcaggcc ctgtggtggc aggaggggca tcttgtttcg tccctgatgt ctgtccagta tggcaggagg atgagaagtg ccagcagttg ggggtaaca cgtcccttga ggaccacagg ccacacccag cccttttgcc tcccaattct ctctcctccg tcccttccct ccaactgctgc ctaatgcaag gcagtggctc agcagcaaga atgctgggttc tacatcccga ggagtgtctg aggtgcgccc cactctgtac agaggctgtt tgggcagcct tgccctccaga gacgagattc cagcttcgga agcccctggt ctaacttggg atctgggaat ggaagggtgct cccatcgagg gggaccctca gagecctgga gactgccagg tgggctgtg gccactgtaa gccaaaaggt ggggaagtcc tgactccagg gtccttgccc caccctgcc tgccacctgg gccctcacag ccagagacct cactgggagg tgagctcagg tgccctttgg aataaagctg cctgatgcaa aaaaaaaaaa aaaaaaaaaa aaa	2079 2139 2199 2259 2319 2379 2439 2499 2559 2619 2672

<210> 10

<211> 658

<212> PRT

<213> Homo Sapien

<400> 10

21/66

Met	Pro	Val	Ala	Glu	Ala	Pro	Gln	Val	Ala	Gly	Gly	Gln	Gly	Asp	Gly
1				5					10					15	
Gly	Asp	Gly	Glu	Glu	Ala	Glu	Pro	Glu	Gly	Met	Phe	Lys	Ala	Cys	Glu
			20					25					30		
Asp	Ser	Lys	Arg	Lys	Ala	Arg	Gly	Tyr	Leu	Arg	Leu	Val	Pro	Leu	Phe
		35					40					45			
Val	Leu	Leu	Ala	Leu	Leu	Val	Leu	Ala	Ser	Ala	Gly	Val	Leu	Leu	Trp
	50					55					60				
Tyr	Phe	Leu	Gly	Tyr	Lys	Ala	Glu	Val	Met	Val	Ser	Gln	Val	Tyr	Ser
65					70				75					80	
Gly	Ser	Leu	Arg	Val	Leu	Asn	Arg	His	Phe	Ser	Gln	Asp	Leu	Thr	Arg
			85						90					95	
Arg	Glu	Ser	Ser	Ala	Phe	Arg	Ser	Glu	Thr	Ala	Lys	Ala	Gln	Lys	Met
			100					105					110		
Leu	Lys	Glu	Leu	Ile	Thr	Ser	Thr	Arg	Leu	Gly	Thr	Tyr	Tyr	Asn	Ser
		115					120					125			
Ser	Ser	Val	Tyr	Ser	Phe	Gly	Val	Tyr	Gly	Cys	Ser	Arg	Gln	Glu	Pro
		130				135					140				
Val	Val	Glu	Val	Leu	Ala	Ser	Gly	Ala	Ile	Met	Ala	Val	Val	Trp	Lys
145					150				155					160	
Lys	Gly	Leu	His	Ser	Tyr	Tyr	Asp	Pro	Phe	Val	Leu	Ser	Val	Gln	Pro
			165						170					175	
Val	Val	Phe	Gln	Ala	Cys	Glu	Val	Asn	Leu	Thr	Leu	Asp	Asn	Arg	Leu
		180						185					190		
Asp	Ser	Gln	Gly	Val	Leu	Ser	Thr	Pro	Tyr	Phe	Pro	Ser	Tyr	Tyr	Ser
		195					200					205			
Pro	Gln	Thr	His	Cys	Ser	Trp	His	Leu	Thr	Val	Pro	Ser	Leu	Asp	Tyr
		210				215					220				
Gly	Leu	Ala	Leu	Trp	Phe	Asp	Ala	Tyr	Ala	Leu	Arg	Arg	Gln	Lys	Tyr
225					230				235					240	
Asp	Leu	Pro	Cys	Thr	Gln	Gly	Gln	Trp	Thr	Ile	Gln	Asn	Arg	Arg	Leu
			245						250					255	
Cys	Gly	Leu	Arg	Ile	Leu	Gln	Pro	Tyr	Ala	Glu	Arg	Ile	Pro	Val	Val
		260						265					270		
Ala	Thr	Ala	Gly	Ile	Thr	Ile	Asn	Phe	Thr	Ser	Gln	Ile	Ser	Leu	Thr
		275					280					285			
Gly	Pro	Gly	Val	Arg	Val	His	Tyr	Gly	Leu	Tyr	Asn	Gln	Ser	Asp	Pro
		290				295					300				
Cys	Pro	Gly	Glu	Phe	Leu	Cys	Ser	Val	Asn	Gly	Leu	Cys	Val	Pro	Ala
305					310				315					320	
Cys	Asp	Gly	Val	Lys	Asp	Cys	Pro	Asn	Gly	Leu	Asp	Glu	Arg	Asn	Cys
			325						330					335	
Val	Cys	Arg	Ala	Thr	Phe	Gln	Cys	Lys	Glu	Asp	Ser	Thr	Cys	Ile	Ser
		340						345					350		
Leu	Pro	Lys	Val	Cys	Asp	Gly	Gln	Pro	Asp	Cys	Leu	Asn	Gly	Ser	Asp
		355					360					365			
Glu	Glu	Gln	Cys	Gln	Glu	Gly	Val	Pro	Cys	Gly	Thr	Phe	Thr	Phe	Gln
	370					375					380				
Cys	Glu	Asp	Arg	Ser	Cys	Val	Lys	Lys	Pro	Asn	Pro	Gln	Cys	Asp	Gly
385					390				395					400	
Arg	Pro	Asp	Cys	Arg	Asp	Gly	Ser	Asp	Glu	Glu	His	Cys	Glu	Cys	Gly
			405						410					415	
Leu	Gln	Gly	Pro	Ser	Ser	Arg	Ile	Val	Gly	Gly	Ala	Val	Ser	Ser	Glu
		420						425					430		
Gly	Glu	Trp	Pro	Trp	Gln	Ala	Ser	Leu	Gln	Val	Arg	Gly	Arg	His	Ile
		435					440					445			
Cys	Gly	Gly	Ala	Leu	Ile	Ala	Asp	Arg	Trp	Val	Ile	Thr	Ala	Ala	His
	450					455					460				
Cys	Phe	Gln	Glu	Asp	Ser	Met	Ala	Ser	Thr	Val	Leu	Trp	Thr	Val	Phe
465					470				475					480	
Leu	Gly	Lys	Val	Trp	Gln	Asn	Ser	Arg	Trp	Pro	Gly	Glu	Val	Ser	Phe
			485						490					495	

22/66

Lys Val Ser Arg Leu Leu Leu His Pro Tyr His Glu Glu Asp Ser His
 500 505 510
 Asp Tyr Asp Val Ala Leu Leu Gln Leu Asp His Pro Val Val Arg Ser
 515 520 525
 Ala Ala Val Arg Pro Val Cys Leu Pro Ala Arg Ser His Phe Phe Glu
 530 535 540
 Pro Gly Leu His Cys Trp Ile Thr Gly Trp Gly Ala Leu Arg Glu Gly
 545 550 555 560
 Gly Pro Ile Ser Asn Ala Leu Gln Lys Val Asp Val Gln Leu Ile Pro
 565 570 575
 Gln Asp Leu Cys Ser Glu Val Tyr Arg Tyr Gln Val Thr Pro Arg Met
 580 585 590
 Leu Cys Ala Gly Tyr Arg Lys Gly Lys Lys Asp Ala Cys Gln Gly Asp
 595 600 605
 Ser Gly Gly Pro Leu Val Cys Lys Ala Leu Ser Gly Arg Trp Phe Leu
 610 615 620
 Ala Gly Leu Val Ser Trp Gly Leu Gly Cys Gly Arg Pro Asn Tyr Phe
 625 630 635 640
 Gly Val Tyr Thr Arg Ile Thr Gly Val Ile Ser Trp Ile Gln Gln Val
 645 650 655
 Val Thr

<210> 11
 <211> 1656
 <212> DNA
 <213> Homo Sapien

<220>
 <221> CDS
 <222> (268) ... (1629)
 <223> DNA sequence encoding a transmembrane serine
 protease (MTSP-6) protein

<400> 11
 cgcccgggca ggtcagtaac actgtggcct actatctctt ccgtgggtgcc atctacattt 60
 ttgggactcg ggaattatga ctgtttttgg ttaatcgata ctgaatgcgc tttgtgtgga 120
 ctgtcgaatt tcaaagattt accgtatgac caagatgcac ctgatgctac aagtataaat 180
 aggggaacaa atgctttctg ttcttctctg gctaaggagg tagaggtgga ggcggagccg 240
 gatgtcagag gtcctgaaat agtcacc atg ggg gaa aat gat ccg cct gct gtt 294
 Met Gly Glu Asn Asp Pro Pro Ala Val
 1 5

 gaa gcc ccc ttc tca ttc cga tcg ctt ttt ggc ctt gat gat ttg aaa 342
 Glu Ala Pro Phe Ser Phe Arg Ser Leu Phe Gly Leu Asp Asp Leu Lys
 10 15 20 25

 ata agt cct gtt gca cca gat gca gat gct gtt gct gca cag atc ctg 390
 Ile Ser Pro Val Ala Pro Asp Ala Asp Ala Val Ala Ala Gln Ile Leu
 30 35 40

 tca ctg ctg cca ttg aag ttt ttt cca atc atc gtc att ggg atc att 438
 Ser Leu Leu Pro Leu Lys Phe Phe Pro Ile Ile Val Ile Gly Ile Ile
 45 50 55

 gca ttg ata tta gca ctg gcc att ggt ctg ggc atc cac ttc gac tgc 486
 Ala Leu Ile Leu Ala Leu Ala Ile Gly Leu Gly Ile His Phe Asp Cys
 60 65 70

 tca ggg aag tac aga tgt cgc tca tcc ttt aag tgt atc gag ctg ata 534
 Ser Gly Lys Tyr Arg Cys Arg Ser Ser Ser Lys Cys Ile Glu Leu Ile
 75 80 85

gct cga tgt gac gga gtc tcg gat tgc aaa gac ggg gag gac gag tac	582
Ala Arg Cys Asp Gly Val Ser Asp Cys Lys Asp Gly Glu Asp Glu Tyr	
90 95 100 105	
cgc tgt gtc cgg gtg ggt ggt cag aat gcc gtg ctc cag gtg ttc aca	630
Arg Cys Val Arg Val Gly Gly Gln Asn Ala Val Leu Gln Val Phe Thr	
110 115 120	
gct gct tcg tgg aag acc atg tgc tcc gat gac tgg aag ggt cac tac	678
Ala Ala Ser Trp Lys Thr Met Cys Ser Asp Asp Trp Lys Gly His Tyr	
125 130 135	
gca aat gtt gcc tgt gcc caa ctg ggt ttc cca agc tat gta agt tca	726
Ala Asn Val Ala Cys Ala Gln Leu Gly Phe Pro Ser Tyr Val Ser Ser	
140 145 150	
gat aac ctc aga gtg agc tcg cta gag ggg cag ttc cgg gag gag ttt	774
Asp Asn Leu Arg Val Ser Ser Leu Glu Gly Gln Phe Arg Glu Glu Phe	
155 160 165	
gtg tcc atc gat cac ctc ttg cca gat gac aag gtg act gca tta cac	822
Val Ser Ile Asp His Leu Leu Pro Asp Asp Lys Val Thr Ala Leu His	
170 175 180 185	
cac tca gta tat gtg agg gag gga tgt gcc tct ggc cac gtg gtt acc	870
His Ser Val Tyr Val Arg Glu Gly Cys Ala Ser Gly His Val Val Thr	
190 195 200	
ttg cag tgc aca gcc tgt ggt cat aga agg ggc tac agc tca cgc atc	918
Leu Gln Cys Thr Ala Cys Gly His Arg Arg Gly Tyr Ser Ser Arg Ile	
205 210 215	
gtg ggt gga aac atg tcc ttg ctc tcg cag tgg ccc tgg cag gcc agc	966
Val Gly Gly Asn Met Ser Leu Leu Ser Gln Trp Pro Trp Gln Ala Ser	
220 225 230	
ctt cag ttc cag ggc tac cac ctg tgc ggg ggc tct gtc atc acg ccc	1014
Leu Gln Phe Gln Gly Tyr His Leu Cys Gly Gly Ser Val Ile Thr Pro	
235 240 245	
ctg tgg atc atc act gct gca cac tgt gtt tat gac ttg tac ctc ccc	1062
Leu Trp Ile Ile Thr Ala Ala His Cys Val Tyr Asp Leu Tyr Leu Pro	
250 255 260 265	
aag tca tgg acc atc cag gtg ggt cta gtt tcc ctg ttg gac aat cca	1110
Lys Ser Trp Thr Ile Gln Val Gly Leu Val Ser Leu Leu Asp Asn Pro	
270 275 280	
gcc cca tcc cac ttg gtg gag aag att gtc tac cac agc aag tac aag	1158
Ala Pro Ser His Leu Val Glu Lys Ile Val Tyr His Ser Lys Tyr Lys	
285 290 295	
cca aag agg ctg ggc aat gac atc gcc ctt atg aag ctg gcc ggg cca	1206
Pro Lys Arg Leu Gly Asn Asp Ile Ala Leu Met Lys Leu Ala Gly Pro	
300 305 310	
ctc acg ttc aat gaa atg atc cag cct gtg tgc ctg ccc aac tct gaa	1254
Leu Thr Phe Asn Glu Met Ile Gln Pro Val Cys Leu Pro Asn Ser Glu	
315 320 325	
gag aac ttc ccc gat gga aaa gtg tgc tgg acg tca gga tgg ggg gcc	1302

24/66

Glu Asn Phe Pro Asp Gly Lys Val Cys Trp Thr Ser Gly Trp Gly Ala
 330 335 340 345
 aca gag gat gga ggt gac gcc tcc cct gtc ctg aac cac gcg gcc gtc 1350
 Thr Glu Asp Gly Gly Asp Ala Ser Pro Val Leu Asn His Ala Ala Val 350 355 360
 cct ttg att tcc aac aag atc tgc aac cac agg gac gtg tac ggt ggc 1398
 Pro Leu Ile Ser Asn Lys Ile Cys Asn His Arg Asp Val Tyr Gly Gly 365 370 375
 atc atc tcc ccc tcc atg ctc tgc gcg ggc tac ctg acg ggt ggc gtg 1446
 Ile Ile Ser Pro Ser Met Leu Cys Ala Gly Tyr Leu Thr Gly Gly Val 380 385 390
 gac agc tgc cag ggg gac agc ggg ggg ccc ctg gtg tgt caa gag agg 1494
 Asp Ser Cys Gln Gly Asp Gly Gly Pro Leu Val Cys Gln Glu Arg 395 400 405
 agg ctg tgg aag tta gtg gga gcg acc agc ttt ggc atc ggc tgc gca 1542
 Arg Leu Trp Lys Leu Val Gly Ala Thr Ser Phe Gly Ile Gly Cys Ala 410 415 420 425
 gag gtg aac aag cct ggg gtg tac acc cgt gtc acc tcc ttc ctg gac 1590
 Glu Val Asn Lys Pro Gly Val Tyr Thr Arg Val Thr Ser Phe Leu Asp 430 435 440
 tgg atc cac gag cag atg gag aga gac cta aaa acc tga agaggaaggg 1639
 Trp Ile His Glu Gln Met Glu Arg Asp Leu Lys Thr * 445 450
 gataagtagc cacctga 1656
 <210> 12
 <211> 453
 <212> PRT
 <213> Homo Sapien
 <400> 12
 Met Gly Glu Asn Asp Pro Pro Ala Val Glu Ala Pro Phe Ser Phe Arg
 1 5 10 15
 Ser Leu Phe Gly Leu Asp Asp Leu Lys Ile Ser Pro Val Ala Pro Asp
 20 25 30
 Ala Asp Ala Val Ala Ala Gln Ile Leu Ser Leu Leu Pro Leu Lys Phe
 35 40 45
 Phe Pro Ile Ile Val Ile Gly Ile Ile Ala Leu Ile Leu Ala Leu Ala
 50 55 60
 Ile Gly Leu Gly Ile His Phe Asp Cys Ser Gly Lys Tyr Arg Cys Arg
 65 70 75 80
 Ser Ser Phe Lys Cys Ile Glu Leu Ile Ala Arg Cys Asp Gly Val Ser
 85 90 95
 Asp Cys Lys Asp Gly Glu Asp Glu Tyr Arg Cys Val Arg Val Gly Gly
 100 105 110
 Gln Asn Ala Val Leu Gln Val Phe Thr Ala Ala Ser Trp Lys Thr Met
 115 120 125
 Cys Ser Asp Asp Trp Lys Gly His Tyr Ala Asn Val Ala Cys Ala Gln
 130 135 140
 Leu Gly Phe Pro Ser Tyr Val Ser Ser Asp Asn Leu Arg Val Ser Ser
 145 150 155 160
 Leu Glu Gly Gln Phe Arg Glu Glu Phe Val Ser Ile Asp His Leu Leu
 165 170 175
 Pro Asp Asp Lys Val Thr Ala Leu His His Ser Val Tyr Val Arg Glu

25/66

180				185				190							
Gly	Cys	Ala	Ser	Gly	His	Val	Val	Thr	Leu	Gln	Cys	Thr	Ala	Cys	Gly
		195					200					205			
His	Arg	Arg	Gly	Tyr	Ser	Ser	Arg	Ile	Val	Gly	Gly	Asn	Met	Ser	Leu
		210				215					220				
Leu	Ser	Gln	Trp	Pro	Trp	Gln	Ala	Ser	Leu	Gln	Phe	Gln	Gly	Tyr	His
225				230						235				240	
Leu	Cys	Gly	Gly	Ser	Val	Ile	Thr	Pro	Leu	Trp	Ile	Ile	Thr	Ala	Ala
				245				250						255	
His	Cys	Val	Tyr	Asp	Leu	Tyr	Leu	Pro	Lys	Ser	Trp	Thr	Ile	Gln	Val
			260					265						270	
Gly	Leu	Val	Ser	Leu	Leu	Asp	Asn	Pro	Ala	Pro	Ser	His	Leu	Val	Glu
		275					280					285			
Lys	Ile	Val	Tyr	His	Ser	Lys	Tyr	Lys	Pro	Lys	Arg	Leu	Gly	Asn	Asp
		290				295					300				
Ile	Ala	Leu	Met	Lys	Leu	Ala	Gly	Pro	Leu	Thr	Phe	Asn	Glu	Met	Ile
305				310						315					320
Gln	Pro	Val	Cys	Leu	Pro	Asn	Ser	Glu	Glu	Asn	Phe	Pro	Asp	Gly	Lys
				325				330						335	
Val	Cys	Trp	Thr	Ser	Gly	Trp	Gly	Ala	Thr	Glu	Asp	Gly	Gly	Asp	Ala
			340					345						350	
Ser	Pro	Val	Leu	Asn	His	Ala	Ala	Val	Pro	Leu	Ile	Ser	Asn	Lys	Ile
		355					360					365			
Cys	Asn	His	Arg	Asp	Val	Tyr	Gly	Gly	Ile	Ile	Ser	Pro	Ser	Met	Leu
		370				375					380				
Cys	Ala	Gly	Tyr	Leu	Thr	Gly	Gly	Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser
385				390						395					400
Gly	Gly	Pro	Leu	Val	Cys	Gln	Glu	Arg	Arg	Leu	Trp	Lys	Leu	Val	Gly
				405				410						415	
Ala	Thr	Ser	Phe	Gly	Ile	Gly	Cys	Ala	Glu	Val	Asn	Lys	Pro	Gly	Val
			420					425						430	
Tyr	Thr	Arg	Val	Thr	Ser	Phe	Leu	Asp	Trp	Ile	His	Glu	Gln	Met	Glu
		435					440					445			
Arg	Asp	Leu	Lys	Thr											
	450														

```
<210> 13
<211> 23
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Oligonucleotide Primer

```
<221> misc_feature
<222> (0) .. (0)
<223> N= Inosine
```

<400> 13
tggtrtnvtnw sngcnrcnca ytg

23

```
<210> 14
<211> 30
<212> DNA
<213> Artificial Sequence
```

<220>
<223> Oligonucleotide Primer

```
<221> misc_feature
<222> (0) ... (0)
<223> N= Inosine
```

<400> 14
nggnccnccn swrtcnccyt nrcanghrtc 30

<210> 15
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 15
tcaâcgcgagaa gatgatgtgt gcaggcatcc 30

<210> 16
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 16
gggacagggg ctgtaaggca gggaatgag 29

<210> 17
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 17
cccgcagcca tagccccagc taacg 25

<210> 18
<211> 27
<212> DNA
<213> Artificial Sequence

<400> 18
gcagacgatg cgtaccaggg ggaagtc 27

<210> 19
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 19
ctcgagaaaa gagtgggtggg tggggaggag gcctctgtg 39

<210> 20
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 20
gcggccgcat tacagctcag ccttccagac 30
<210> 21
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<223> Oligonucleotide Primer
<400> 21
cctccacggt gctgtggacc gtgttcc 27
<210> 22
<211> 24
<212> DNA
<213> Artificial Sequence
<220>
<223> Oligonucleotide Primer
<400> 22
cctcgcgcaa ggcgccccag cccg 24
<210> 23
<211> 32
<212> DNA
<213> Artificial Sequence
<220>
<223> Oligonucleotide Primer
<400> 23
gcgtggcgtc acctggtagc gatagacctc gc 32
<210> 24
<211> 27
<212> DNA
<213> Artificial Sequence
<220>
<223> Oligonucleotide Primer
<400> 24
cctccacggt gctgtggacc gtgttcc 27
<210> 25
<211> 24
<212> DNA
<213> Artificial Sequence
<220>
<223> Oligonucleotide Primer
<400> 25
cctcgcgcaa ggcgccccag cccg 24
<210> 26
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 26
tcatcgcca gaggtgatc agtgag 26

<210> 27
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 27
cctcctcagt gcataggcat caaaccag 28

<210> 28
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 28
tctctcgaga aaagaattgt tggaggagct gtgtcctccg ag 42

<210> 29
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 29
aggtgggcct tgctttgcag gggggcagtt c 31

<210> 30
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 30
tcacgcatcg tgggtggaac atgtcc 26

<210> 31
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 31
accacctcc atctgctcgt ggatcc 26

<210> 32

<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 32
ccacagcctc ctctcttgac acaccag 27

<210> 33
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 33
acgcccctgt ggatcatcac tgctgc 26

<210> 34
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 34
tccctccctc acatatactg agtggtg 27

<210> 35
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 35
cgactgctca gggaagtcag atgtcg 26

<210> 36
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 36
gcggccgcac tataccccag tggtctcttt gatcca 36

<210> 37
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 37
ctggtgtgtc aagagaggag gctgtgg 27

<210> 38
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide primer

<400> 38
actcaggtgg ctacttatcc ccttcctc 28

<210> 39
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 39
tctctcgaga aaagagtggg ggggtggggag gaggcctctg tg 42

<210> 40
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 40
attcgcggcc gcattacagc tcagccttcc agac 34

<210> 41
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 41
tctctcgaga aaagaattgt tgggtggagct gtgtcctccg ag 42

<210> 42
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 42
attcgcggcc gctcaggtca ccacttgtcg gatccag 37

<210> 43
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Oligonucleotide Primer

<400> 43

ctcgagaaac gcacgtggg tggaaacatg tccttg

36

<210> 44

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide Primer

<400> 44

actcaggtgg ctacttatcc ccttcctc

28

<210> 45

<211> 9276

<212> DNA

<213> Pichia pastoris

<400> 45

agatctaaca	tccaaagacg	aaagggtgaa	tgaacacctt	ttgccatccg	acatccacag	60
gtccattctc	acacataagt	gccaaacgca	acaggagggg	atacactagc	agcagaccgt	120
tgcaaacgca	ggacctccac	tcctcttctc	ctcaacaccc	acttttgcca	tcgaaaaacc	180
agcccagtta	ttgggcttga	ttggagctcg	ctcattccaa	ttccttctat	taggctacta	240
acaccatgac	tttattagcc	tgtctatcct	ggccccctcg	gcgaggttca	tgtttgttta	300
tttccgaatg	caacaagctc	cgcattacac	ccgaacatca	ctccagatga	gggctttctg	360
agtgtggggg	caaatagttt	catgttcccc	aaatggccca	aaactgacag	tttaaaccgt	420
gtcttgggaac	ctaataatgac	aaaagcgtga	tctcatccaa	gatgaactaa	gtttgggttcg	480
ttgaaatgct	aacggccagt	tggtcaaaaa	gaaacttcca	aaagtcgcca	taccgtttgt	540
cttgtttggg	attgattgac	gaatgctcaa	aaataatctc	attaatgctt	agcgcagctc	600
ctctatcgct	tctgaacccc	gggtgcacctg	tgccgaaacg	caaatgggga	aacacccgct	660
ttttggatga	ttatgcattg	tctccacatt	gtatgcttcc	aagattctgg	tgggaatact	720
gctgatagcc	taacgttcat	gatcaaaatt	taactgttct	aacccctact	tgacagcaat	780
atataaacag	aaggaagctg	ccctgtctta	aacctttttt	tttatcatca	ttattagctt	840
actttcataa	ttgcgactgg	ttccaattga	caagcttttg	attttaacga	cttttaacga	900
caacttgaga	agatcaaaaa	acaactaatt	attcgagga	tccaaacgat	gagatttctc	960
tcaattttta	ctgcagtttt	attcgcagca	tcctccgcat	tagctgctcc	agtcaaacact	1020
acaacagaag	atgaaacggc	acaaattccg	gctgaagctg	tcacgggtta	ctcagattta	1080
gaaggggatt	tcgatgttgc	tgttttgcca	ttttccaaca	gcacaaataa	cgggttattg	1140
tttataaata	ctactattgc	cagcattgct	gctaagaag	aaggggtatc	tctcgagaaa	1200
agagaggctg	aagcttacgt	agaattccct	agggcgggcg	cgaatttaatt	cgccttagac	1260
atgactgttc	ctcagttcaa	gttgggcact	tacgagaaga	ccggtcttgc	tagattctaa	1320
tcaagaggat	gtcagaatgc	catttgcctg	agagatgcag	gcttcatttt	tgatactttt	1380
ttatttgtaa	cctatatagt	ataggatttt	ttttgtcatt	ttgtttcttc	tctgacgagc	1440
ttgctcctga	tcagcctatc	tgcagctga	tgaatatctt	gtggtagggg	tttgggaaaa	1500
tcattcgagt	ttgatgtttt	tcttggtatt	tcccactcct	cttcagagta	cagaagatta	1560
agtgagaagt	tcgtttgtgc	aagcttatcg	ataagcttta	atgcggtagt	ttatcacagt	1620
taaattgcta	acgcagtcag	gcaccgtgta	tgaatctaa	caatgcgctc	atcgtcatcc	1680
tcggcaccgt	caccctggat	gctgtaggca	taggcttggg	tatgccggta	ctgccggggc	1740
tcttgcggga	tatcgtccat	tccgacagca	tcgccagtc	ctatggcggtg	ctgctagcgc	1800
tatatgcgtt	gatgcaattt	ctatgcgcac	ccgttctcgg	agcactgtcc	gaccgctttg	1860
gccgcggccc	agtcctgctc	gcttcgctac	ttggagccac	tatcgactac	gcgatcatgg	1920
cgaccacacc	cgtcctgtgg	atctatcgaa	tctaaatgta	agttaaaatc	tctaaataat	1980
taataaagtc	ccagtttctc	catacgaacc	ttaacagcat	tcggtgagc	atctagacct	2040
tcaacagcag	ccagatctgg	cactgcttgg	ccaatatgtt	tcagtccctc	aggagttacg	2100
tcttgtgaag	tgatgaaactt	ctggaagggt	gcagtgttaa	ctccgctgta	ttgacgggca	2160
tatccgtacg	ttggcaaaagt	gtggttggtg	ccggaggagt	aatctccaca	actctctgga	2220
gagttaggcac	caacaaacac	agatccagcg	tgttgtactt	gatcaacata	agaagaagca	2280
ttctcgattt	gcaggatcaa	gtgttcagga	gcgtactgat	tggacatttc	caaagcctgc	2340

tcgtagggtt	caaccgatag	ggttgtagag	tgtgcaatac	acttgcgtag	aatttcaacc	2400
cttggcaact	gcacagcttg	gttgtgaaca	gcattcttcaa	ttctggcaag	ctccttgtct	2460
gtcatatcga	cagccaacag	aatcacctgg	gaatcaatac	catgttcagc	ttgagacaga	2520
aggctgagg	caacgaaatc	tggatcagcg	tatttatcag	caataactag	aacttcagaa	2580
ggcccagcag	gcatgtcaat	actacacagg	gctgatgtgt	cattttgaac	catcatcttg	2640
gcagcagtaa	cgaaactggt	tcttggaacca	aataatttgt	cacacttagg	aacagtttct	2700
gttccgtaag	ccatagcagc	tactgcctgg	gcgcctcctg	ctagcacgat	acacttagca	2760
ccaaccttgt	gggcaacgta	gatgacttct	ggggtaaggg	taccatcctt	cttaggtgga	2820
gatgcacaaa	caatttcttt	gcaaccagca	actttggcag	gaacacccag	catcagggaa	2880
gtggaaggca	gaattgcggg	tccaaccagga	atatagaggc	caactttctc	aatagggtct	2940
gcaaaacgag	agcagactac	accagggcaa	gtctcaactt	gcaacgtctc	cgtaggttga	3000
gcttcatgga	atttctctgac	gttatctata	gagagatcaa	tggctctctt	aacgttatct	3060
ggcaattgca	taagttcctc	tgggaaagga	gcttctaaca	caggtgtctt	caaagcgact	3120
ccatcaaaact	tggcagttag	ttctaaaagg	gttttgtcac	cattttgacg	aacattgtcg	3180
acaattggtt	tgactaattc	cataatctgt	tccgttttct	ggataggacg	acgaagggca	3240
tcttcaattt	cttgtgagga	ggccttagaa	acgtcaattt	tgcacaattc	aatacgacct	3300
tcagaaggga	cttcttttag	tttggattct	ttctttagggt	gttcttgggt	gtatcctggc	3360
ttggcatctc	cttctcttct	agtgaacctt	agggaacttca	tatccagggt	tctctccacc	3420
tcgtccaacg	tcacacogta	cttggcacat	ctaactaatg	caaaataaaa	taagtcagca	3480
cattcccagg	cttatctctc	cttggattta	gcttctgcaa	gttcatcagc	ttctctcccta	3540
attttagcgt	tcaacaaaaac	ttcgtcgtca	aataaccggt	tgggtataaga	accttctgga	3600
gcattgctct	tacgatccca	caaggtgggt	tccatggctc	taagaccett	tgattggcca	3660
aaacaggaag	tgcgttccaa	gtgacagaaa	ccaacacctg	tttgttcaac	cacaaatttc	3720
aagcagtcct	catcacaaac	caattcgata	cccagcaact	tttgagttgc	tccagatgta	3780
gcacctttat	accacaaacc	gtgacgacga	gattggtaga	ctccagtttg	tgtccttata	3840
gctcccgga	tagacttttt	ggacgagtag	accaggccca	acgagtaatt	agaagagtca	3900
gccaacaaag	tagtgaattag	accatcgggg	cggtcagtag	tcaaagacgc	caacaaaatt	3960
tcactgacag	ggaacttttt	gacatcttca	gaaagttcgt	attcagtagt	caattgccga	4020
gcatcaataa	tggggattat	accagaagca	acagttggaag	tcacatctac	caactttgcg	4080
gtctcagaaa	aagcataaac	agttctacta	ccgccttagg	tgaacttttt	caaatcgccc	4140
agtggagaag	aaaaaggcac	agcgatacta	gcattagcgg	gcaaggatgc	aactttatca	4200
accaggtgct	tatagataac	cctagcgcct	gggatcatcc	tttggacaac	tctttctgcc	4260
aaatctaggt	ccaaaatcac	ttcattgata	caattattgt	acaacttgag	caagttgtcg	4320
atcagctcct	caaattgggtc	ctctgtaacg	gatgactcaa	cttgacacatt	aacttgaagc	4380
tcagtcgatt	gagtgaaact	gatcaggttg	tgacagctgg	cagcagcata	gggaaacacg	4440
gcttttcccta	ccaaactcaa	ggaattatca	aactctgcaa	cacttgcgta	tgcaggtagc	4500
aagggaatag	tcataacttga	agtccgacag	tgagtgtagt	cttgagaaat	tctgaagccg	4560
tatttttatt	atcagtgagt	cagtcacatc	gagatcctct	acgccggacg	catcggtggc	4620
gacctcgagg	gggggggggg	gcgctgaggt	ctgctcggtg	aagaaggtgt	tgctgactca	4680
taccaggcct	gaatcgcccc	atcatccagc	cagaaagtga	gggagccacg	ggtgatgaga	4740
gctttgttgt	aggtggacca	gttgggtgatt	ttgaactttt	gctttgccac	ggaacgggtct	4800
gcgttgtcgg	gaagatgcgt	gatctgatcc	ttcaactcag	caaaagttcg	atttattcaa	4860
caaagccgcc	gtcccgtcaa	gtcagcgtaa	tgctctgcca	gtgttacaac	caattaacca	4920
attctgatta	gaaaaactca	tcgagcatca	aatgaaaactg	caattttattc	atatcaggat	4980
tatcaatacc	atatttttga	aaaagccgtt	tctgtaatga	aggagaaaac	tcaccgaggc	5040
agttccatag	gatggcaaga	tcttggtatc	ggtctgcgat	tccgactcgt	ccaacatcaa	5100
tacaacctat	taatttcccc	tcgtcaaaaa	taaggttatc	aagtgagaaa	tcaccatgag	5160
tgacgactga	atccggtgag	aatggcaaaa	gcttatgcac	ttctttccag	acttggtcaa	5220
caggccagcc	attacgctcg	tcataaaaat	cactcgcatc	aaccaaaccg	ttattcattc	5280
gtgattgcgc	ctgagcgaga	cgaataacgc	gatcgctgtt	aaaaggacaa	ttacaaacag	5340
gaatcgaatg	caaccggcgc	aggaacactg	ccagcgcac	aacaatattt	tcacctgaat	5400
caggatattc	ttctaatacc	tggaaatgctg	ttttcccggg	gatcgagtg	gtgagtaacc	5460
atgcattcat	aggagtacgg	ataaaatgct	tgatggtcgg	aagaggcata	aattccgtca	5520
gccagtttag	tctgaccatc	tcattctgtaa	catcattggc	aacgctacct	ttgccatgtt	5580
tcagaacaac	ctctggcgca	tcgggcttcc	catacaactc	atagattgtc	gcacctgatt	5640
gcccagacatt	atcgcgagcc	catttatacc	catataaatc	agcatccatg	ttggaattta	5700
atcgcgccct	cgagcaagac	gtttcccgtt	gaatatggct	cataacacct	cttgatttac	5760
tgtttatgta	agcagacagt	tttattgttc	atgatgatat	atttttatct	tgtgcaatgt	5820
aacatcagag	attttgagac	acaacgtggc	tttccccccc	ccccctgcag	gtcggcatca	5880
ccggcgccac	aggtgcgggt	gctggcgcc	atatcgccga	catcaccgat	ggggaagatc	5940
gggctcgcca	cttcgggctc	atgagcgctt	gtttcggcgt	gggtatggtg	gcaggcccg	6000
tggccggggg	actggtgggc	gccatctcct	tgcattgcacc	attccttgcg	gcggcggtgc	6060

tcaacggcct	caacctaata	ctgggctgct	tcctaatagca	ggagtcgcat	aagggagagc	6120
gtcagatc	tatgattgga	agtatgggaa	tggtgatacc	cgcattcttc	agtgtcttga	6180
ggctctctat	cagattatgc	ccaactaaag	caaccggagg	aggagatttc	atggtaaatt	6240
tctctgactt	ttgggtcatca	gtagactcga	actgtgagac	tatctcgggt	atgacagcag	6300
aaatgtcctt	cttggagaca	gtaaatgaag	ccccaccaat	aaagaaatcc	ttgttatcag	6360
gaacaaactt	cttgtttcga	actttttcgg	tgccctgaac	tataaaatgt	agagtggata	6420
tgctcggtag	gaatggagcg	ggcaaatgct	taccttctgg	accttcaaga	ggtagttagg	6480
gtttgttagat	actgtgcca	acttcagtga	caacgtttgct	atttcgttca	aaccattccg	6540
aatccagaga	aatcaaagtt	gtttgtctac	tattgatcca	agccagtgcg	gtcttgaaac	6600
tgacaatagt	gtgctcgtgt	tttgaggtca	tctttgtatg	aataaatcta	gtctttgatc	6660
taaaataatct	tgacgagcca	aggcgataaa	tacccaaactc	taaaactctt	ttaaaacggt	6720
aaaaggacaa	gtatgtctgc	ctgtattaaa	ccccaaatca	gctcgtagtc	tgatcctcat	6780
caacttgagg	ggcactatct	tgtttttagag	aaatttgccg	agatgcgata	tcgagaaaaa	6840
ggtagcgtga	ttttaaacgt	gaaatttatac	tcaagatctc	tgccctcgcg	gttctcgggtga	6900
tgacgggtgaa	aacctctgac	acatgcagct	cccgagagacg	gtcacagctt	gtctgtaagc	6960
ggatgcccggg	agcagacaag	ccggtcaggg	cgcgtcagcg	gggtgtggcg	gggtgcgggg	7020
cgcagccatg	acccagtcac	gtagcgatag	cggagtgtat	actggcttaa	ctatgcggca	7080
tcagagcaga	ttgtactgag	agtgcacat	atgcggtgtg	aaataccgca	cagatgcgta	7140
aggagaaaat	accgcacag	gctgtcttcc	gcttctctgc	tcactgactc	gctgcgctcg	7200
gtcgttcggc	tgccggcagc	ggtagcagct	cactcaaagg	cggtaatacag	gttatccaca	7260
gaatcagggg	ataacgcagg	aaagaacatg	tgagcaaaag	gccagcaaaa	ggccaggaac	7320
cgtaaaaagg	ccgcgttgct	ggcgtttttc	cataggtctc	gccccctga	cgagcatcac	7380
aaaaatcgac	gctcaagtc	gaggtggcga	aaccggacag	gactataaag	ataccaggcg	7440
tttccccctg	gaagctccct	cgtgcgctct	cctgttccga	ccctgccgct	taccggatac	7500
ctgtccgcct	ttctcccttc	gggaagcgtg	gcgcttctc	aatgtctcag	ctgtagggtat	7560
ctcagttcgg	tgtaggtcgt	tcgctccaag	ctgggctgtg	tgacgaacc	ccccgttcag	7620
cccagccgct	gcgccttctc	cggttaactat	cgtcttgagt	ccaaccgggt	aagacacgac	7680
ttatcgccac	tgccagcagc	cactgttaac	aggattagca	gagcagggtg	tgtaggcgggt	7740
gtcacagagt	tcttgaagtg	gtggcctaac	tacggctaca	ctagaaggag	agtatttggt	7800
atctgcgtc	tggtgaagcc	agttaccttc	ggaaaaagga	ttggtagctc	ttgatccggc	7860
aaacaaacca	ccgtcgttag	cgggtggttt	tttgtttgca	agcagcagat	tacgcgcaga	7920
aaaaaaggat	ctcaagaaga	tcctttgatc	ttttctacgg	ggctcgacgc	tcagtggaaac	7980
gaaaaactcac	gttaagggat	tttgggtcatg	agattatcaa	aaaggatctt	cacctagatc	8040
cttttaaat	aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	aacttgggtct	8100
gacagttacc	aatgcttaat	cagtgaaggca	cctatctcag	cgatctgtct	atttcgttca	8160
tccatagttg	ccgtactccc	cgtcgtgtag	ataactacga	tacgggaggg	cttaccatct	8220
ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	cggctccaga	tttatcagca	8280
ataaaccagc	cagccggaag	ggccgagcgc	agaagtggtc	ctgcaacttt	atccgcctcc	8340
atccagctca	ttaattgttg	ccgggaagct	agagtaagta	gttcgccagt	taatagtttg	8400
cgcaacgttg	ttgccattgc	tgccagcatc	gtggtgtcac	gctcgtcgtt	tggtatggct	8460
tcattcagct	ccggttccca	acgatcaagg	cgagttacat	gatcccccat	gttgtgcaaa	8520
aaagcggtta	gctccttcgg	tcctccgatac	gttgtcagaa	gtaagtggc	cgcagtgtta	8580
tcactcatgg	ttatggcagc	actgcataat	tctcttactg	tcatgccatc	cgtaagatgc	8640
ttttctgtga	ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	8700
agttgctctt	gcccggcgtc	aacacgggat	aataccgcgc	cacatagcag	aactttaaaa	8760
gtgctcatca	ttggaaaaacg	ttcttcgggg	cgaaaaactct	caaggatctt	accgctgttg	8820
agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	cttcagcatc	ttttactttc	8880
accagcgttt	ctgggtgagc	aaaaacagga	aggcaaaatg	ccgcaaaaaa	gggaataagg	8940
gcgacacgga	aatgttgaat	actcatactc	ttcctttttc	aatattattg	aagcattttat	9000
caggggttatt	gtctcatgag	cggatacata	tttgaatgta	tttagaaaaa	taaacaaaata	9060
gggggttccgc	gcacatttcc	ccgaaaaagtg	ccacctgacg	tctaagaaac	cattattatc	9120
atgacattaa	ctataaaaaa	taggcgtatc	acagggcctt	ttcgtcttca	agaatttaatt	9180
ctcatgtttg	acagcttatac	atcgataagc	tgactcatgt	tggtattgtg	aaatagacgc	9240
agatcgggaa	cactgaaaaa	taacagttat	tattcgc			9276

<210> 46

<211> 3908

<212> DNA

<213> Escherichia coli

<400> 46

agcgcccaat acgcaaacgc cctctccccg cgcgttgggc gattcattaa tgcagctggc

60

acgacaggtt	tcccgaactg	aaagcgggca	gtgagcgcaa	cgcaattaat	gtgagttagc	120
tcactcatta	ggcaccaccg	gctttaact	ttatgcttcc	ggctcgtatg	ttgtgtggaa	180
ttgtgagcgg	ataacaattt	cacacaggaa	acagctatga	ccatgattac	gccaagcttg	240
gtaccgagct	cggatccact	agtaacggcc	gccagtggtg	tggaaatcgc	ccttaagggc	300
gaattctgca	gatatccatc	acactggcgg	ccgctcgagc	atgcatctag	agggcccaat	360
tcgcccctata	gtgagtcgta	ttacaattca	ctggccgctg	ttttacaacg	tcgtgactgg	420
gaaaaccctg	gcgttaccca	acttaatcgc	cttgacgac	atcccccttt	cgccagctgg	480
cgtaatagcg	aagaggcccg	caccgatcgc	ccttcccaac	agttgcgcag	cctgaatggc	540
gaatgggacg	cgccctgtag	cgccgcatla	agcgcggcgg	gtgtgggtgg	tacgcgcagc	600
gtgaccgcta	cacttgccag	cgccctagcg	cccgtccttt	tcgctttctt	cccttccctt	660
ctcgccacgt	tcgcccggct	tcccgtcaa	gctctaaatc	gggggctccc	tttaggggtt	720
cgatttagag	ctttacggca	cctcgaccgc	aaaaaacttg	atttgggtga	tgggttcacgt	780
agtgggcat	cgccctgata	gacgggtttt	cgccctttga	cgttggagtc	cacgttcttt	840
aatagtgagc	tcttggtcca	aactggaaaca	acactcaacc	ctatcgcggt	ctattctttt	900
gatttataag	ggattttgcc	gatttcggcc	tattggttaa	aaaaatgagct	gatttaacaa	960
attcagggcg	caagggctgc	taaaaggaaac	ggaacacgta	gaaagccagt	ccgcagaaac	1020
gggtgctgacc	ccggatgaat	gtcagctact	gggtatctg	gacaaggga	aacgcaagcg	1080
caaagagaaa	gcaggtagct	tgcagtgggc	ttacatggcg	atagctagac	tggcggttt	1140
tatggacagc	aagcgaaccg	gaattgccag	ctggggcgcc	ctctggtaag	gttgggaagc	1200
cctgcaaat	aaactggatg	gctttcttgc	cgccaaagat	ctgatggcg	aggggatcaa	1260
gatctgatca	agagacagga	tgaggatcgt	ttcgcatgat	tgaacaagat	ggattgcacg	1320
caggttctcc	ggccgcttgg	gtggagaggc	tattcggcta	tgactgggca	caacagacaa	1380
tcggctgctc	tgatgcgcgc	gtgttcgcgc	tgtagcgca	ggggcgcccg	gttctttttg	1440
tcaagaccga	cctgtccggg	ggcctgaatg	aactgcagga	cgaggcagcg	cggtctatcgt	1500
ggctggccac	gacggcggtt	ccttgcgcag	ctgtgctcga	cgttgtcact	gaagcgggaa	1560
gggactggct	gctattgggc	gaagtgcggg	ggcaggatct	cctgtcatct	cgcttgcctc	1620
ctgcccagaa	agatccatc	atggctgatg	caatgcggcg	gctgcatacg	cttgatccgg	1680
ctacctgccc	attcgaccac	caagcgaaac	atcgcatcga	gcgagcacgt	actcggatgg	1740
aagccggctc	tgtcgatcag	gatgatctgg	acgaagagca	tcaggggctc	gcgccagccg	1800
aactgttcgg	caggctcaag	gcgcgcagtc	ccgacggcga	ggatctcgtc	gtgatccatg	1860
gcgatgcctg	cttgccgaat	atcatggtgg	aaaatggccg	cttttctgga	ttcaacgact	1920
gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttgatatac	cgtgatattg	1980
ctgaagagct	tggcgcgcaa	tgggtcgacc	gcttccctcg	gctttacggg	atcgccgctc	2040
ccgattcgca	gcgcacgccc	ttctatcgcc	ttcttgacga	gttcttctga	attgaaaaag	2100
gaagagtatg	agtattcaac	atctccgtgt	cgcccttatt	cccttttttg	cggcattttg	2160
ccttctctgt	tttgctcacc	cagaaacgct	ggtgaaagta	aaagatgctg	aagatcagtt	2220
gggtgcacga	gtgggttaca	tcgaactgga	tctcaaacgc	ggtaagatcc	ttgagagttt	2280
tcgcccggaa	gaacgttttc	caatgatgag	cacttttaaa	gttctgctat	gtcatacact	2340
attatcccg	attgacgcgc	ggcaagagca	actcggctcg	cgggcgcggt	attctcagaa	2400
tgacttggtt	gagtactcac	cagtcacaga	aaagcatctt	acggatggca	tgacagtaag	2460
agaattatgc	agtgtgcgca	taaccatgag	tgataaacact	gcggccaact	tacttctgac	2520
aacgatcgga	ggaccgaagg	agctaaccgc	ttttttgcac	aacatggggg	atcatgtaac	2580
tcgcttgat	cgttgggaac	cggagctgaa	tgaagccata	ccaaacgcag	agagtgcac	2640
cacgatgcct	gtagcaatgc	caacaacggt	gcgcaaaact	ttactggcg	aactacttac	2700
tctagcttcc	ggcaacaat	taatagactg	gatggaggcg	gataaagtgg	caggaccact	2760
tctgctctcg	ggccttccgg	ctggctgggt	tattgctgat	aaatctggag	ccggtgagcg	2820
tgggtctcgc	ggtatcattg	cagcactggg	gccagatggg	aagccctccc	gtatcgtagt	2880
tatctacacg	acggggagtc	aggcaactat	ggatgaacga	aatagacaga	tcgctgagat	2940
aggtgcctca	ctgattaaag	attggtaact	gtcagaccaa	gtttactcat	atatacttta	3000
gattgattta	aaacttcatt	tttaatttaa	aaggatctag	gtgaagatcc	tttttgataa	3060
tctcatgacc	aaaatccctt	aacgtgagtt	ttcgttccac	tgagcgtcag	accccgtaga	3120
aaagatcaaa	ggatcttctt	gagatccttt	ttttctgcgc	gtaatctgct	gcttgcaaac	3180
aaaaaaacca	ccgctaccag	cggtggtttg	tttgccggat	caagagctac	caactctttt	3240
tccgaaggta	actggcttca	gcagagcgca	tacaccaaat	actgtccttc	tagtgtagcc	3300
gtagttaggc	caccacttca	agaactctgt	agcaccgct	acatacctcg	ctctgcta	3360
cctgttacca	gtggctgctg	ccagtggcga	taagtctgtg	cttaccgggt	tggactcaag	3420
acgatagtta	ccggataaag	cgcagcggtc	gggctgaacg	gggggttcgt	gcacacagcc	3480
cagcttgag	cgaacgacct	acaccgaact	gagataccta	cagcgtgagc	attgagaaag	3540
cgccacgctt	cccgaaggga	gaaaggcgga	caggtatccg	gtaagcggca	gggtcggaac	3600
aggagagcgc	acgagggagc	ttccaggggg	aaacgcctgg	tatctttata	gtcctgtcgg	3660
gtttcgccac	ctctgacttg	agcgtcgatt	tttctgatgc	tcgctcagggg	ggcggagcct	3720
atggaaaaac	gccagcaacg	cgcccttttt	acggttctctg	gccttttgct	ggccttttgc	3780

```

tcacatgttc tttcctgcgt tatccctga ttctgtggat aaccgtatta cgcctttga 3840
gtgagctgat accgctcgcc gcagccgaac gaccgagcgc agcgagtcag tgagcgagga 3900
agcggaag 3908

```

```

<210> 47
<211> 46
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Oligonucleotide primer

```

```

<400> 47
ggaattccat atgccgcgct ttaaagtggg ggggtggggag gagggc 46

```

```

<210> 48
<211> 32
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Oligonucleotide primer

```

```

<400> 48
cgcgataccc gttacagctc agccttccag ac 32

```

```

<210> 49
<211> 3147
<212> DNA
<213> Homo Sapien

```

```

<220>
<221> CDS
<222> (1865)...(2590)
<223> Nucleic acid sequence of protease domain of MTSP1

```

```

<400> 49
tcaagagcgg cctcggggta ccatggggag cgatcggggc cgcaagggcg gagggggccc 60
gaaggacttc ggcgcgggac tcaagtacaa ctcccggcac gagaaagtga atggccttga 120
ggaaggcgtg gaggctcctgc cagtcaacaa cgtcaagaag gtggaaaagc atggcccggg 180
gcgctgggtg gtgctggcag ccgtgctgat cggcctcctc ttggtcctgc tgggggatcg 240
cttcctgggtg tggcatttgc agtaccggga cgtgcgtgtc cagaaggctc tcaatggcta 300
catgaggatc acaaatgaga attttgtgga tgcctacgag aactccaaact cactgagtt 360
tgtaagcctg gccagcaagg tgaaggacgc gctgaagctg ctgtacagcg gagtccatt 420
cctgggcccc taccacaagg agtcggctgt gacggccttc agcgagggca gcgtcatcgc 480
ctactactgg tctgagttca gcatcccgcg gcacctggtg gaggaggcgc agcgcgtcat 540
ggccgaggag cgcgtagtcg tgctgccccg gcgggcgcgc tccttgaagt cctttgtggt 600
cacctcagtg gtggctttcc ccacggactc caaaacagta cagaggaccg aggacaacag 660
ctgcagcttt ggcctgcacg cccgcggtgt ggagctgatg cgcttcacca cgcctggctt 720
ccctgacagc ccctaccccg ctcatgcccg ctgccagtgg gccctgcggg ggagcgccga 780
ctcagtgtcg agcctcacct tccgcagctt tgaccttgcg tcctgcgacg agcgcggcag 840
cgacctggtg acggtgtaca acaccctgag ccccatggag cccacgcgcc tgggtgcagtt 900
gtgtggcacc taccctccct cctacaacct gaccttcac tcctcccaga acgtcctgct 960
catcacactg ataaccaaca ctgagcggcg gcatcccggc tttgaggcca ccttcttcca 1020
gctgcctagg atgagcagct gtggaggccg cttacgtaaa gccagggga cattcaacag 1080
cccctactac ccaggccact accacccaa cattgactgc acatggaaca ttgaggtgcc 1140
caacaaccag catgtgaagg tgagcttcaa attcttctac ctgctggagc cggcggtgcc 1200
tgccggcacc tgcccccaagg actacgtgga gatcaatggg gagaataact gcggagagag 1260
gtcccagttc gtcgtcacca gcaacagcaa caagatcaca gttcgcttcc actcagatca 1320
gtcctacacc gacaccgggt tcttagctga atacctctc tacgactcca gtgacccatg 1380
cccgggcag ttcacgtgcc gcacggggcg gtgtatccgg aaggagctgc gctgtgatgg 1440
ctgggcccag tgcaccgacc acagcgatga gctcaactgc agttgcgacg ccggccacca 1500

```

gttcacgtgc aagaacaagt tctgcaagcc cctcttcttg gtctgcgaca gtgtgaacga	1560
ctgcggagac aacagcgacg agcaggggtg cagttgtccg gcccagacct tcaggtgttc	1620
caatgggaag tgctctctga aaagccagca gtgcaatggg aaggacgact gtggggacgg	1680
gtccgacgag gcctcctgcc ccaaggtgaa cgtcgtcact tgtaccaaac acacctaccg	1740
ctgcctcaat gggctctgct tgagcaaggg caaccctgag tgtgacggga aggaggactg	1800
tagcgacggc tcagatgaga aggactgcga ctgtgggctg cggtcattca cgagacaggc	1860
tcgt gtt gtt ggg ggc acg gat gcg gat gag ggc gag tgg ccc tgg cag	1909
Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln	
1 5 10 15	
gta agc ctg cat gct ctg ggc cag ggc cac atc tgc ggt gct tcc ctc	1957
Val Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu	
20 25 30	
atc tct ccc aac tgg ctg gtc tct gcc gca cac tgc tac atc gat gac	2005
Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp	
35 40 45	
aga gga ttc agg tac tca gac ccc acg cag tgg acg gcc ttc ctg ggc	2053
Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly	
50 55 60	
ttg cac gac cag agc cag cgc agc gcc cct ggg gtg cag gag cgc agg	2101
Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg	
65 70 75	
ctc aag cgc atc atc tcc cac ccc ttc ttc aat gac ttc acc ttc gac	2149
Leu Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp	
80 85 90 95	
tat gac atc gcg ctg ctg gag ctg gag aaa ccg gca gag tac agc tcc	2197
Tyr Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser	
100 105 110	
atg gtg cgg ccc atc tgc ctg ccg gac gcc tcc cat gtc ttc cct gcc	2245
Met Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala	
115 120 125	
ggc aag gcc atc tgg gtc acg ggc tgg gga cac acc cag tat gga ggc	2293
Gly Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly	
130 135 140	
act ggc gcg ctg atc ctg caa aag ggt gag atc cgc gtc atc aac cag	2341
Thr Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln	
145 150 155	
acc acc tgc gag aac ctc ctg ccg cag cag atc acg ccg cgc atg atg	2389
Thr Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met	
160 165 170 175	
tgc gtg ggc ttc ctc agc ggc ggc gtg gac tcc tgc cag ggt gat tcc	2437
Cys Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser	
180 185 190	
ggg gga ccc ctg tcc agc gtg gag gcg gat ggg cgg atc ttc cag gcc	2485
Gly Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala	
195 200 205	
ggt gtg gtg agc tgg gga gac ggc tgc gct cag agg aac aag cca ggc	2533
Gly Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly	
210 215 220	

gtg tac aca agg ctc cct ctg ttt cgg gac tgg atc aaa gag aac act 2581
 Val Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr
 225 230 235

ggg gta tag gggccggggc caccctaaatg tgtacacctg cggggccacc 2630
 Gly Val *
 240

catcgctccac cccagtgtgc acgcctgcag gctggagact ggaccgctga ctgcaccagc 2690
 gccccagaa catacactgt gaactcaatc tccagggctc caaatctgcc tagaaaacct 2750
 ctgccttcct cagcctccaa agtggagctg ggaggtagaa ggggaggaca ctggtggttc 2810
 tactgaccca actgggggca aaggtttgaa gacacagcct cccccgccag cccaagctg 2870
 ggccgaggcg cgtttgtgta tatctgcctc cctgtctgt aaggagcagc gggaacggag 2930
 ctccggagcc tcctcagtga aggtggtggg gctgccggat ctgggctgtg gggcccttgg 2990
 gccacgctct tgaggaagcc caggctcgga ggaccctgga aaacagacgg gtctgagact 3050
 gaaattgttt taccagctcc cagggtggac ttcagtgtgt gtatttgtgt aaatgggtaa 3110
 aacaatttat ttctttttta aaaaaaaaaa aaaaaaa 3147

<210> 50
 <211> 241
 <212> PRT
 <213> Homo Sapien

<400> 50
 Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val
 1 5 10 15
 Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu Ile
 20 25 30
 Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp Asp Arg
 35 40 45
 Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe Leu Gly Leu
 50 55 60
 His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln Glu Arg Arg Leu
 65 70 75
 Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp Phe Thr Phe Asp Tyr
 85 90 95
 Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro Ala Glu Tyr Ser Ser Met
 100 105 110
 Val Arg Pro Ile Cys Leu Pro Asp Ala Ser His Val Phe Pro Ala Gly
 115 120 125
 Lys Ala Ile Trp Val Thr Gly Trp Gly His Thr Gln Tyr Gly Gly Thr
 130 135 140
 Gly Ala Leu Ile Leu Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr
 145 150 155
 Thr Cys Glu Asn Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys
 165 170 175
 Val Gly Phe Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly
 180 185 190
 Gly Pro Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly
 195 200 205
 Val Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val
 210 215 220
 Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr Gly
 225 230 235 240
 Val

<210> 51
 <211> 46
 <212> DNA
 <213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 51
tctctcgaga aaagagtggg ggggtgggtgg ggaggaggcc tctgtg 46

<210> 52
<211> 43
<212> DNA
<213> Artificial sequence

<400> 52
gctcctcatc aaagaagggc agagagatgg gcctgactgt gcc 43

<210> 53
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 53
attcggggcc gcattacagc tcagccttcc agac 34

<210> 54
<211> 43
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 54
ggcacagtca ggcccatctc tctgcccttc tttgatgagg agc 43

<210> 55
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 55
cacccttct tcaatgactt caccttcg 28

<210> 56
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Oligonucleotide Primer

<400> 56
tacctctcct acgactcc 18

<210> 57
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
 <223> Oligonucleotide Primer

 <400> 57
 gaggttctcg caggtggtct ggttg 25

 <210> 58
 <211> 39
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Oligonucleotide Primer

 <400> 58
 ctcgagaaaa gagttggttg gggcacggat gcggatgag 39

 <210> 59
 <211> 11
 <212> PRT
 <213> Homo Sapien

 <400> 59
 Phe Glu Val Phe Ser Gln Ser Ser Ser Leu Gly
 1 5 10

 <210> 60
 <211> 32
 <212> PRT
 <213> Homo Sapien

 <400> 60
 Glu Ile Val Ala Pro Arg Glu Arg Ala Asp Arg Arg Gly Arg Lys Leu
 1 5 10 15
 Leu Cys Trp Arg Lys Pro Thr Lys Met Lys Gly Pro Arg Pro Ser His
 20 25 30

 <210> 61
 <211> 4933
 <212> DNA
 <213> Homo Sapien

 <220>
 <221> CDS
 <222> (94)...(3222)
 <223> Nucleotide sequence encoding corin

 <300>
 <308> GenBank AF133845
 <309> 1999-05-24

 <400> 61
 aaatcatccg tagtgctcc ccgggggaca cgtagaggag agaaaagcga ccaagataaa 60
 agtggacaga agaataagcg agacttttta tcc atg aaa cag tct cct gcc ctc 114
 Met Lys Gln Ser Pro Ala Leu
 1 5

 gct ccg gaa gag cgc tac cgc aga gcc ggg tcc cca aag ccg gtc ttg 162
 Ala Pro Glu Glu Arg Tyr Arg Arg Ala Gly Ser Pro Lys Pro Val Leu
 10 15 20

 aga gct gat gac aat aac atg ggc aat ggc tgc tct cag aag ctg gcg 210

gga atc tgc atc ccc ggg aaa ctg caa tgt aat ggc tac aac gac tgt Gly Ile Cys Ile Pro Gly Lys Leu Gln Cys Asn Gly Tyr Asn Asp Cys 280 285 290 295	978
gac gac tgg agt gac gag gct cat tgc aac tgc agc gag aat ctg ttt Asp Asp Trp Ser Asp Glu Ala His Cys Asn Cys Ser Glu Asn Leu Phe 300 305 310	1026
cac tgt cac aca ggc aag tgc ctt aat tac agc ctt gtg tgt gat gga His Cys His Thr Gly Lys Cys Leu Asn Tyr Ser Leu Val Cys Asp Gly 315 320 325	1074
tat gat gac tgt ggg gat ttg agt gat gag caa aac tgt gat tgc aat Tyr Asp Asp Cys Gly Asp Leu Ser Asp Glu Gln Asn Cys Asp Cys Asn 330 335 340	1122
ccc aca aca gag cat cgc tgc ggg gac ggg cgc tgc atc gcc atg gag Pro Thr Thr Glu His Arg Cys Gly Asp Gly Arg Cys Ile Ala Met Glu 345 350 355	1170
tgg gtg tgt gat ggt gac cac gac tgt gtg gat aag tcc gac gag gtc Trp Val Cys Asp Gly Asp His Asp Cys Val Asp Lys Ser Asp Glu Val 360 365 370 375	1218
aac tgc tcc tgt cac agc cag ggt ctg gtg gaa tgc aga aat gga caa Asn Cys Ser Cys His Ser Gln Gly Leu Val Glu Cys Arg Asn Gly Gln 380 385 390	1266
tgt atc ccc agc acg ttt caa tgt gat ggt gac gag gac tgc aag gat Cys Ile Pro Ser Thr Phe Gln Cys Asp Gly Asp Glu Asp Cys Lys Asp 395 400 405	1314
ggg agt gat gag gag aac tgc agc gtc att cag act tca tgt caa gaa Gly Ser Asp Glu Glu Asn Cys Ser Val Ile Gln Thr Ser Cys Gln Glu 410 415 420	1362
gga gac caa aga tgc ctc tac aat ccc tgc ctt gat tca tgt ggt ggt Gly Asp Gln Arg Cys Leu Tyr Asn Pro Cys Leu Asp Ser Cys Gly Gly 425 430 435	1410
agc tct ctc tgt gac ccg aac aac agt ctg aat aac tgt agt caa tgt Ser Ser Leu Cys Asp Pro Asn Asn Ser Leu Asn Asn Cys Ser Gln Cys 440 445 450 455	1458
gaa cca att aca ttg gaa ctc tgc atg aat ttg ccc tac aac agt aca Glu Pro Ile Thr Leu Glu Leu Cys Met Asn Leu Pro Tyr Asn Ser Thr 460 465 470	1506
agt tat cca aat tat ttt ggc cac agg act caa aag gaa gca tcc atc Ser Tyr Pro Asn Tyr Phe Gly His Arg Thr Gln Lys Glu Ala Ser Ile 475 480 485	1554
agc tgg gag tct tct ctt ttc cct gca ctt gtt caa acc aac tgt tat Ser Trp Glu Ser Ser Leu Phe Pro Ala Leu Val Gln Thr Asn Cys Tyr 490 495 500	1602
aaa tac ctc atg ttc ttt tct tgc acc att ttg gta cca aaa tgt gat Lys Tyr Leu Met Phe Phe Ser Cys Thr Ile Leu Val Pro Lys Cys Asp 505 510 515	1650
gtg aat aca ggc gag cgt atc cct cct tgc agg gca ttg tgt gaa cac	1698

Val 520	Asn	Thr	Gly	Glu	Arg 525	Ile	Pro	Pro	Cys	Arg 530	Ala	Leu	Cys	Glu	His 535	
tct Ser	aaa Lys	gaa Glu	cgc Arg	tgt Cys	gag Glu	tct Ser	gtt Val	ctt Leu	ggg Gly	att Ile	gtg Val	ggc Gly	cta Leu	cag Gln	tgg Trp	1746
cct Pro	gaa Glu	gac Asp	aca Thr	gat Asp	tgc Cys	agt Ser	caa Gln	ttt Phe	cca Pro	gag Glu	gaa Glu	aat Asn	tca Ser	gac Asp	aat Asn	1794
caa Gln	acc Thr	tgc Cys	ctg Leu	atg Met	cct Pro	gat Asp	gaa Gln	tat Tyr	gtg Val	gaa Glu	gaa Glu	tgc Cys	tca Ser	cct Pro	agt Ser	1842
cat His	ttc Phe	aag Lys	tgc Cys	cgc Arg	tca Ser	gga Gly	cag Gln	tgt Cys	gtt Val	ctg Leu	gct Ala	tcc Ser	aga Arg	aga Arg	tgt Cys	1890
gat Asp	ggc Gly	cag Gln	gcc Ala	gac Asp	tgt Cys	gac Asp	gat Asp	gac Asp	agt Ser	gat Asp	gag Glu	gaa Glu	aac Asn	tgt Cys	ggt Gly	1938
tgt Cys	aaa Lys	gag Glu	aga Arg	gat Asp	ctt Leu	tgg Trp	gaa Glu	tgt Cys	cca Pro	tcc Ser	aat Asn	aaa Lys	caa Gln	tgt Cys	ttg Leu	1986
aag Lys	cac His	aca Thr	gtg Val	atc Ile	tgc Cys	gat Asp	ggg Gly	ttc Phe	cca Pro	gac Asp	tgc Cys	cct Pro	gat Asp	tac Tyr	atg Met	2034
gac Asp	gag Glu	aaa Lys	aac Asn	tgc Cys	tca Ser	ttt Phe	tgc Cys	caa Gln	gat Asp	gat Asp	gag Glu	ctg Leu	gaa Glu	tgt Cys	gca Ala	2082
aac Asn	cat His	gcg Ala	tgt Cys	gtg Val	tca Ser	cgt Arg	gac Asp	ctg Leu	tgg Trp	tgt Cys	gat Asp	ggg Gly	gaa Glu	gcc Ala	gac Asp	2130
tgc Cys	tca Ser	gac Asp	agt Ser	tca Ser	gat Asp	gaa Glu	tgg Trp	gac Asp	tgt Cys	gtg Val	acc Thr	ctc Leu	tct Ser	ata Ile	aat Asn	2178
gtg Val	aac Asn	tcc Ser	tct Ser	tcc Ser	ttt Phe	ctg Leu	atg Met	gtt Val	cac His	aga Arg	gct Ala	gcc Ala	aca Thr	gaa Glu	cac His	2226
cat His	gtg Val	tgt Cys	gca Ala	gat Asp	ggc Gly	tgg Trp	cag Gln	gag Glu	ata Ile	ttg Leu	agt Ser	cag Gln	ctg Leu	gcc Ala	tgc Cys	2274
aag Lys	cag Gln	atg Met	ggt Gly	tta Leu	gga Gly	gaa Glu	cca Pro	tct Ser	gtg Val	acc Thr	aaa Lys	ttg Leu	ata Ile	cag Gln	gaa Glu	2322
cag Gln	gag Glu	aaa Lys	gag Glu	ccg Pro	cgg Arg	tgg Trp	ctg Leu	aca Thr	tta Leu	cac His	tcc Ser	aac Asn	tgg Trp	gag Glu	agc Ser	2370
ctc Leu	aat Asn	ggg Gly	acc Thr	act Thr	tta Leu	cat His	gaa Glu	ctt Leu	cta Leu	gta Val	aat Asn	ggg Gly	cag Gln	tct Ser	tgt Cys	2418

gag	agc	aga	agt	aaa	att	tct	ctt	ctg	tgt	act	aaa	caa	gac	tgt	ggg	2466
Glu	Ser	Arg	Ser	Lys	Ile	Ser	Leu	Leu	Cys	Thr	Lys	Gln	Asp	Cys	Gly	
				780					785					790		
cgc	cgc	cct	gct	gcc	cga	atg	aac	aaa	agg	atc	ctt	gga	ggt	cgg	acg	2514
Arg	Arg	Pro	Ala	Ala	Arg	Met	Asn	Lys	Arg	Ile	Leu	Gly	Gly	Arg	Thr	
			795					800					805			
agt	cgc	cct	gga	agg	tgg	cca	tgg	cag	tgt	tct	ctg	cag	agt	gaa	ccc	2562
Ser	Arg	Pro	Gly	Arg	Trp	Pro	Trp	Gln	Cys	Ser	Leu	Gln	Ser	Glu	Pro	
		810					815					820				
agt	gga	cat	atc	tgt	ggc	tgt	gtc	ctc	att	gcc	aag	aag	tgg	gtt	ctg	2610
Ser	Gly	His	Ile	Cys	Gly	Cys	Val	Leu	Ile	Ala	Lys	Lys	Trp	Val	Leu	
	825					830					835					
aca	gtt	gcc	cac	tgc	ttc	gag	ggg	aga	gag	aat	gct	gca	gtt	tgg	aaa	2658
Thr	Val	Ala	His	Cys	Phe	Glu	Gly	Arg	Glu	Asn	Ala	Ala	Val	Trp	Lys	
	840				845					850					855	
gtg	gtg	ctt	ggc	atc	aac	aat	cta	gac	cat	cca	tca	gtg	ttc	atg	cag	2706
Val	Val	Leu	Gly	Ile	Asn	Asn	Leu	Asp	His	Pro	Ser	Val	Phe	Met	Gln	
				860					865					870		
aca	cgc	ttt	gtg	aag	acc	atc	atc	ctg	cat	ccc	cgc	tac	agt	cga	gca	2754
Thr	Arg	Phe	Val	Lys	Thr	Ile	Ile	Leu	His	Pro	Arg	Tyr	Ser	Arg	Ala	
			875					880					885			
gtg	gtg	gac	tat	gac	atc	agc	atc	gtt	gag	ctg	agt	gaa	gac	atc	agt	2802
Val	Val	Asp	Tyr	Asp	Ile	Ser	Ile	Val	Glu	Leu	Ser	Glu	Asp	Ile	Ser	
		890					895					900				
gag	act	ggc	tac	gtc	cgg	cct	gtc	tgc	ttg	ccc	aac	ccg	gag	cag	tgg	2850
Glu	Thr	Gly	Tyr	Val	Arg	Pro	Val	Cys	Leu	Pro	Asn	Pro	Glu	Gln	Trp	
	905					910					915					
cta	gag	cct	gac	acg	tac	tgc	tat	atc	aca	ggc	tgg	ggc	cac	atg	ggc	2898
Leu	Glu	Pro	Asp	Thr	Tyr	Cys	Tyr	Ile	Thr	Gly	Trp	Gly	His	Met	Gly	
	920				925					930					935	
aat	aaa	atg	cca	ttt	aag	ctg	caa	gag	gga	gag	gtc	cgc	att	att	tct	2946
Asn	Lys	Met	Pro	Phe	Lys	Leu	Gln	Glu	Gly	Glu	Val	Arg	Ile	Ile	Ser	
			940					945						950		
ctg	gaa	cat	tgt	cag	tcc	tac	ttt	gac	atg	aag	acc	atc	acc	act	cgg	2994
Leu	Glu	His	Cys	Gln	Ser	Tyr	Phe	Asp	Met	Lys	Thr	Ile	Thr	Thr	Arg	
			955					960					965			
atg	ata	tgt	gct	ggc	tat	gag	tct	ggc	aca	gtt	gat	tca	tgc	atg	ggg	3042
Met	Ile	Cys	Ala	Gly	Tyr	Glu	Ser	Gly	Thr	Val	Asp	Ser	Cys	Met	Gly	
		970					975					980				
gac	agc	ggg	ggg	cct	ctt	gtt	tgt	gag	aag	cct	gga	gga	cgg	tgg	aca	3090
Asp	Ser	Gly	Gly	Pro	Leu	Val	Cys	Glu	Lys	Pro	Gly	Gly	Arg	Trp	Thr	
		985				990					995					
tta	ttt	gga	tta	act	tca	tgg	ggc	tcc	gtc	tgc	ttt	tcc	aaa	gtc	ctg	3138
Leu	Phe	Gly	Leu	Thr	Ser	Trp	Gly	Ser	Val	Cys	Phe	Ser	Lys	Val	Leu	
	1000				1005					1010					1015	
ggg	cct	ggc	gtt	tat	agt	aat	gtg	tca	tat	ttc	gtc	gaa	tgg	att	aaa	3186

Gly Pro Gly Val Tyr Ser Asn Val Ser Tyr Phe Val Glu Trp Ile Lys
 1020 1025 1030

aga cag att tac atc cag acc ttt ctc cta aac taa ttataaggat 3232
 Arg Gln Ile Tyr Ile Gln Thr Phe Leu Leu Asn *
 1035 1040

gatcagagac ttttgccagc taaactaaaa gaaaatggcc ttcttgactg tgaagagctg 3292
 cctgcagaga gctgtacaga agcacttttc atggacagaa atgctcaatc gtgcactgca 3352
 aatttgcattg tttgttttgg actaattttt ttcaatttat tttttcacct tcatttttct 3412
 cttattttcaa gttcaatgaa agacttttaca aaagcaaaca aagcagactt tgtccttttg 3472
 ccaggcctaa ccatgactgc agcacaaaaat tatcgactct ggcgagattt aaaatcaggt 3532
 gctacagtaa ccagttatgg aatgggtctct ttatcctat cacaaaaaaa gacatagata 3592
 tttaggctga ttaattatct ctaccagttt ttgtttctca agctcagtgac atagtggtaa 3652
 atttcagtggt taacattgga gacttgcttt tctttttctt tttttatacc ccacaattct 3712
 tttttattac acttcgaatt ttaggggtaca cgagcacaaac gtgcagggtta gttacatatg 3772
 tatacatgtg ccatgttggt gtgctgaacc cagtaactcg tcatttgatt tattaaaagc 3832
 caagataatt tacatgttta aagtattttac tattaccccc ttctaattgtt tgcataattc 3892
 tgagaactga taaaagacag caataaaaaga ccagtgtcat ccatttaggt agcaagacat 3952
 attgaatgca aagtctcttta gatatacaata ttaacacttg acattatttg acccccatt 4012
 ctggatgtat atcaagatca taattttata gaagagtctc tatagaactg tcctcatagc 4072
 tgggttttgtt caggatataat gagttggctg attgagactg caacaactac atctatatatt 4132
 atgggcaata ttttgtttta cttatgtggc aaagaactgg atattaaact ttgcaaaaaga 4192
 gaatttagat gagagatgca attttttaaa aagaaaatta atttgcattc ctcgtttaat 4252
 taaattttatt tttcagtttt cttgctgtca tccataccaa caaagtcata aagagcatat 4312
 ttttagagcac agtaagactt tgcattggagt aaaacatttt gtaattttcc tcaaaaagatg 4372
 ttttaatatct ggtttcttct cattggtaat taaaatttta gaaatgattt ttagctctag 4432
 gccactttac gcaactcaat ttctgaagca attagtggta aaaagtattt ttccccacta 4492
 aaaaaacttta aaacacaaat cttcatatat acttaattta attagtcagg catccatttt 4552
 gcctttttaa caactaggat tccctactaa cctccaccag caacctggac tgcctcagca 4612
 ttccaaatag atactacctg caattttata catgtatttt tgtatctttt ctgtgtgtaa 4672
 acatagttag aattcaaaaa gttgtagcaa ttctatact attcatctcc tgcctctcag 4732
 tttgtataaa cctaaggaga gtgtgaaatc cagcaactga attgtgggtc cgattgtatg 4792
 aaagtccaag aacatatgtc agttttgtta cagttgtagc tacatactca atgtatcaac 4852
 ttttagcctg ctcaacttga gctcagtgaa atatatatat tatacttatt ttaaataatt 4912
 ctttaatacaa ataaaaatgg a

<210> 62
 <211> 1042
 <212> PRT
 <213> Homo Sapien

<400> 62
 Met Lys Gln Ser Pro Ala Leu Ala Pro Glu Glu Arg Tyr Arg Arg Ala
 1 5 10 15
 Gly Ser Pro Lys Pro Val Leu Arg Ala Asp Asp Asn Asn Met Gly Asn
 20 25 30
 Gly Cys Ser Gln Lys Leu Ala Thr Ala Asn Leu Leu Arg Phe Leu Leu
 35 40 45
 Leu Val Leu Ile Pro Cys Ile Cys Ala Leu Val Leu Leu Val Ile
 50 55 60
 Leu Leu Ser Tyr Val Gly Thr Leu Gln Lys Val Tyr Phe Lys Ser Asn
 65 70 75 80
 Gly Ser Glu Pro Leu Val Thr Asp Gly Glu Ile Gln Gly Ser Asp Val
 85 90 95
 Ile Leu Thr Asn Thr Ile Tyr Asn Gln Ser Thr Val Val Ser Thr Ala
 100 105 110
 His Pro Asp Gln His Val Pro Ala Trp Thr Thr Asp Ala Ser Leu Pro
 115 120 125
 Gly Asp Gln Ser His Arg Asn Thr Ser Ala Cys Met Asn Ile Thr His
 130 135 140
 Ser Gln Cys Gln Met Leu Pro Tyr His Ala Thr Leu Thr Pro Leu Leu

145					150					155					160
Ser	Val	Val	Arg	Asn	Met	Glu	Met	Glu	Lys	Phe	Leu	Lys	Phe	Phe	Thr
				165					170					175	
Tyr	Leu	His	Arg	Leu	Ser	Cys	Tyr	Gln	His	Ile	Met	Leu	Phe	Gly	Cys
			180					185					190		
Thr	Leu	Ala	Phe	Pro	Glu	Cys	Ile	Ile	Asp	Gly	Asp	Asp	Ser	His	Gly
		195					200					205			
Leu	Leu	Pro	Cys	Arg	Ser	Phe	Cys	Glu	Ala	Ala	Lys	Glu	Gly	Cys	Glu
	210					215					220				
Ser	Val	Leu	Gly	Met	Val	Asn	Tyr	Ser	Trp	Pro	Asp	Phe	Leu	Arg	Cys
225					230					235				240	
Ser	Gln	Phe	Arg	Asn	Gln	Thr	Glu	Ser	Ser	Asn	Val	Ser	Arg	Ile	Cys
				245					250					255	
Phe	Ser	Pro	Gln	Gln	Glu	Asn	Gly	Lys	Gln	Leu	Leu	Cys	Gly	Arg	Gly
			260				265						270		
Glu	Asn	Phe	Leu	Cys	Ala	Ser	Gly	Ile	Cys	Ile	Pro	Gly	Lys	Leu	Gln
	275						280					285			
Cys	Asn	Gly	Tyr	Asn	Asp	Cys	Asp	Asp	Trp	Ser	Asp	Glu	Ala	His	Cys
	290					295					300				
Asn	Cys	Ser	Glu	Asn	Leu	Phe	His	Cys	His	Thr	Gly	Lys	Cys	Leu	Asn
305					310					315				320	
Tyr	Ser	Leu	Val	Cys	Asp	Gly	Tyr	Asp	Asp	Cys	Gly	Asp	Leu	Ser	Asp
			325						330					335	
Glu	Gln	Asn	Cys	Asp	Cys	Asn	Pro	Thr	Thr	Glu	His	Arg	Cys	Gly	Asp
		340						345					350		
Gly	Arg	Cys	Ile	Ala	Met	Glu	Trp	Val	Cys	Asp	Gly	Asp	His	Asp	Cys
	355						360					365			
Val	Asp	Lys	Ser	Asp	Glu	Val	Asn	Cys	Ser	Cys	His	Ser	Gln	Gly	Leu
	370					375					380				
Val	Glu	Cys	Arg	Asn	Gly	Gln	Cys	Ile	Pro	Ser	Thr	Phe	Gln	Cys	Asp
385					390					395				400	
Gly	Asp	Glu	Asp	Cys	Lys	Asp	Gly	Ser	Asp	Glu	Glu	Asn	Cys	Ser	Val
				405					410				415		
Ile	Gln	Thr	Ser	Cys	Gln	Glu	Gly	Asp	Gln	Arg	Cys	Leu	Tyr	Asn	Pro
	420							425					430		
Cys	Leu	Asp	Ser	Cys	Gly	Gly	Ser	Ser	Leu	Cys	Asp	Pro	Asn	Asn	Ser
	435						440					445			
Leu	Asn	Asn	Cys	Ser	Gln	Cys	Glu	Pro	Ile	Thr	Leu	Glu	Leu	Cys	Met
	450					455					460				
Asn	Leu	Pro	Tyr	Asn	Ser	Thr	Ser	Tyr	Pro	Asn	Tyr	Phe	Gly	His	Arg
465					470				475					480	
Thr	Gln	Lys	Glu	Ala	Ser	Ile	Ser	Trp	Glu	Ser	Ser	Leu	Phe	Pro	Ala
				485					490					495	
Leu	Val	Gln	Thr	Asn	Cys	Tyr	Lys	Tyr	Leu	Met	Phe	Phe	Ser	Cys	Thr
	500						505						510		
Ile	Leu	Val	Pro	Lys	Cys	Asp	Val	Asn	Thr	Gly	Glu	Arg	Ile	Pro	Pro
	515						520					525			
Cys	Arg	Ala	Leu	Cys	Glu	His	Ser	Lys	Glu	Arg	Cys	Glu	Ser	Val	Leu
	530					535					540				
Gly	Ile	Val	Gly	Leu	Gln	Trp	Pro	Glu	Asp	Thr	Asp	Cys	Ser	Gln	Phe
545					550					555				560	
Pro	Glu	Glu	Asn	Ser	Asp	Asn	Gln	Thr	Cys	Leu	Met	Pro	Asp	Glu	Tyr
			565						570					575	
Val	Glu	Glu	Cys	Ser	Pro	Ser	His	Phe	Lys	Cys	Arg	Ser	Gly	Gln	Cys
	580						585						590		
Val	Leu	Ala	Ser	Arg	Arg	Cys	Asp	Gly	Gln	Ala	Asp	Cys	Asp	Asp	Asp
	595						600					605			
Ser	Asp	Glu	Glu	Asn	Cys	Gly	Cys	Lys	Glu	Arg	Asp	Leu	Trp	Glu	Cys
	610					615					620				
Pro	Ser	Asn	Lys	Gln	Cys	Leu	Lys	His	Thr	Val	Ile	Cys	Asp	Gly	Phe
625					630					635				640	
Pro	Asp	Cys	Pro	Asp	Tyr	Met	Asp	Glu	Lys	Asn	Cys	Ser	Phe	Cys	Gln

Asp	Asp	Glu	Leu	645	Glu	Cys	Ala	Asn	His	650	Ala	Cys	Val	Ser	Arg	655	Asp	Leu
			660						665							670		
Trp	Cys	Asp	Gly	Glu	Ala	Asp	Cys	Ser	Asp	Ser	Ser	Asp	Glu	Trp	Asp			
		675					680						685					
Cys	Val	Thr	Leu	Ser	Ile	Asn	Val	Asn	Ser	Ser	Ser	Phe	Leu	Met	Val			
		690				695						700						
His	Arg	Ala	Ala	Thr	Glu	His	His	Val	Cys	Ala	Asp	Gly	Trp	Gln	Glu			
705					710				715					720				
Ile	Leu	Ser	Gln	Leu	Ala	Cys	Lys	Gln	Met	Gly	Leu	Gly	Glu	Pro	Ser			
			725						730					735				
Val	Thr	Lys	Leu	Ile	Gln	Glu	Gln	Glu	Lys	Glu	Pro	Arg	Trp	Leu	Thr			
			740					745						750				
Leu	His	Ser	Asn	Trp	Glu	Ser	Leu	Asn	Gly	Thr	Thr	Leu	His	Glu	Leu			
		755					760					765						
Leu	Val	Asn	Gly	Gln	Ser	Cys	Glu	Ser	Arg	Ser	Lys	Ile	Ser	Leu	Leu			
		770				775					780							
Cys	Thr	Lys	Gln	Asp	Cys	Gly	Arg	Arg	Pro	Ala	Ala	Arg	Met	Asn	Lys			
785					790				795					800				
Arg	Ile	Leu	Gly	Gly	Arg	Thr	Ser	Arg	Pro	Gly	Arg	Trp	Pro	Trp	Gln			
			805						810					815				
Cys	Ser	Leu	Gln	Ser	Glu	Pro	Ser	Gly	His	Ile	Cys	Gly	Cys	Val	Leu			
		820					825						830					
Ile	Ala	Lys	Lys	Trp	Val	Leu	Thr	Val	Ala	His	Cys	Phe	Glu	Gly	Arg			
		835					840					845						
Glu	Asn	Ala	Ala	Val	Trp	Lys	Val	Val	Leu	Gly	Ile	Asn	Asn	Leu	Asp			
		850				855					860							
His	Pro	Ser	Val	Phe	Met	Gln	Thr	Arg	Phe	Val	Lys	Thr	Ile	Ile	Leu			
865					870				875					880				
His	Pro	Arg	Tyr	Ser	Arg	Ala	Val	Val	Asp	Tyr	Asp	Ile	Ser	Ile	Val			
			885						890					895				
Glu	Leu	Ser	Glu	Asp	Ile	Ser	Glu	Thr	Gly	Tyr	Val	Arg	Pro	Val	Cys			
			900				905						910					
Leu	Pro	Asn	Pro	Glu	Gln	Trp	Leu	Glu	Pro	Asp	Thr	Tyr	Cys	Tyr	Ile			
		915					920						925					
Thr	Gly	Trp	Gly	His	Met	Gly	Asn	Lys	Met	Pro	Phe	Lys	Leu	Gln	Glu			
		930				935					940							
Gly	Glu	Val	Arg	Ile	Ile	Ser	Leu	Glu	His	Cys	Gln	Ser	Tyr	Phe	Asp			
945					950				955					960				
Met	Lys	Thr	Ile	Thr	Thr	Arg	Met	Ile	Cys	Ala	Gly	Tyr	Glu	Ser	Gly			
			965						970					975				
Thr	Val	Asp	Ser	Cys	Met	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	Cys	Glu			
			980					985					990					
Lys	Pro	Gly	Gly	Arg	Trp	Thr	Leu	Phe	Gly	Leu	Thr	Ser	Trp	Gly	Ser			
		995				1000						1005						
Val	Cys	Phe	Ser	Lys	Val	Leu	Gly	Pro	Gly	Val	Tyr	Ser	Asn	Val	Ser			
		1010				1015					1020							
Tyr	Phe	Val	Glu	Trp	Ile	Lys	Arg	Gln	Ile	Tyr	Ile	Gln	Thr	Phe	Leu			
1025					1030					1035				1040				
Leu	Asn																	

<210> 63

<211> 3696

<212> DNA

<213> Homo Sapien

<220>

<221> CDS

<222> (41)...(3100)

<223> Nucleotide sequence encoding human entorkinase

<300>

<308> GenBank HSU09860

<309> 1995-06-03

<400> 63

```

accagacagt tcttaaatta gcaagccttc aaaacacaaa atg ggg tcg aaa aga      55
                                     Met Gly Ser Lys Arg
                                     1       5

ggc ata tct tct agg cat cat tct ctc agc tcc tat gaa atc atg ttt      103
Gly Ile Ser Ser Arg His His Ser Leu Ser Ser Tyr Glu Ile Met Phe
                                10                20

gca gct ctc ttt gcc ata ttg gta gtg ctc tgt gct gga tta att gca      151
Ala Ala Leu Phe Ala Ile Leu Val Val Leu Cys Ala Gly Leu Ile Ala
                                25                30                35

gta tcc tgc ctg aca atc aag gaa tcc caa cga ggt gca gca ctt gga      199
Val Ser Cys Leu Thr Ile Lys Glu Ser Gln Arg Gly Ala Ala Leu Gly
                                40                45                50

cag agt cat gaa gcc aga gcg aca ttt aaa ata aca tcc gga gtt aca      247
Gln Ser His Glu Ala Arg Ala Thr Phe Lys Ile Thr Ser Gly Val Thr
                                55                60                65

tat aat cct aat ttg caa gac aaa ctc tca gtg gat ttc aaa gtt ctt      295
Tyr Asn Pro Asn Leu Gln Asp Lys Leu Ser Val Asp Phe Lys Val Leu
                                70                75                80                85

gct ttt gac ctt cag caa atg ata gat gag atc ttt cta tca agc aat      343
Ala Phe Asp Leu Gln Gln Met Ile Asp Glu Ile Phe Leu Ser Ser Asn
                                90                95                100

ctg aag aat gaa tat aag aac tca aga gtt tta caa ttt gaa aat ggc      391
Leu Lys Asn Glu Tyr Lys Asn Ser Arg Val Leu Gln Phe Glu Asn Gly
                                105                110                115

agc att ata gtc gta ttt gac ctt ttc ttt gcc cag tgg gtg tca gat      439
Ser Ile Ile Val Val Phe Asp Leu Phe Phe Ala Gln Trp Val Ser Asp
                                120                125                130

caa aat gta aaa gaa gaa ctg att caa ggc ctt gaa gca aat aaa tcc      487
Gln Asn Val Lys Glu Glu Leu Ile Gln Gly Leu Glu Ala Asn Lys Ser
                                135                140                145

agc caa ctg gtc act ttc cat att gat ttg aac agc gtt gat atc cta      535
Ser Gln Leu Val Thr Phe His Ile Asp Leu Asn Ser Val Asp Ile Leu
                                150                155                160                165

gac aag cta aca acc acc agt cat ctg gca act cca gga aat gtc tca      583
Asp Lys Leu Thr Thr Thr Ser His Leu Ala Thr Pro Gly Asn Val Ser
                                170                175                180

ata gag tgc ctg cct ggt tca agt cct tgt act gat gct cta acg tgt      631
Ile Glu Cys Leu Pro Gly Ser Ser Pro Cys Thr Asp Ala Leu Thr Cys
                                185                190                195

ata aaa gct gat tta ttt tgt gat gga gaa gta aac tgt cca gat ggt      679
Ile Lys Ala Asp Leu Phe Cys Asp Gly Glu Val Asn Cys Pro Asp Gly
                                200                205                210

tct gac gaa gac aat aaa atg tgt gcc aca gtt tgt gat gga aga ttt      727

```

Ser 215	Asp	Glu	Asp	Asn	Lys	Met 220	Cys	Ala	Thr	Val	Cys 225	Asp	Gly	Arg	Phe	
ttg	tta	act	gga	tca	tct	ggg	tct	ttc	cag	gct	act	cat	tat	cca	aaa	775
Leu 230	Leu	Thr	Gly	Ser	Ser 235	Gly	Ser	Phe	Gln	Ala 240	Thr	His	Tyr	Pro	Lys 245	
cct	tct	gaa	aca	agt	gtt	gtc	tgc	cag	tgg	atc	ata	cgt	gta	aac	caa	823
Pro	Ser	Glu	Thr	Ser 250	Val	Val	Cys	Gln	Trp 255	Ile	Ile	Arg	Val	Asn 260	Gln	
gga	ctt	tcc	att	aaa	ctg	agc	ttc	gat	gat	ttt	aat	aca	tat	tat	aca	871
Gly	Leu	Ser	Ile 265	Lys	Leu	Ser	Phe	Asp 270	Asp	Phe	Asn	Thr	Tyr 275	Tyr	Thr	
gat	ata	tta	gat	att	tat	gaa	ggg	gta	gga	tca	agc	aag	att	tta	aga	919
Asp	Ile	Leu	Asp 280	Ile	Tyr	Glu	Gly 285	Val	Gly	Ser	Ser	Lys	Ile	Leu	Arg	
gct	tct	att	tgg	gaa	act	aat	cct	ggc	aca	ata	aga	att	ttt	tcc	aac	967
Ala	Ser	Ile	Trp	Glu	Thr	Asn 300	Pro	Gly	Thr	Ile	Arg 305	Ile	Phe	Ser	Asn	
caa	gtt	act	gcc	acc	ttt	ctt	ata	gaa	tct	gat	gaa	agt	gat	tat	gtt	1015
Gln 310	Val	Thr	Ala	Thr	Phe 315	Leu	Ile	Glu	Ser	Asp 320	Glu	Ser	Asp	Tyr	Val 325	
ggc	ttt	aat	gca	aca	tat	act	gca	ttt	aac	agc	agt	gag	ctt	aat	aat	1063
Gly	Phe	Asn	Ala 330	Thr	Tyr	Thr	Ala	Phe	Asn 335	Ser	Ser	Ser	Glu	Leu	Asn 340	
tat	gag	aaa	att	aat	tgt	aac	ttt	gag	gat	ggc	ttt	tgt	ttc	tgg	gtc	1111
Tyr	Glu	Lys	Ile 345	Asn	Cys	Asn	Phe	Glu	Asp 350	Gly	Phe	Cys	Phe	Trp	Val 355	
cag	gat	cta	aat	gat	gat	aat	gaa	tgg	gaa	agg	att	cag	gga	agc	acc	1159
Gln	Asp	Leu	Asn 360	Asp	Asp	Asn	Glu 365	Trp	Glu	Arg	Ile	Gln 370	Gly	Ser	Thr	
ttt	tct	cct	ttt	act	gga	ccc	aat	ttt	gac	cac	act	ttt	ggc	aat	gct	1207
Phe	Ser	Pro	Phe 375	Thr	Gly	Pro 380	Asn	Phe	Asp	His	Thr 385	Phe	Gly	Asn	Ala	
tca	gga	ttt	tac	att	tct	acc	cca	act	gga	cca	gga	ggg	aga	caa	gaa	1255
Ser	Gly	Phe	Tyr	Ile	Ser 395	Thr	Pro	Thr	Gly	Pro 400	Gly	Gly	Arg	Gln	Glu 405	
cga	gtg	ggg	ctt	tta	agc	ctc	cct	ttg	gac	ccc	act	ttg	gag	cca	gct	1303
Arg	Val	Gly	Leu 410	Leu	Ser	Leu	Pro	Leu	Asp 415	Pro	Thr	Leu	Glu	Pro 420	Ala	
tgc	ctt	agt	ttc	tgg	tat	cat	atg	tat	ggg	gaa	aat	gtc	cat	aaa	tta	1351
Cys	Leu	Ser	Phe 425	Trp	Tyr	His	Met	Tyr 430	Gly	Glu	Asn	Val	His 435	Lys	Leu	
agc	att	aat	atc	agc	aat	gac	caa	aat	atg	gag	aag	aca	gtt	ttc	caa	1399
Ser	Ile	Asn 440	Ile	Ser	Asn	Asp	Gln 445	Asn	Met	Glu	Lys	Thr 450	Val	Phe	Gln	
aag	gaa	gga	aat	tat	gga	gac	aat	tgg	aat	tat	gga	caa	gta	acc	cta	1447
Lys	Glu	Gly	Asn 455	Tyr	Gly	Asp 460	Asn	Trp	Asn	Tyr	Gly 465	Gln	Val	Thr	Leu	

aat gaa aca gtt .aaa ttt aag gtt gct ttt aat gct ttt aaa aac aag Asn Glu Thr Val Lys Phe Lys Val Ala Phe Asn Ala Phe Lys Asn Lys 470 475 480 485	1495
atc ctg agt gat att gcg ttg gat gac att agc cta aca tat ggg att Ile Leu Ser Asp Ile Ala Leu Asp Asp Ile Ser Leu Thr Tyr Gly Ile 490 495 500	1543
tgc aat ggg agt ctt tat cca gaa cca act ttg gtg cca act cct cca Cys Asn Gly Ser Leu Tyr Pro Glu Pro Thr Leu Val Pro Thr Pro Pro 505 510 515	1591
cca gaa ctt cct acg gac tgt gga gga cct ttt gag ctg tgg gag cca Pro Glu Leu Pro Thr Asp Cys Gly Gly Pro Phe Glu Leu Trp Glu Pro 520 525 530	1639
aat aca aca ttc agt tct acg aac ttt cca aac agc tac cct aat ctg Asn Thr Thr Phe Ser Ser Thr Asn Phe Pro Asn Ser Tyr Pro Asn Leu 535 540 545	1687
gct ttc tgt gtt tgg att tta aat gca caa aaa gga aag aat ata caa Ala Phe Cys Val Trp Ile Leu Asn Ala Gln Lys Gly Lys Asn Ile Gln 550 555 560 565	1735
ctt cat ttt caa gaa ttt gac tta gaa aat att aac gat gta gtt gaa Leu His Phe Gln Glu Phe Asp Leu Glu Asn Ile Asn Asp Val Val Glu 570 575 580	1783
ata aga gat ggt gaa gaa gct gat tcc ttg ctc tta gct gtg tac aca Ile Arg Asp Gly Glu Glu Ala Asp Ser Leu Leu Leu Ala Val Tyr Thr 585 590 595	1831
ggg cct ggc cca gta aag gat gtg ttc tct acc acc aac aga atg act Gly Pro Gly Pro Val Lys Asp Val Phe Ser Thr Thr Asn Arg Met Thr 600 605 610	1879
gtg ctt ctc atc act aac gat gtg ttg gca aga gga ggg ttt aaa gca Val Leu Leu Ile Thr Asn Asp Val Leu Ala Arg Gly Gly Phe Lys Ala 615 620 625	1927
aac ttt act act ggc tat cac ttg ggg att cca gag cca tgc aag gca Asn Phe Thr Thr Gly Tyr His Leu Gly Ile Pro Glu Pro Cys Lys Ala 630 635 640 645	1975
gac cat ttt caa tgt aaa aat gga gag tgt gtt cca ctg gtg aat ctc Asp His Phe Gln Cys Lys Asn Gly Glu Cys Val Pro Leu Val Asn Leu 650 655 660	2023
tgt gac ggt cat ctg cac tgt gag gat ggc tca gat gaa gca gat tgt Cys Asp Gly His Leu His Cys Glu Asp Gly Ser Asp Glu Ala Asp Cys 665 670 675	2071
gtg cgt ttt ttc aat ggc aca acg aac aac aat ggt tta gtg cgg ttc Val Arg Phe Phe Asn Gly Thr Thr Asn Asn Asn Gly Leu Val Arg Phe 680 685 690	2119
aga atc cag agc ata tgg cat aca gct tgt gct gag aac tgg acc acc Arg Ile Gln Ser Ile Trp His Thr Ala Cys Ala Glu Asn Trp Thr Thr 695 700 705	2167
cag att tca aat gat gtt tgt caa ctg ctg gga cta ggg agt gga aac	2215

Gln 710	Ile	Ser	Asn	Asp	Val 715	Cys	Gln	Leu	Leu	Gly 720	Leu	Gly	Ser	Gly	Asn 725	
tca	tca	aag	cca	atc	ttc	tct	acc	gat	ggg	gga	cca	ttt	gtc	aaa	tta	2263
Ser	Ser	Lys	Pro	Ile	Phe	Ser	Thr	Asp	Gly	Gly	Pro	Phe	Val	Lys	Leu	
				730					735					740		
aac	aca	gca	cct	gat	ggc	cac	tta	ata	cta	aca	ccc	agt	caa	cag	tgt	2311
Asn	Thr	Ala	Pro	Asp	Gly	His	Leu	Ile	Leu	Thr	Pro	Ser	Gln	Gln	Cys	
			745					750						755		
tta	cag	gat	tcc	ttg	att	cgg	tta	cag	tgt	aac	cat	aaa	tct	tgt	gga	2359
Leu	Gln	Asp	Ser	Leu	Ile	Arg	Leu	Gln	Cys	Asn	His	Lys	Ser	Cys	Gly	
			760					765						770		
aaa	aaa	ctg	gca	gct	caa	gac	atc	acc	cca	aag	att	gtt	gga	gga	agt	2407
Lys	Lys	Leu	Ala	Ala	Gln	Asp	Ile	Thr	Pro	Lys	Ile	Val	Gly	Gly	Ser	
		775				780										
aat	gcc	aaa	gaa	ggg	gcc	tgg	ccc	tgg	gtt	gtg	ggg	ctg	tat	tat	ggc	2455
Asn	Ala	Lys	Glu	Gly	Ala	Trp	Pro	Trp	Val	Val	Gly	Leu	Tyr	Tyr	Gly	
				795					800						805	
ggc	cga	ctg	ctc	tgc	ggc	gca	tct	ctc	gtc	agc	agt	gac	tgg	ctg	gtg	2503
Gly	Arg	Leu	Leu	Cys	Gly	Ala	Ser	Leu	Val	Ser	Ser	Asp	Trp	Leu	Val	
				810					815					820		
tcc	gac	gca	cac	tgc	gtg	tat	ggg	aga	aac	tta	gag	cca	tcc	aag	tgg	2551
Ser	Ala	Ala	His	Cys	Val	Tyr	Gly	Arg	Asn	Leu	Glu	Pro	Ser	Lys	Trp	
			825					830						835		
aca	gca	atc	cta	ggc	ctg	cat	atg	aaa	tca	aac	ctg	acc	tct	cct	caa	2599
Thr	Ala	Ile	Leu	Gly	Leu	His	Met	Lys	Ser	Asn	Leu	Thr	Ser	Pro	Gln	
			840					845						850		
aca	gtc	cct	cga	tta	ata	gat	gaa	att	gtc	ata	aac	cct	cat	tac	aat	2647
Thr	Val	Pro	Arg	Leu	Ile	Asp	Glu	Ile	Val	Ile	Asn	Pro	His	Tyr	Asn	
			855			860						865				
agg	cga	aga	aag	gac	aac	gac	att	gcc	atg	atg	cat	ctg	gaa	ttt	aaa	2695
Arg	Arg	Arg	Lys	Asp	Asn	Asp	Ile	Ala	Met	Met	His	Leu	Glu	Phe	Lys	
				875						880					885	
gtg	aat	tac	aca	gat	tac	ata	caa	cct	att	tgt	tta	ccg	gaa	gaa	aat	2743
Val	Asn	Tyr	Thr	Asp	Tyr	Ile	Gln	Pro	Ile	Cys	Leu	Pro	Glu	Glu	Asn	
				890					895						900	
caa	gtt	ttt	cct	cca	gga	aga	aat	tgt	tct	att	gct	ggg	tgg	ggg	acg	2791
Gln	Val	Phe	Pro	Pro	Gly	Arg	Asn	Cys	Ser	Ile	Ala	Gly	Trp	Gly	Thr	
			905					910						915		
gtt	gta	tat	caa	ggg	act	act	gca	aac	ata	ttg	caa	gaa	gct	gat	gtt	2839
Val	Val	Tyr	Gln	Gly	Thr	Thr	Ala	Asn	Ile	Leu	Gln	Glu	Ala	Asp	Val	
			920				925						930			
cct	ctt	cta	tca	aat	gag	aga	tgc	caa	cag	cag	atg	cca	gaa	tat	aac	2887
Pro	Leu	Leu	Ser	Asn	Glu	Arg	Cys	Gln	Gln	Gln	Met	Pro	Glu	Tyr	Asn	
			935			940					945					
att	act	gaa	aat	atg	ata	tgt	gca	ggc	tat	gaa	gaa	gga	gga	ata	gat	2935
Ile	Thr	Glu	Asn	Met	Ile	Cys	Ala	Gly	Tyr	Glu	Glu	Gly	Gly	Ile	Asp	
					955					960					965	

tct tgt cag ggg gat tca gga gga cca tta atg tgc caa gaa aac aac 2983
 Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Cys Gln Glu Asn Asn
 970 975 980

agg tgg ttc ctt gct ggt gtg acc tca ttt gga tac aag tgt gcc ctg 3031
 Arg Trp Phe Leu Ala Gly Val Thr Ser Phe Gly Tyr Lys Cys Ala Leu
 985 990 995

cct aat cgc ccc gga gtg tat gcc agg gtc tca agg ttt acc gaa tgg 3079
 Pro Asn Arg Pro Gly Val Tyr Ala Arg Val Ser Arg Phe Thr Glu Trp
 1000 1005 1010

ata caa agt ttt cta cat tag cgcatttctt aaactaaaca ggaaagtgcg 3130
 Ile Gln Ser Phe Leu His *
 1015

attattttcc cattctactc tagaaagcat ggaaattaag tgtttcgtac aaaaatttta 3190
 aaaagttacc aaagggttttt attcttacct atgtcaatga aatgctaggg ggccagggaa 3250
 acaaaatttt aaaaataata aaattcacca tagcaataca gaataacttt aaaataccat 3310
 taaatacatt tgtatttcat tgtgaacagg tatttcttca cagatctcat ttttaaaatt 3370
 cttaatgatt atttttatta ctactgttg tttaaaggga tgttatttta aagcatatac 3430
 catacactta agaaatttga gcagaattta aaaaagaaag aaaataaatt gtttttccca 3490
 aagtatgtca ctgttggaata taaactgcca taaattttct agttccagtt tagtttgctg 3550
 ctattagcag aaactcaatt gtttctctgt cttttctatc aaaaattttca acatatgcat 3610
 aaccttagta ttttcccaac caatagaaac tattttattgt aagcttatgt cacaggcctg 3670
 gactaaattg attttacgtt cctctt 3696

<210> 64
 <211> 1019
 <212> PRT
 <213> Homo Sapien

<400> 64
 Met Gly Ser Lys Arg Gly Ile Ser Ser Arg His His Ser Leu Ser Ser
 1 5 10 15
 Tyr Glu Ile Met Phe Ala Ala Leu Phe Ala Ile Leu Val Val Leu Cys
 20 25 30
 Ala Gly Leu Ile Ala Val Ser Cys Leu Thr Ile Lys Glu Ser Gln Arg
 35 40 45
 Gly Ala Ala Leu Gly Gln Ser His Glu Ala Arg Ala Thr Phe Lys Ile
 50 55 60
 Thr Ser Gly Val Thr Tyr Asn Pro Asn Leu Gln Asp Lys Leu Ser Val
 65 70 75 80
 Asp Phe Lys Val Leu Ala Phe Asp Leu Gln Gln Met Ile Asp Glu Ile
 85 90 95
 Phe Leu Ser Ser Asn Leu Lys Asn Glu Tyr Lys Asn Ser Arg Val Leu
 100 105 110
 Gln Phe Glu Asn Gly Ser Ile Ile Val Val Phe Asp Leu Phe Phe Ala
 115 120 125
 Gln Trp Val Ser Asp Gln Asn Val Lys Glu Glu Leu Ile Gln Gly Leu
 130 135 140
 Glu Ala Asn Lys Ser Ser Gln Leu Val Thr Phe His Ile Asp Leu Asn
 145 150 155 160
 Ser Val Asp Ile Leu Asp Lys Leu Thr Thr Thr Ser His Leu Ala Thr
 165 170 175
 Pro Gly Asn Val Ser Ile Glu Cys Leu Pro Gly Ser Ser Pro Cys Thr
 180 185 190
 Asp Ala Leu Thr Cys Ile Lys Ala Asp Leu Phe Cys Asp Gly Glu Val
 195 200 205
 Asn Cys Pro Asp Gly Ser Asp Glu Asp Asn Lys Met Cys Ala Thr Val
 210 215 220

Cys	Asp	Gly	Arg	Phe	Leu	Leu	Thr	Gly	Ser	Ser	Gly	Ser	Phe	Gln	Ala
225					230					235					240
Thr	His	Tyr	Pro	Lys	Pro	Ser	Glu	Thr	Ser	Val	Val	Cys	Gln	Trp	Ile
				245					250					255	
Ile	Arg	Val	Asn	Gln	Gly	Leu	Ser	Ile	Lys	Leu	Ser	Phe	Asp	Asp	Phe
			260					265					270		
Asn	Thr	Tyr	Tyr	Thr	Asp	Ile	Leu	Asp	Ile	Tyr	Glu	Gly	Val	Gly	Ser
		275				280						285			
Ser	Lys	Ile	Leu	Arg	Ala	Ser	Ile	Trp	Glu	Thr	Asn	Pro	Gly	Thr	Ile
	290					295					300				
Arg	Ile	Phe	Ser	Asn	Gln	Val	Thr	Ala	Thr	Phe	Leu	Ile	Glu	Ser	Asp
305				310					315					320	
Glu	Ser	Asp	Tyr	Val	Gly	Phe	Asn	Ala	Thr	Tyr	Thr	Ala	Phe	Asn	Ser
				325					330					335	
Ser	Glu	Leu	Asn	Asn	Tyr	Glu	Lys	Ile	Asn	Cys	Asn	Phe	Glu	Asp	Gly
			340					345					350		
Phe	Cys	Phe	Trp	Val	Gln	Asp	Leu	Asn	Asp	Asp	Asn	Glu	Trp	Glu	Arg
		355					360					365			
Ile	Gln	Gly	Ser	Thr	Phe	Ser	Pro	Phe	Thr	Gly	Pro	Asn	Phe	Asp	His
	370					375					380				
Thr	Phe	Gly	Asn	Ala	Ser	Gly	Phe	Tyr	Ile	Ser	Thr	Pro	Thr	Gly	Pro
385				390					395					400	
Gly	Gly	Arg	Gln	Glu	Arg	Val	Gly	Leu	Leu	Ser	Leu	Pro	Leu	Asp	Pro
				405				410						415	
Thr	Leu	Glu	Pro	Ala	Cys	Leu	Ser	Phe	Trp	Tyr	His	Met	Tyr	Gly	Glu
			420					425					430		
Asn	Val	His	Lys	Leu	Ser	Ile	Asn	Ile	Ser	Asn	Asp	Gln	Asn	Met	Glu
		435					440					445			
Lys	Thr	Val	Phe	Gln	Lys	Glu	Gly	Asn	Tyr	Gly	Asp	Asn	Trp	Asn	Tyr
	450					455					460				
Gly	Gln	Val	Thr	Leu	Asn	Glu	Thr	Val	Lys	Phe	Lys	Val	Ala	Phe	Asn
465				470					475					480	
Ala	Phe	Lys	Asn	Lys	Ile	Leu	Ser	Asp	Ile	Ala	Leu	Asp	Asp	Ile	Ser
			485					490						495	
Leu	Thr	Tyr	Gly	Ile	Cys	Asn	Gly	Ser	Leu	Tyr	Pro	Glu	Pro	Thr	Leu
			500					505					510		
Val	Pro	Thr	Pro	Pro	Pro	Glu	Leu	Pro	Thr	Asp	Cys	Gly	Gly	Pro	Phe
		515					520					525			
Glu	Leu	Trp	Glu	Pro	Asn	Thr	Thr	Phe	Ser	Ser	Thr	Asn	Phe	Pro	Asn
	530					535					540				
Ser	Tyr	Pro	Asn	Leu	Ala	Phe	Cys	Val	Trp	Ile	Leu	Asn	Ala	Gln	Lys
545				550						555				560	
Gly	Lys	Asn	Ile	Gln	Leu	His	Phe	Gln	Glu	Phe	Asp	Leu	Glu	Asn	Ile
			565						570					575	
Asn	Asp	Val	Val	Glu	Ile	Arg	Asp	Gly	Glu	Glu	Ala	Asp	Ser	Leu	Leu
			580					585					590		
Leu	Ala	Val	Tyr	Thr	Gly	Pro	Gly	Pro	Val	Lys	Asp	Val	Phe	Ser	Thr
		595					600					605			
Thr	Asn	Arg	Met	Thr	Val	Leu	Leu	Ile	Thr	Asn	Asp	Val	Leu	Ala	Arg
	610					615					620				
Gly	Gly	Phe	Lys	Ala	Asn	Phe	Thr	Thr	Gly	Tyr	His	Leu	Gly	Ile	Pro
625				630					635					640	
Glu	Pro	Cys	Lys	Ala	Asp	His	Phe	Gln	Cys	Lys	Asn	Gly	Glu	Cys	Val
			645						650					655	
Pro	Leu	Val	Asn	Leu	Cys	Asp	Gly	His	Leu	His	Cys	Glu	Asp	Gly	Ser
			660				665						670		
Asp	Glu	Ala	Asp	Cys	Val	Arg	Phe	Asn	Gly	Thr	Thr	Asn	Asn	Asn	
		675					680					685			
Gly	Leu	Val	Arg	Phe	Arg	Ile	Gln	Ser	Ile	Trp	His	Thr	Ala	Cys	Ala
	690					695					700				
Glu	Asn	Trp	Thr	Thr	Gln	Ile	Ser	Asn	Asp	Val	Cys	Gln	Leu	Leu	Gly
705					710					715					720

Leu Gly Ser Gly Asn Ser Ser Lys Pro Ile Phe Ser Thr Asp Gly Gly
 725 730 735
 Pro Phe Val Lys Leu Asn Thr Ala Pro Asp Gly His Leu Ile Leu Thr
 740 745 750
 Pro Ser Gln Gln Cys Leu Gln Asp Ser Leu Ile Arg Leu Gln Cys Asn
 755 760 765
 His Lys Ser Cys Gly Lys Lys Leu Ala Ala Gln Asp Ile Thr Pro Lys
 770 775 780
 Ile Val Gly Gly Ser Asn Ala Lys Glu Gly Ala Trp Pro Trp Val Val
 785 790 795 800
 Gly Leu Tyr Tyr Gly Gly Arg Leu Leu Cys Gly Ala Ser Leu Val Ser
 805 810 815
 Ser Asp Trp Leu Val Ser Ala Ala His Cys Val Tyr Gly Arg Asn Leu
 820 825 830
 Glu Pro Ser Lys Trp Thr Ala Ile Leu Gly Leu His Met Lys Ser Asn
 835 840 845
 Leu Thr Ser Pro Gln Thr Val Pro Arg Leu Ile Asp Glu Ile Val Ile
 850 855 860
 Asn Pro His Tyr Asn Arg Arg Lys Asp Asn Asp Ile Ala Met Met
 865 870 875 880
 His Leu Glu Phe Lys Val Asn Tyr Thr Asp Tyr Ile Gln Pro Ile Cys
 885 890 895
 Leu Pro Glu Glu Asn Gln Val Phe Pro Pro Gly Arg Asn Cys Ser Ile
 900 905 910
 Ala Gly Trp Gly Thr Val Val Tyr Gln Gly Thr Thr Ala Asn Ile Leu
 915 920 925
 Gln Glu Ala Asp Val Pro Leu Leu Ser Asn Glu Arg Cys Gln Gln Gln
 930 935 940
 Met Pro Glu Tyr Asn Ile Thr Glu Asn Met Ile Cys Ala Gly Tyr Glu
 945 950 955 960
 Glu Gly Gly Ile Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Met
 965 970 975
 Cys Gln Glu Asn Asn Arg Trp Phe Leu Ala Gly Val Thr Ser Phe Gly
 980 985 990
 Tyr Lys Cys Ala Leu Pro Asn Arg Pro Gly Val Tyr Ala Arg Val Ser
 995 1000 1005
 Arg Phe Thr Glu Trp Ile Gln Ser Phe Leu His
 1010 1015

<210> 65
 <211> 1500
 <212> DNA
 <213> Homo Sapien

<220>
 <221> CDS
 <222> (62)...(1318)
 <223> Nucleotide sequence encoding human airway
 trypsin-like protease

<300>
 <308> GenBank AB002134
 <309> 1998-06-04

<400> 65
 gagtggaat ctcaaagcag ttgagtaggc agaaaaaaga acctcttcat taaggattaa 60
 a atg tat agg cca gca cgt gta act tcg act tca aga ttt ctg aat cca 109
 Met Tyr Arg Pro Ala Arg Val Thr Ser Thr Ser Arg Phe Leu Asn Pro
 1 5 10 15
 tat gta gta tgt ttc att gtc gtc gca ggg gta gtg atc ctg gca gtc 157
 Tyr Val Val Cys Phe Ile Val Val Ala Gly Val Val Ile Leu Ala Val

acc Thr	ata Ile	gct Ala 35	cta Leu	ctt Leu	gtt Val	tac Tyr	ttt Phe 40	tta Leu	gct Ala	ttt Phe	gat Asp	caa Gln 45	aaa Lys	tct Ser	tac Tyr	205
ttt Phe	tat Tyr 50	agg Arg	agc Ser	agt Ser	ttt Phe	caa Gln 55	ctc Leu	cta Leu	aat Asn	gtt Val	gaa Glu 60	tat Tyr	aat Asn	agt Ser	cag Gln	253
tta Leu 65	aat Asn	tca Ser	cca Pro	gct Ala	aca Thr 70	cag Gln	gaa Glu	tac Tyr	agg Arg	act Thr 75	ttg Leu	agt Ser	gga Gly	aga Arg	att Ile 80	301
gaa Glu	tct Ser	ctg Leu	att Ile	act Thr 85	aaa Lys	aca Thr	ttc Phe	aaa Lys	gaa Glu 90	tca Ser	aat Asn	tta Leu	aga Arg	aat Asn 95	cag Gln	349
ttc Phe	atc Ile	aga Arg	gct Ala 100	cat His	gtt Val	gcc Ala	aaa Lys	ctg Leu 105	agg Arg	caa Gln	gat Asp	ggg Gly 110	agt Ser	ggg Gly	gtg Val	397
aga Arg	gcg Ala	gat Asp 115	gtt Val	gtc Val	atg Met	aaa Lys	ttt Phe 120	caa Gln	ttc Phe	act Thr	aga Arg	aat Asn 125	aac Asn	aat Asn	gga Gly	445
gca Ala	tca Ser 130	atg Met	aaa Lys	agc Ser	aga Arg	att Ile 135	gag Glu	tct Ser	gtt Val	tta Leu 140	cga Arg	caa Gln	atg Met	ctg Leu	aat Asn	493
aac Asn 145	tct Ser	gga Gly	aac Asn	ctg Leu	gaa Glu 150	ata Ile	aac Asn	cct Pro	tca Ser	act Thr 155	gag Glu	ata Ile	aca Thr	tca Ser	ctt Leu 160	541
act Thr	gac Asp	cag Gln	gct Ala	gca Ala 165	gca Ala	aat Asn	tgg Trp	ctt Leu 170	att Ile	aat Asn	gaa Glu	tgt Cys	ggg Gly	gcc Ala 175	ggg Gly	589
cca Pro	gac Asp	cta Leu	ata Ile 180	aca Thr	ttg Leu	tct Ser	gag Glu	cag Gln 185	aga Arg	atc Ile	ctt Leu	gga Gly	ggc Gly 190	act Thr	gag Glu	637
gct Ala	gag Glu	gag Glu 195	gga Gly	agc Ser	tgg Trp	ccg Pro	tgg Trp 200	caa Gln	gtc Val	agt Ser	ctg Leu	cgg Arg 205	ctc Leu	aat Asn	aat Asn	685
gcc Ala	cac His 210	cac His	tgt Cys	gga Gly	ggc Gly	agc Ser 215	ctg Leu	atc Ile	aat Asn	aac Asn 220	atg Met	tgg Trp	atc Ile	ctg Leu	aca Thr	733
gca Ala 225	gct Ala	cac His	tgc Cys	ttc Phe	aga Arg 230	agc Ser	aac Asn	tct Ser	aat Asn	cct Pro 235	cgt Arg	gac Asp	tgg Trp	att Ile	gcc Ala 240	781
acg Thr	tct Ser	ggg Gly	att Ile	tcc Ser 245	aca Thr	aca Thr	ttt Phe	cct Pro	aaa Lys 250	cta Leu	aga Arg	atg Met	aga Arg	gta Val 255	aga Arg	829
aat Asn	att Ile	tta Leu	att Ile 260	cat His	aac Asn	aat Asn	tat Tyr 265	aaa Lys 265	tct Ser	gca Ala	act Thr	cat His	gaa Glu 270	aat Asn	gac Asp	877

att gca ctt gtg aga ctt gag aac agt gtc acc ttt acc aaa gat atc 925
 Ile Ala Leu Val Arg Leu Glu Asn Ser Val Thr Phe Thr Lys Asp Ile
 275 280 285

cat agt gtg tgt ctc cca gct gct acc cag aat att cca cct ggc tct 973
 His Ser Val Cys Leu Pro Ala Ala Thr Gln Asn Ile Pro Pro Gly Ser
 290 295 300

act gct tat gta aca gga tgg ggc gct caa gaa tat gct ggc cac aca 1021
 Thr Ala Tyr Val Thr Gly Trp Gly Ala Gln Glu Tyr Ala Gly His Thr
 305 310 315 320

gtt cca gag cta agg caa gga cag gtc aga ata ata agt aat gat gta 1069
 Val Pro Glu Leu Arg Gln Gly Gln Val Arg Ile Ile Ser Asn Asp Val
 325 330 335

tgt aat gca cca cat agt tat aat gga gcc atc ttg tct gga atg ctg 1117
 Cys Asn Ala Pro His Ser Tyr Asn Gly Ala Ile Leu Ser Gly Met Leu
 340 345 350

tgt gct gga gta cct caa ggt gga gtg gac gca tgt cag ggt gac tct 1165
 Cys Ala Gly Val Pro Gln Gly Gly Val Asp Ala Cys Gln Gly Asp Ser
 355 360 365

ggt ggc cca cta gta caa gaa gac tca cgg cgg ctt tgg ttt att gtg 1213
 Gly Gly Pro Leu Val Gln Glu Asp Ser Arg Arg Leu Trp Phe Ile Val
 370 375 380

ggg ata gta agc tgg gga gat cag tgt ggc ctg ccg gat aag cca gga 1261
 Gly Ile Val Ser Trp Gly Asp Gln Cys Gly Leu Pro Asp Lys Pro Gly
 385 390 395 400

gtg tat act cga gtg aca gcc tac ctt gac tgg att agg caa caa act 1309
 Val Tyr Thr Arg Val Thr Ala Tyr Leu Asp Trp Ile Arg Gln Gln Thr
 405 410 415

ggg atc tag tgcaacaagt gcatccctgt tgcaaagtct gtatgcaggt 1358
 Gly Ile *

gtgcctgtct taaattccaa agctttacat ttcaactgaa aaagaaacta gaaatgtcct 1418
 aatttaacat cttgtttacat aaatatgggt taacaaacac tgtttaacct ttctttatta 1478
 ttaaagggtt tctattttct cc 1500

<210> 66

<211> 418

<212> PRT

<213> Homo Sapien

<400> 66

Met Tyr Arg Pro Ala Arg Val Thr Ser Thr Ser Arg Phe Leu Asn Pro
 1 5 10 15
 Tyr Val Val Cys Phe Ile Val Val Ala Gly Val Val Ile Leu Ala Val
 20 25 30
 Thr Ile Ala Leu Leu Val Tyr Phe Leu Ala Phe Asp Gln Lys Ser Tyr
 35 40 45
 Phe Tyr Arg Ser Ser Phe Gln Leu Leu Asn Val Glu Tyr Asn Ser Gln
 50 55 60
 Leu Asn Ser Pro Ala Thr Gln Glu Tyr Arg Thr Leu Ser Gly Arg Ile
 65 70 75 80
 Glu Ser Leu Ile Thr Lys Thr Phe Lys Glu Ser Asn Leu Arg Asn Gln
 85 90 95

Phe Ile Arg Ala His Val Ala Lys Leu Arg Gln Asp Gly Ser Gly Val
 100 105 110
 Arg Ala Asp Val Val Met Lys Phe Gln Phe Thr Arg Asn Asn Gly
 115 120 125
 Ala Ser Met Lys Ser Arg Ile Glu Ser Val Leu Arg Gln Met Leu Asn
 130 135 140
 Asn Ser Gly Asn Leu Glu Ile Asn Pro Ser Thr Glu Ile Thr Ser Leu
 145 150 155 160
 Thr Asp Gln Ala Ala Asn Trp Leu Ile Asn Glu Cys Gly Ala Gly
 165 170 175
 Pro Asp Leu Ile Thr Leu Ser Glu Gln Arg Ile Leu Gly Gly Thr Glu
 180 185 190
 Ala Glu Glu Gly Ser Trp Pro Trp Gln Val Ser Leu Arg Leu Asn Asn
 195 200 205
 Ala His His Cys Gly Gly Ser Leu Ile Asn Asn Met Trp Ile Leu Thr
 210 215 220
 Ala Ala His Cys Phe Arg Ser Asn Ser Asn Pro Arg Asp Trp Ile Ala
 225 230 235 240
 Thr Ser Gly Ile Ser Thr Thr Phe Pro Lys Leu Arg Met Arg Val Arg
 245 250 255
 Asn Ile Leu Ile His Asn Asn Tyr Lys Ser Ala Thr His Glu Asn Asp
 260 265 270
 Ile Ala Leu Val Arg Leu Glu Asn Ser Val Thr Phe Thr Lys Asp Ile
 275 280 285
 His Ser Val Cys Leu Pro Ala Ala Thr Gln Asn Ile Pro Pro Gly Ser
 290 295 300
 Thr Ala Tyr Val Thr Gly Trp Gly Ala Gln Glu Tyr Ala Gly His Thr
 305 310 315 320
 Val Pro Glu Leu Arg Gln Gly Gln Val Arg Ile Ile Ser Asn Asp Val
 325 330 335
 Cys Asn Ala Pro His Ser Tyr Asn Gly Ala Ile Leu Ser Gly Met Leu
 340 345 350
 Cys Ala Gly Val Pro Gln Gly Gly Val Asp Ala Cys Gln Gly Asp Ser
 355 360 365
 Gly Gly Pro Leu Val Gln Glu Asp Ser Arg Arg Leu Trp Phe Ile Val
 370 375 380
 Gly Ile Val Ser Trp Gly Asp Gln Cys Gly Leu Pro Asp Lys Pro Gly
 385 390 395 400
 Val Tyr Thr Arg Val Thr Ala Tyr Leu Asp Trp Ile Arg Gln Gln Thr
 405 410 415
 Gly Ile

<210> 67
 <211> 1783
 <212> DNA
 <213> Homo Sapien

<220>
 <221> CDS
 <222> (246)...(1499)
 <223> Nucleotide sequence encoding human hepsin

<300>
 <308> GenBank M18930
 <309> 1993-06-11

<400> 67
 tcgagcccgc tttccaggga cctacctga gggccacag gtgaggcagc ctggcctagc 60
 agggcccacg ccaccgcctc tgctccagg ccgcccgtg ctgcggggcc accatgctcc 120
 tgcccaggcc tggagactga cccgaccctg gcactacctc gaggctccgc cccacctgc 180
 tggaccccag ggtcccaccc tggcccagga ggtagccag ggaatcatta acaagaggca 240

gtgac atg gcg cag aag gag ggt ggc cgg act gtg cca tgc tgc tcc aga	290
Met Ala Gln Lys Glu Gly Gly Arg Thr Val Pro Cys Cys Ser Arg	
1 5 10 15	
ccc aag gtg gca gct ctc act gcg ggg acc ctg cta ctt ctg aca gcc	338
Pro Lys Val Ala Ala Leu Thr Ala Gly Thr Leu Leu Leu Thr Ala	
20 25 30	
atc ggg gcg gca tcc tgg gcc att gtg gct gtt ctc ctc agg agt gac	386
Ile Gly Ala Ala Ser Trp Ala Ile Val Ala Val Leu Leu Arg Ser Asp	
35 40 45	
cag gag ccg ctg tac cca gtg cag gtc agc tct gcg gac gct cgg ctc	434
Gln Glu Pro Leu Tyr Pro Val Gln Val Ser Ser Ala Asp Ala Arg Leu	
50 55 60	
atg gtc ttt gac aag acg gaa ggg acg tgg cgg ctg ctg tgc tcc tcg	482
Met Val Phe Asp Lys Thr Glu Gly Thr Trp Arg Leu Leu Cys Ser Ser	
65 70 75	
cgc tcc aac gcc agg gta gcc gga ctc agc tgc gag gag atg ggc ttc	530
Arg Ser Asn Ala Arg Val Ala Gly Leu Ser Cys Glu Glu Met Gly Phe	
80 85 90 95	
ctc agg gca ctg acc cac tcc gag ctg gac gtg cga acg gcg ggc gcc	578
Leu Arg Ala Leu Thr His Ser Glu Leu Asp Val Arg Thr Ala Gly Ala	
100 105 110	
aat ggc acg tcg ggc ttc ttc tgt gtg gac gag ggg agg ctg ccc cac	626
Asn Gly Thr Ser Gly Phe Phe Cys Val Asp Glu Gly Arg Leu Pro His	
115 120 125	
acc cag agg ctg ctg gag gtc atc tcc gtg tgt gat tgc ccc aga ggc	674
Thr Gln Arg Leu Leu Glu Val Ile Ser Val Cys Asp Cys Pro Arg Gly	
130 135 140	
cgt ttc ttg gcc gcc atc tgc caa gac tgt ggc cgc agg aag ctg ccc	722
Arg Phe Leu Ala Ala Ile Cys Gln Asp Cys Gly Arg Arg Lys Leu Pro	
145 150 155	
gtg gac cgc atc gtg gga ggc cgg gac acc agc ttg ggc cgg tgg ccg	770
Val Asp Arg Ile Val Gly Gly Arg Asp Thr Ser Leu Gly Arg Trp Pro	
160 165 170 175	
tgg caa gtc agc ctt cgc tat gat gga gca cac ctc tgt ggg gga tcc	818
Trp Gln Val Ser Leu Arg Tyr Asp Gly Ala His Leu Cys Gly Gly Ser	
180 185 190	
ctg ctc tcc ggg gac tgg gtg ctg aca gcc gcc cac tgc ttc ccg gag	866
Leu Leu Ser Gly Asp Trp Val Leu Thr Ala Ala His Cys Phe Pro Glu	
195 200 205	
cgg aac cgg gtc ctg tcc cga tgg cga gtg ttt gcc ggt gcc gtg gcc	914
Arg Asn Arg Val Leu Ser Arg Trp Arg Val Phe Ala Gly Ala Val Ala	
210 215 220	
cag gcc tct ccc cac ggt ctg cag ctg ggg gtg cag gct gtg gtc tac	962
Gln Ala Ser Pro His Gly Leu Gln Leu Gly Val Gln Ala Val Val Tyr	
225 230 235	
cac ggg ggc tat ctt ccc ttt cgg gac ccc aac agc gag gag aac agc	1010
His Gly Gly Tyr Leu Pro Phe Arg Asp Pro Asn Ser Glu Glu Asn Ser	

240	245	250	255	
aac gat att gcc ctg gtc cac ctc tcc agt ccc ctg ccc ctc aca gaa				1058
Asn Asp Ile Ala Leu Val His Leu Ser Ser Pro Leu Pro Leu Thr Glu				
	260	265	270	
tac atc cag cct gtg tgc ctc cca gct gcc ggc cag gcc ctg gtg gat				1106
Tyr Ile Gln Pro Val Cys Leu Pro Ala Ala Gly Gln Ala Leu Val Asp				
	275	280	285	
ggc aag atc tgt acc gtg acg ggc tgg ggc aac acg cag tac tat ggc				1154
Gly Lys Ile Cys Thr Val Thr Gly Trp Gly Asn Thr Gln Tyr Tyr Gly				
	290	295	300	
caa cag gcc ggg gta ctc cag gag gct cga gtc ccc ata atc agc aat				1202
Gln Gln Ala Gly Val Leu Gln Glu Ala Arg Val Pro Ile Ile Ser Asn				
	305	310	315	
gat gtc tgc aat ggc gct gac ttc tat gga aac cag atc aag ccc aag				1250
Asp Val Cys Asn Gly Ala Asp Phe Tyr Gly Asn Gln Ile Lys Pro Lys				
	320	325	330	335
atg ttc tgt gct ggc tac ccc gag ggt ggc att gat gcc tgc cag ggc				1298
Met Phe Cys Ala Gly Tyr Pro Glu Gly Gly Ile Asp Ala Cys Gln Gly				
	340	345	350	
gac agc ggt ggt ccc ttt gtg tgt gag gac agc atc tct cgg acg cca				1346
Asp Ser Gly Gly Pro Phe Val Cys Glu Asp Ser Ile Ser Arg Thr Pro				
	355	360	365	
cgt tgg cgg ctg tgt ggc att gtg agt tgg ggc act ggc tgt gcc ctg				1394
Arg Trp Arg Leu Cys Gly Ile Val Ser Trp Gly Thr Gly Cys Ala Leu				
	370	375	380	
gcc cag aag cca ggc gtc tac acc aaa gtc agt gac ttc cgg gag tgg				1442
Ala Gln Lys Pro Gly Val Tyr Thr Lys Val Ser Asp Phe Arg Glu Trp				
	385	390	395	
atc ttc cag gcc ata aag act cac tcc gaa gcc agc ggc atg gtg acc				1490
Ile Phe Gln Ala Ile Lys Thr His Ser Glu Ala Ser Gly Met Val Thr				
	400	405	410	415
cag ctc tga ccggtggctt ctgctgcgc agcctccagg gcccgaggtg				1539
Gln Leu *				
atccccggtgg tgggatccac gctggggcga ggatgggacg tttttcttct tgggccccgt				1599
ccacaggtcc aaggacaccc tccctccagg gtctctctct ccacagtggc gggcccactc				1659
agccccgaga ccacccaacc tcacctctct gacccccatg taaatattgt tctgctgtct				1719
gggactcctg tctaggtgcc cctgatgatg ggatgctctt taaataataa agatggtttt				1779
gatt				1783

<210> 68
 <211> 417
 <212> PRT
 <213> Homo Sapien

<400> 68
 Met Ala Gln Lys Glu Gly Gly Arg Thr Val Pro Cys Cys Ser Arg Pro
 1 5 10 15
 Lys Val Ala Ala Leu Thr Ala Gly Thr Leu Leu Leu Thr Ala Ile
 20 25 30

```

Gly Ala Ala Ser Trp Ala Ile Val Ala Val Leu Leu Arg Ser Asp Gln
      35      40      45
Glu Pro Leu Tyr Pro Val Gln Val Ser Ser Ala Asp Ala Arg Leu Met
      50      55      60
Val Phe Asp Lys Thr Glu Gly Thr Trp Arg Leu Leu Cys Ser Ser Arg
      65      70      75      80
Ser Asn Ala Arg Val Ala Gly Leu Ser Cys Glu Glu Met Gly Phe Leu
      85      90      95
Arg Ala Leu Thr His Ser Glu Leu Asp Val Arg Thr Ala Gly Ala Asn
      100      105      110
Gly Thr Ser Gly Phe Phe Cys Val Asp Glu Gly Arg Leu Pro His Thr
      115      120      125
Gln Arg Leu Leu Glu Val Ile Ser Val Cys Asp Cys Pro Arg Gly Arg
      130      135      140
Phe Leu Ala Ala Ile Cys Gln Asp Cys Gly Arg Arg Lys Leu Pro Val
      145      150      155      160
Asp Arg Ile Val Gly Gly Arg Asp Thr Ser Leu Gly Arg Trp Pro Trp
      165      170      175
Gln Val Ser Leu Arg Tyr Asp Gly Ala His Leu Cys Gly Gly Ser Leu
      180      185      190
Leu Ser Gly Asp Trp Val Leu Thr Ala Ala His Cys Phe Pro Glu Arg
      195      200      205
Asn Arg Val Leu Ser Arg Trp Arg Val Phe Ala Gly Ala Val Ala Gln
      210      215      220
Ala Ser Pro His Gly Leu Gln Leu Gly Val Gln Ala Val Val Tyr His
      225      230      235      240
Gly Gly Tyr Leu Pro Phe Arg Asp Pro Asn Ser Glu Glu Asn Ser Asn
      245      250      255
Asp Ile Ala Leu Val His Leu Ser Ser Pro Leu Pro Leu Thr Glu Tyr
      260      265      270
Ile Gln Pro Val Cys Leu Pro Ala Ala Gly Gln Ala Leu Val Asp Gly
      275      280      285
Lys Ile Cys Thr Val Thr Gly Trp Gly Asn Thr Gln Tyr Tyr Gly Gln
      290      295      300
Gln Ala Gly Val Leu Gln Glu Ala Arg Val Pro Ile Ile Ser Asn Asp
      305      310      315      320
Val Cys Asn Gly Ala Asp Phe Tyr Gly Asn Gln Ile Lys Pro Lys Met
      325      330      335
Phe Cys Ala Gly Tyr Pro Glu Gly Gly Ile Asp Ala Cys Gln Gly Asp
      340      345      350
Ser Gly Gly Pro Phe Val Cys Glu Asp Ser Ile Ser Arg Thr Pro Arg
      355      360      365
Trp Arg Leu Cys Gly Ile Val Ser Trp Gly Thr Gly Cys Ala Leu Ala
      370      375      380
Gln Lys Pro Gly Val Tyr Thr Lys Val Ser Asp Phe Arg Glu Trp Ile
      385      390      395      400
Phe Gln Ala Ile Lys Thr His Ser Glu Ala Ser Gly Met Val Thr Gln
      405      410      415
Leu

```

<210> 69

<211> 2479

<212> DNA

<213> Homo sapien

<220>

<221> CDS

<222> (57)...(1535)

<223> Nucleotide sequence encoding human serine protease

<300>

<308> GenBank U75329

<309> 1997-10-10

<400> 69

```

gtcatattga acattccaga tacctatcat tactcgatgc tgttgataac agcaag atg      59
                                   Met
                                   1

gct ttg aac tca ggg tca cca cca gct att gga cct tac tat gaa aac      107
Ala Leu Asn Ser Gly Ser Pro Pro Ala Ile Gly Pro Tyr Tyr Glu Asn
                    5                                10                    15

cat gga tac caa ccg gaa aac ccc tat ccc gca cag ccc act gtg gtc      155
His Gly Tyr Gln Pro Glu Asn Pro Tyr Pro Ala Gln Pro Thr Val Val
                    20                                25                    30

ccc act gtc tac gag gtg cat ccg gct cag tac tac ccg tcc ccc gtg      203
Pro Thr Val Tyr Glu Val His Pro Ala Gln Tyr Tyr Pro Ser Pro Val
                    35                                40                    45

ccc cag tac gcc ccg agg gtc ctg acg cag gct tcc aac ccc gtc gtc      251
Pro Gln Tyr Ala Pro Arg Val Leu Thr Gln Ala Ser Asn Pro Val Val
                    50                                55                    60                    65

tgc acg cag ccc aaa tcc cca tcc ggg aca gtg tgc acc tca aag act      299
Cys Thr Gln Pro Lys Ser Pro Ser Gly Thr Val Cys Thr Ser Lys Thr
                    70                                75                    80

aag aaa gca ctg tgc atc acc ttg acc ctg ggg acc ttc ctc gtg gga      347
Lys Lys Ala Leu Cys Ile Thr Leu Thr Leu Gly Thr Phe Leu Val Gly
                    85                                90                    95

gct gcg ctg gcc gct ggc cta ctc tgg aag ttc atg ggc agc aag tgc      395
Ala Ala Leu Ala Ala Gly Leu Leu Trp Lys Phe Met Gly Ser Lys Cys
                    100                               105                    110

tcc aac tct ggg ata gag tgc gac tcc tca ggt acc tgc atc aac ccc      443
Ser Asn Ser Gly Ile Glu Cys Asp Ser Ser Gly Thr Cys Ile Asn Pro
                    115                               120                    125

tct aac tgg tgt gat ggc gtg tca cac tgc ccc ggc ggg gag gac gag      491
Ser Asn Trp Cys Asp Gly Val Ser His Cys Pro Gly Gly Glu Asp Glu
                    130                               135                    140                    145

aat cgg tgt gtt cgc ctc tac gga cca aac ttc atc ctt cag atg tac      539
Asn Arg Cys Val Arg Leu Tyr Gly Pro Asn Phe Ile Leu Gln Met Tyr
                    150                               155                    160

tca tct cag agg aag tcc tgg cac cct gtg tgc caa gac gac tgg aac      587
Ser Ser Gln Arg Lys Ser Trp His Pro Val Cys Gln Asp Asp Trp Asn
                    165                               170                    175

gag aac tac ggg cgg gcg gcc tgc agg gac atg ggc tat aag aat aat      635
Glu Asn Tyr Gly Arg Ala Ala Cys Arg Asp Met Gly Tyr Lys Asn Asn
                    180                               185                    190

ttt tac tct agc caa gga ata gtg gat gac agc gga tcc acc agc ttt      683
Phe Tyr Ser Ser Gln Gly Ile Val Asp Asp Ser Gly Ser Thr Ser Phe
                    195                               200                    205

atg aaa ctg aac aca agt gcc ggc aat gtc gat atc tat aaa aaa ctg      731
Met Lys Leu Asn Thr Ser Ala Gly Asn Val Asp Ile Tyr Lys Lys Leu

```

210					215					220					225	
tac	cac	agt	gat	gcc	tgt	tct	tca	aaa	gca	gtg	gtt	tct	tta	cgc	tgt	779
Tyr	His	Ser	Asp	Ala	Cys	Ser	Ser	Lys	Ala	Val	Val	Ser	Leu	Arg	Cys	
				230					235					240		
tta	gcc	tgc	ggg	gtc	aac	ttg	aac	tca	agc	cgc	cag	agc	agg	atc	gtg	827
Leu	Ala	Cys	Gly	Val	Asn	Leu	Asn	Ser	Ser	Arg	Gln	Ser	Arg	Ile	Val	
			245					250					255			
ggc	ggt	gag	agc	gcg	ctc	ccg	ggg	gcc	tgg	ccc	tgg	cag	gtc	agc	ctg	875
Gly	Gly	Glu	Ser	Ala	Leu	Pro	Gly	Ala	Trp	Pro	Trp	Gln	Val	Ser	Leu	
		260					265				270					
cac	gtc	cag	aac	gtc	cac	gtg	tgc	gga	ggc	tcc	atc	atc	acc	ccc	gag	923
His	Val	Gln	Asn	Val	His	Val	Cys	Gly	Gly	Ser	Ile	Ile	Thr	Pro	Glu	
	275					280					285					
tgg	atc	gtg	aca	gcc	gcc	cac	tgc	gtg	gaa	aaa	cct	ctt	aac	aat	cca	971
Trp	Ile	Val	Thr	Ala	Ala	His	Cys	Val	Glu	Lys	Pro	Leu	Asn	Asn	Pro	
	290				295					300					305	
tgg	cat	tgg	acg	gca	ttt	gcg	ggg	att	ttg	aga	caa	tct	ttc	atg	ttc	1019
Trp	His	Trp	Thr	Ala	Phe	Ala	Gly	Ile	Leu	Arg	Gln	Ser	Phe	Met	Phe	
				310					315					320		
tat	gga	gcc	gga	tac	caa	gta	caa	aaa	gtg	att	tct	cat	cca	aat	tat	1067
Tyr	Gly	Ala	Gly	Tyr	Gln	Val	Gln	Lys	Val	Ile	Ser	His	Pro	Asn	Tyr	
				325				330					335			
gac	tcc	aag	acc	aag	aac	aat	gac	att	gcg	ctg	atg	aag	ctg	cag	aag	1115
Asp	Ser	Lys	Thr	Lys	Asn	Asn	Asp	Ile	Ala	Leu	Met	Lys	Leu	Gln	Lys	
		340					345					350				
cct	ctg	act	ttc	aac	gac	cta	gtg	aaa	cca	gtg	tgt	ctg	ccc	aac	cca	1163
Pro	Leu	Thr	Phe	Asn	Asp	Leu	Val	Lys	Pro	Val	Cys	Leu	Pro	Asn	Pro	
		355				360					365					
ggc	atg	atg	ctg	cag	cca	gaa	cag	ctc	tgc	tgg	att	tcc	ggg	tgg	ggg	1211
Gly	Met	Met	Leu	Gln	Pro	Glu	Gln	Leu	Cys	Trp	Ile	Ser	Gly	Trp	Gly	
	370				375					380					385	
gcc	acc	gag	gag	aaa	ggg	aag	acc	tca	gaa	gtg	ctg	aac	gct	gcc	aag	1259
Ala	Thr	Glu	Glu	Lys	Gly	Lys	Thr	Ser	Glu	Val	Leu	Asn	Ala	Ala	Lys	
				390				395						400		
gtg	ctt	ctc	att	gag	aca	cag	aga	tgc	aac	agc	aga	tat	gtc	tat	gac	1307
Val	Leu	Leu	Ile	Glu	Thr	Gln	Arg	Cys	Asn	Ser	Arg	Tyr	Val	Tyr	Asp	
			405					410					415			
aac	ctg	atc	aca	cca	gcc	atg	atc	tgt	gcc	ggc	ttc	ctg	cag	ggg	aac	1355
Asn	Leu	Ile	Thr	Pro	Ala	Met	Ile	Cys	Ala	Gly	Phe	Leu	Gln	Gly	Asn	
		420					425					430				
gtc	gat	tct	tgc	cag	ggt	gac	agt	gga	ggg	cct	ctg	gtc	act	tcg	aac	1403
Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	Thr	Ser	Asn	
						440					445					
aac	aat	atc	tgg	tgg	ctg	ata	ggg	gat	aca	agc	tgg	ggt	tct	ggc	tgt	1451
Asn	Asn	Ile	Trp	Trp	Leu	Ile	Gly	Asp	Thr	Ser	Trp	Gly	Ser	Gly	Cys	
					455					460					465	

gcc aaa gct tac aga cca gga gtg tac ggg aat gtg atg gta ttc acg 1499
 Ala Lys Ala Tyr Arg Pro Gly Val Tyr Gly Asn Val Met Val Phe Thr
 470 475 480

gac tgg att tat cga caa atg aag gca aac ggc taa tccacatggt 1545
 Asp Trp Ile Tyr Arg Gln Met Lys Ala Asn Gly *
 485 490

cttcgtcctt gacgtcgctt tacaagaaaa caatggggct gggtttgctt ccccggtgcat 1605
 gatttactct tagagatgat tcagaggtca cttcattttt attaaacagt gaacttgtct 1665
 ggctttggca ctctctgcca tactgtgcag gctgcagtgg ctccccctgcc cagcctgctc 1725
 tccctaaccc cttgtccgca aggggtgatg gccggctggt tgtgggcaet ggcgggtcaat 1785
 tgtggaagga agaggggttg aggctgcccc cattgagatc ttctgtctga gtcctttcca 1845
 gggggccaatt ttggatgagc atggagctgt cacttctcag ctgctggatg acttgagatg 1905
 aaaaaggaga gacatggaaa gggagacagc caggtggcac ctgcagcggc tgccctcttg 1965
 ggccacttgg tagtgtcccc agcctacttc acaaggggat tttgctgatg ggttcttaga 2025
 gccttagcag ccctggatgg tggccagaaa taaagggacc agcccttcat ggggtggtgac 2085
 gtggtagtca cttgtaaggg gaacagaaac atttttgctt ttatgggggtg agaatataga 2145
 cagtgccctt ggtgcgaggg aagcaattga aaaggaactt gccctgagca ctctggtgc 2205
 aggtctccac ctgcacattg ggtggggctc ctgggagggg gactcagcct tctcctcat 2265
 cctccctgac cctgctccta gcacctgga gagtgaatgc cccttggtcc ctggcagggc 2325
 gccaaagttg gcaccatgtc ggcctcttca ggcctgatag tcattggaaa ttgaggtcca 2385
 tgggggaaat caaggatgct cagtttaagg tacactgttt ccatgttatg tttctacaca 2445
 ttgatgggtg tgacctgag ttcaaagcca tctt 2479

<210> 70
 <211> 492
 <212> PRT
 <213> Homo sapien

<400> 70
 Met Ala Leu Asn Ser Gly Ser Pro Pro Ala Ile Gly Pro Tyr Tyr Glu
 1 5 10 15
 Asn His Gly Tyr Gln Pro Glu Asn Pro Tyr Pro Ala Gln Pro Thr Val
 20 25 30
 Val Pro Thr Val Tyr Glu Val His Pro Ala Gln Tyr Tyr Pro Ser Pro
 35 40 45
 Val Pro Gln Tyr Ala Pro Arg Val Leu Thr Gln Ala Ser Asn Pro Val
 50 55 60
 Val Cys Thr Gln Pro Lys Ser Pro Ser Gly Thr Val Cys Thr Ser Lys
 65 70 75 80
 Thr Lys Lys Ala Leu Cys Ile Thr Leu Thr Gly Thr Phe Leu Val
 85 90 95
 Gly Ala Ala Leu Ala Ala Gly Leu Leu Trp Lys Phe Met Gly Ser Lys
 100 105 110
 Cys Ser Asn Ser Gly Ile Glu Cys Asp Ser Ser Gly Thr Cys Ile Asn
 115 120 125
 Pro Ser Asn Trp Cys Asp Gly Val Ser His Cys Pro Gly Gly Glu Asp
 130 135 140
 Glu Asn Arg Cys Val Arg Leu Tyr Gly Pro Asn Phe Ile Leu Gln Met
 145 150 155 160
 Tyr Ser Ser Gln Arg Lys Ser Trp His Pro Val Cys Gln Asp Asp Trp
 165 170 175
 Asn Glu Asn Tyr Gly Arg Ala Ala Cys Arg Asp Met Gly Tyr Lys Asn
 180 185 190
 Asn Phe Tyr Ser Ser Gln Gly Ile Val Asp Asp Ser Gly Ser Thr Ser
 195 200 205
 Phe Met Lys Leu Asn Thr Ser Ala Gly Asn Val Asp Ile Tyr Lys Lys
 210 215 220
 Leu Tyr His Ser Asp Ala Cys Ser Ser Lys Ala Val Val Ser Leu Arg
 225 230 235 240
 Cys Leu Ala Cys Gly Val Asn Leu Asn Ser Ser Arg Gln Ser Arg Ile

Val	Gly	Gly	Glu	Ser	Ala	Leu	Pro	Gly	Ala	Trp	Pro	Trp	Gln	Val	Ser
			245					250					255		
Leu	His	Val	Gln	Asn	Val	His	Val	Cys	Gly	Gly	Ser	Ile	Ile	Thr	Pro
		275					280					285			
Glu	Trp	Ile	Val	Thr	Ala	Ala	His	Cys	Val	Glu	Lys	Pro	Leu	Asn	Asn
	290						295				300				
Pro	Trp	His	Trp	Thr	Ala	Phe	Ala	Gly	Ile	Leu	Arg	Gln	Ser	Phe	Met
	305				310				315					320	
Phe	Tyr	Gly	Ala	Gly	Tyr	Gln	Val	Gln	Lys	Val	Ile	Ser	His	Pro	Asn
			325						330					335	
Tyr	Asp	Ser	Lys	Thr	Lys	Asn	Asn	Asp	Ile	Ala	Leu	Met	Lys	Leu	Gln
			340					345					350		
Lys	Pro	Leu	Thr	Phe	Asn	Asp	Leu	Val	Lys	Pro	Val	Cys	Leu	Pro	Asn
		355					360					365			
Pro	Gly	Met	Met	Leu	Gln	Pro	Glu	Gln	Leu	Cys	Trp	Ile	Ser	Gly	Trp
	370					375					380				
Gly	Ala	Thr	Glu	Glu	Lys	Gly	Lys	Thr	Ser	Glu	Val	Leu	Asn	Ala	Ala
	385				390					395				400	
Lys	Val	Leu	Leu	Ile	Glu	Thr	Gln	Arg	Cys	Asn	Ser	Arg	Tyr	Val	Tyr
			405						410				415		
Asp	Asn	Leu	Ile	Thr	Pro	Ala	Met	Ile	Cys	Ala	Gly	Phe	Leu	Gln	Gly
		420					425					430			
Asn	Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	Thr	Ser
	435						440					445			
Asn	Asn	Asn	Ile	Trp	Trp	Leu	Ile	Gly	Asp	Thr	Ser	Trp	Gly	Ser	Gly
	450					455				460					
Cys	Ala	Lys	Ala	Tyr	Arg	Pro	Gly	Val	Tyr	Gly	Asn	Val	Met	Val	Phe
	465				470					475				480	
Thr	Asp	Trp	Ile	Tyr	Arg	Gln	Met	Lys	Ala	Asn	Gly				
			485						490						

<210> 71
 <211> 2079
 <212> DNA
 <213> Homo sapien

<220>
 <221> CDS
 <222> (251)...(1522)
 <223> Nucleotide sequence encoding transmembrane
 protease, serine 4 (TMPRSS4)

<300>
 <308> GenBank NM016425
 <309> 2000-11-06

<400> 71
 gagaggcagc agcttggttca gcggacaagg atgctgggcg tgaggggacca aggcctgccc 60
 tgcactcggg cctcctccag ccagtgtgta ccagggaactt ctgacctgct ggccagccag 120
 gacctgtgtg gggaggccct cctgtgcct tggggtgaca atctcagctc caggctacag 180
 ggagaccggg aggatcacag agccagcatg gtacaggatc ctgacagtga tcaacctctg 240
 aacagcctcg atg tca aac ccc tgc gca aac ccc gta tcc cca tgg aga 289
 Met Ser Asn Pro Cys Ala Asn Pro Val Ser Pro Trp Arg
 1 5 10
 cct tca gaa agt gtg ggg atc ccc atc atc ata gca cta ctg agc ctg 337
 Pro Ser Glu Ser Val Gly Ile Pro Ile Ile Ile Ala Leu Leu Ser Leu
 15 20 25
 gcg agt atc atc att gtg gtt gtc ctc atc aag gtg att ctg gat aaa 385
 Ala Ser Ile Ile Ile Val Val Val Leu Ile Lys Val Ile Leu Asp Lys

30	35	40	45	
tac tac ttc ctc tgc ggg cag cct ctc cac ttc atc ccg agg aag cag				433
Tyr Tyr Phe Leu Cys 50	Gly Gln Pro Leu 55	His Phe Ile Pro Arg Lys 60	Gln	
ctg tgt gac gga gag ctg gac tgt ccc ttg ggg gag gac gag gag cac				481
Leu Cys Asp Gly Glu Leu Asp Cys 65	Pro Leu Gly Glu Asp 70	Glu Asp Glu Glu His 75		
tgt gtc aag agc ttc ccc gaa ggg cct gca gtg gca gtc cgc ctc tcc				529
Cys Val Lys Ser Phe Pro Glu Gly 80	Pro Ala Val Ala Val Arg 85	Leu Ser 90		
aag gac cga tcc aca ctg cag gtg ctg gac tcg gcc aca ggg aac tgg				577
Lys Asp Arg Ser Thr Leu Gln Val Leu Asp Ser 95	Ala Thr Gly Asn Trp 100			
ttc tct gcc tgt ttc gac aac ttc aca gaa gct ctc gct gag aca gcc				625
Phe Ser Ala Cys Phe 110	Asn Phe Thr Glu Ala Leu 115	Ala Glu Thr Ala 120		
tgt agg cag atg ggc tac agc agc aaa ccc act ttc aga gct gtg gag				673
Cys Arg Gln Met Gly Tyr Ser Ser Lys 125	Pro Thr Phe Arg Ala Val Glu 130			
att ggc cca gac cag gat ctg gat gtt gtt gaa atc aca gaa aac agc				721
Ile Gly Pro Asp Gln Asp Leu Asp Val 135	Val Glu Ile Thr Glu Asn Ser 140			
cag gag ctt cgc atg cgg aac tca agt ggg ccc tgt ctc tca ggc tcc				769
Gln Glu Leu Arg Met Arg Asn 145	Ser Ser Gly Pro Cys 150	Leu Ser Gly Ser 155		
ctg gtc tcc ctg cac tgt ctt gcc tgt ggg aag agc ctg aag acc ccc				817
Leu Val Ser Leu His Cys 160	Leu Ala Cys Gly Lys 165	Leu Lys Thr Pro 170		
cgt gtg gtg ggt ggg gag gag gcc tct gtg gat tct tgg cct tgg cag				865
Arg Val Val Gly Gly Glu Glu Ala Ser Val Asp Ser Trp Pro Trp Gln 175	180			
gtc agc atc cag tac gac aaa cag cac gtc tgt gga ggg agc atc ctg				913
Val Ser Ile Gln Tyr 185	Lys Gln His Val Cys Gly Gly Ser Ile Leu 190			
gac ccc cac tgg gtc ctc acg gca gcc cac tgc ttc agg aaa cat acc				961
Asp Pro His Trp Val Leu Thr Ala Ala His Cys Phe Arg Lys His Thr 195	200			
gat gtg ttc aac tgg aag gtg cgg gca ggc tca gac aaa ctg ggc agc				1009
Asp Val Phe Asn Trp Lys Val Arg Ala Gly Ser Asp Lys Leu Gly Ser 205	210			
ttc cca tcc ctg gct gtg gcc aag atc atc atc att gaa ttc aac ccc				1057
Phe Pro Ser Leu Ala Val Ala Lys Ile Ile Ile Ile Glu Phe Asn Pro 215	220			
atg tac ccc aaa gac aat gac atc gcc ctc atg aag ctg cag ttc cca				1105
Met Tyr Pro Lys Asp Asn Asp Ile Ala Leu Met Lys Leu Gln Phe Pro 225	230			

```

ctc act ttc tca ggc aca gtc agg ccc atc tgt ctg ccc ttc ttt gat 1153
Leu Thr Phe Ser Gly Thr Val Arg Pro Ile Cys Leu Pro Phe Phe Asp
290 295 300

gag gag ctc act cca gcc acc cca ctc tgg atc att gga tgg ggc ttt 1201
Glu Glu Leu Thr Pro Ala Thr Pro Leu Trp Ile Ile Gly Trp Gly Phe
305 310 315

acg aag cag aat gga ggg aag atg tct gac ata ctg ctg cag gcg tca 1249
Thr Lys Gln Asn Gly Gly Lys Met Ser Asp Ile Leu Leu Gln Ala Ser
320 325 330

gtc cag gtc att gac agc aca cgg tgc aat gca gac gat gcg tac cag 1297
Val Gln Val Ile Asp Ser Thr Arg Cys Asn Ala Asp Asp Ala Tyr Gln
335 340 345

ggg gaa gtc acc gag aag atg atg tgt gca ggc atc ccg gaa ggg ggt 1345
Gly Glu Val Thr Glu Lys Met Met Cys Ala Gly Ile Pro Glu Gly Gly
350 355 360 365

gtg gac acc tgc cag ggt gac agt ggt ggg ccc ctg atg tac caa tct 1393
Val Asp Thr Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Tyr Gln Ser
370 375 380

gac cag tgg cat gtg gtg ggc atc gtt agc tgg ggc tat ggc tgc ggg 1441
Asp Gln Trp His Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Gly
385 390 395

ggc ccg agc acc cca gga gta tac acc aag gtc tca gcc tat ctc aac 1489
Gly Pro Ser Thr Pro Gly Val Tyr Thr Lys Val Ser Ala Tyr Leu Asn
400 405 410

tgg atc tac aat gtc tgg aag gct gag ctg taa tgctgtgtgcc cctttgcagt 1542
Trp Ile Tyr Asn Val Trp Lys Ala Glu Leu *
415 420

gctgggagcc gcttccttcc tgccctgccc acctggggat ccccaaagt cagacacaga 1602
gcaagagtcc ccttgggtac acccctctgc ccacagcctc agcatttctt ggagcagcaa 1662
agggcctcaa ttctgtgaag agaccctcgc agcccagagg cgcccagagg aagtcagcag 1722
ccctagctcg gccacacttg gtgtctccag catcccaggg agagacacag cccactgaac 1782
aaggtctcag gggatttgct aagccaagaa ggaactttcc cacactactg aatggaagca 1842
ggctgtcttg taaaagccca gatcactgtg ggctggagag gagaaggaaa ggtctgcgc 1902
cagccctgtc cgtcttcacc catccccaag cctactagag caagaaacca gttgtaatat 1962
aaaatgcact gccctactgt tggtatgact accgttacct actgttgtca ttgttattac 2022
agctatggcc actattatta aagagctgtg taacatcaaa aaaaaaaaaa aaaaaaa 2079

```

<210> 72

<211> 423

<212> PRT

<213> Homo sapien

<400> 72

```

Met Ser Asn Pro Cys Ala Asn Pro Val Ser Pro Trp Arg Pro Ser Glu
1 5 10 15
Ser Val Gly Ile Pro Ile Ile Ala Leu Leu Ser Leu Ala Ser Ile
20 25 30
Ile Ile Val Val Leu Ile Lys Val Ile Leu Asp Lys Tyr Tyr Phe
35 40 45
Leu Cys Gly Gln Pro Leu His Phe Ile Pro Arg Lys Gln Leu Cys Asp
50 55 60
Gly Glu Leu Asp Cys Pro Leu Gly Glu Asp Glu His Cys Val Lys
65 70 75 80

```

Ser Phe Pro Glu Gly Pro Ala Val Ala Val Arg Leu Ser Lys Asp Arg
 85 90 95
 Ser Thr Leu Gln Val Leu Asp Ser Ala Thr Gly Asn Trp Phe Ser Ala
 100 105 110
 Cys Phe Asp Asn Phe Thr Glu Ala Leu Ala Glu Thr Ala Cys Arg Gln
 115 120 125
 Met Gly Tyr Ser Ser Lys Pro Thr Phe Arg Ala Val Glu Ile Gly Pro
 130 135 140
 Asp Gln Asp Leu Asp Val Val Glu Ile Thr Glu Asn Ser Gln Glu Leu
 145 150 155 160
 Arg Met Arg Asn Ser Ser Gly Pro Cys Leu Ser Gly Ser Leu Val Ser
 165 170 175
 Leu His Cys Leu Ala Cys Gly Lys Ser Leu Lys Thr Pro Arg Val Val
 180 185 190
 Gly Gly Glu Glu Ala Ser Val Asp Ser Trp Pro Trp Gln Val Ser Ile
 195 200 205
 Gln Tyr Asp Lys Gln His Val Cys Gly Gly Ser Ile Leu Asp Pro His
 210 215 220
 Trp Val Leu Thr Ala Ala His Cys Phe Arg Lys His Thr Asp Val Phe
 225 230 235 240
 Asn Trp Lys Val Arg Ala Gly Ser Asp Lys Leu Gly Ser Phe Pro Ser
 245 250 255
 Leu Ala Val Ala Lys Ile Ile Ile Ile Glu Phe Asn Pro Met Tyr Pro
 260 265 270
 Lys Asp Asn Asp Ile Ala Leu Met Lys Leu Gln Phe Pro Leu Thr Phe
 275 280 285
 Ser Gly Thr Val Arg Pro Ile Cys Leu Pro Phe Phe Asp Glu Glu Leu
 290 295 300
 Thr Pro Ala Thr Pro Leu Trp Ile Ile Gly Trp Gly Phe Thr Lys Gln
 305 310 315 320
 Asn Gly Gly Lys Met Ser Asp Ile Leu Leu Gln Ala Ser Val Gln Val
 325 330 335
 Ile Asp Ser Thr Arg Cys Asn Ala Asp Asp Ala Tyr Gln Gly Glu Val
 340 345 350
 Thr Glu Lys Met Met Cys Ala Gly Ile Pro Glu Gly Gly Val Asp Thr
 355 360 365
 Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Tyr Gln Ser Asp Gln Trp
 370 375 380
 His Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys Gly Gly Pro Ser
 385 390 395 400
 Thr Pro Gly Val Tyr Thr Lys Val Ser Ala Tyr Leu Asn Trp Ile Tyr
 405 410 415
 Asn Val Trp Lys Ala Glu Leu
 420

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57194 A3

- (51) International Patent Classification⁷: **C12N 9/64**,
15/57, C07K 16/40, C12Q 1/37, C07K 19/00, A61K 38/48
- (21) International Application Number: **PCT/US01/03471**
- (22) International Filing Date: 2 February 2001 (02.02.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|--------------------------------|----|
| 60/179,982 | 3 February 2000 (03.02.2000) | US |
| 60/183,542 | 18 February 2000 (18.02.2000) | US |
| 60/213,124 | 22 June 2000 (22.06.2000) | US |
| 60/220,970 | 26 July 2000 (26.07.2000) | US |
| 09/657,986 | 8 September 2000 (08.09.2000) | US |
| 60/234,840 | 22 September 2000 (22.09.2000) | US |
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
- | | |
|----------|--------------------------------|
| US | 60/179,982 (CIP) |
| Filed on | 3 February 2000 (03.02.2000) |
| US | 60/183,542 (CIP) |
| Filed on | 18 February 2000 (18.02.2000) |
| US | 60/213,124 (CIP) |
| Filed on | 22 June 2000 (22.06.2000) |
| US | 60/220,970 (CIP) |
| Filed on | 26 July 2000 (26.07.2000) |
| US | 09/657,986 (CIP) |
| Filed on | 8 September 2000 (08.09.2000) |
| US | 60/234,840 (CIP) |
| Filed on | 22 September 2000 (22.09.2000) |
- (71) Applicant (for all designated States except US): **CORVAS INTERNATIONAL, INC.** [US/US]; 3030 Science Park Road, San Diego, CA 92121 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MADISON, Edwin, L.** [US/US]; 501 Stratford Court, Del Mar, CA 92014 (US). **ONG, Edgar, O.** [CA/US]; 8328 Regents Road #3C, San Diego, CA 92122 (US). **YEH, Jiunn-chen** [—/US]; 3128-E Via Alicante Drive, La Jolla, CA 92037 (US).
- (74) Agents: **SEIDMAN, Stephanie, L.** et al.; Heller Ehrman White & McAuliffe LLP, Suite 700, 4250 Executive Square, La Jolla, CA 92037 (US).
- (81) Designated States (national): AE, AG, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
28 March 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID MOLECULES ENCODING TRANSMEMBRANE SERINE PROTEASES, THE ENCODED PROTEINS AND METHODS BASED THEREON

(57) Abstract: Provided herein are polypeptides that include the protease domain of a type II transmembrane serine protease (MTSP) as a single chain. Methods using the polypeptides to identify compounds that modulate the protease activity of an MTSP are provided. Also provided are MTSPs designated MTSP3 and MTSP4 and a form of an MTSP designated MTSP6.

WO 01/57194 A3

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/03471

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/64 C12N15/57 C07K16/40 C12Q1/37 C07K19/00
A61K38/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, BIOSIS, MEDLINE, EMBL, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 42120 A (UNIV ARKANSAS) 26 August 1999 (1999-08-26)	1-35, 38-41, 105,106, 130,132, 142
Y	the whole document	42-54, 137-139
X	WO 99 36550 A (INCYTE PHARMA INC ;CORLEY NEIL C (US); YUE HENRY (US); BANDMAN OLG) 22 July 1999 (1999-07-22) SEQ ID NO 6 and 18 page 21, line 22 -page 22, line 7 -/-	55-68, 70, 105-108, 111-125, 133-135

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 August 2001

Date of mailing of the international search report

11.09.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bilang, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/03471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LIN C-Y ET AL: "Molecular cloning of cDNA for Matriptase, a matrix-degrading serine protease with trypsin-like activity" JOURNAL OF BIOLOGICAL CHEMISTRY, US, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 274, no. 26, 25 June 1999 (1999-06-25), pages 8231-18236, XP002139963 ISSN: 0021-9258 cited in the application	1-35, 38-41, 105, 106, 130, 132, 142
Y	the whole document	42-54, 137-139
X	--- DATABASE EMBL 'Online! Accession Number W22987, 8 October 1997 (1997-10-08) SUNTORY LTD: "Human Serine Protease 67" XP002169836 abstract & PATENT ABSTRACTS OF JAPAN vol. 1997, no. 10, 31 October 1997 (1997-10-31) & JP 09 149790 A (SUNTORY LTD), 10 June 1997 (1997-06-10) abstract	1-9, 11-15, 21-35, 39-41
X	--- DATABASE EMBL 'Online! Accession Number AAY41710, 7 December 1999 (1999-12-07) GENENTECH INC.: "Human PRO618 protein sequence" XP002175683 abstract & WO 99 46281 A (GENENTECH INC.) 16 September 1999 (1999-09-16)	71-87, 89, 105-108, 111
X	--- DATABASE EMBL 'Online! Accession Number AAZ34033, 7 December 1999 (1999-12-07) GENENTECH INC.: "Human PRO618 nucleotide sequence" XP002175684 abstract & WO 99 46281 A (GENENTECH INC.) 16 September 1999 (1999-09-16)	71-87, 89, 105-108, 111
	--- -/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/03471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! Accession Number AAZ33949, 7 December 1999 (1999-12-07) GENENTECH INC.: "Human PR0382 nucleotide sequence" XP002175685 abstract & WO 99 46281 A (GENENTECH INC.) 16 September 1999 (1999-09-16)	90-102, 104-108, 111, 126-129
X	DATABASE EMBL 'Online! Accession Number AAY41694, 7 December 1999 (1999-12-07) GENENTECH INC.: "Human PR0382 protein sequence" XP002175687 abstract & WO 99 46281 A (GENENTECH INC.) 16 September 1999 (1999-09-16)	90-102, 104-108, 111, 126-129
X	DATABASE EMBL 'Online! Accession Number AI469095, 17 March 1999 (1999-03-17) NCI-CGAP: "tm06c09.x1 Homo sapiens cDNA clone IMAGE:2155792" XP002175686 abstract	90-102, 104-108, 111, 126-129
X	TAKEUCHI TOSHIHIKO ET AL: "Reverse biochemistry: Use of macromolecular protease inhibitors to dissect complex biological processes and identify a membrane-type serine protease in epithelial cancer and normal tissue." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 96, no. 20, 28 September 1999 (1999-09-28), pages 11054-11061, XP002169835 ISSN: 0027-8424 cited in the application	105,106
Y	figure 1	1-35, 38-54, 137-139, 142
	-/-	

INTERNATIONAL SEARCH REPORT

Inter national Application No
PCT/US 01/03471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	VALLIE LA E R ET AL: "CLONING AND FUNCTIONAL EXPRESSION OF A CDNA ENCODING THE CATALYTIC SUBUNIT OF BOVINE ENTEROKINASE" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 268, no. 31, 5 November 1993 (1993-11-05), pages 23311-23317, XP000999230 ISSN: 0021-9258 the whole document	1-35, 38-54, 137-142
Y	LU DESHUN ET AL: "Bovine proenteropeptidase is activated by trypsin, and the specificity of enteropeptidase depends on the heavy chain." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 50, 12 December 1997 (1997-12-12), pages 31293-31300, XP000999222 ISSN: 0021-9258 the whole document	1-35, 38-54, 137-142
P,X	WO 01 04141 A (MERCK PATENT GMBH ;HOFMANN UWE (DE); WILBERT OLIVER (DE); MATZKU S) 18 January 2001 (2001-01-18)	55-68, 70, 105-108, 111-125, 133-135, 140-142
P,Y	the whole document	1-35, 38-54, 137-139
P,X	WO 00 52044 A (UNIV ARKANSAS) 8 September 2000 (2000-09-08)	90-102, 104-108, 111-129, 133-135, 140-142
Y	the whole document	1-35, 38-54, 137-139
-/-		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/03471

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>TAKEUCHI TOSHIHIKO ET AL: "Cellular localization of membrane-type serine protease 1 and identification of protease-activated receptor-2 and single-chain urokinase-type plasminogen activator as substrates." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 275, no. 34, 25 August 2000 (2000-08-25), pages 26333-26342, XP001000977 ISSN: 0021-9258 cited in the application the whole document</p>	<p>1-35, 38-54, 105,106, 130,132, 137-139, 142</p>
P,A	<p>LEE SHEAU-LING ET AL: "Activation of hepatocyte growth factor and urokinase/plasminogen activator by matriptase, an epithelial membrane serine protease." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 275, no. 47, 24 November 2000 (2000-11-24), pages 36720-36725, XP000999224 ISSN: 0021-9258 the whole document</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/03471

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 16-18, 36, 37, 69, 88, 103, 109, 110, 136
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
1-54, 130, 132, 137-139, 142 (all partially), 55-129, 131, 133-136, 140, 141 (completely)
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-54, 105, 106, 130, 132, 137-139,
142 (all partially)

Polypeptide comprising the protease domain of Type-II
membrane type serine protease MTSP1, nucleic acid encoding
said polypeptide, and uses thereof.

2. Claims: 1-54, 105-125, 130, 131, 133-139,
140-142 (all partially), 55-70 (completely)

As 1., but MTSP3

3. Claims: 1-54, 105-125, 130, 131, 133-139,
140-142 (all partially), 71-89 (completely)

As 1., but MTSP4

4. Claims: 1-54, 105-125, 130, 131, 133-139,
140-142 (all partially), 90-104,
126-129 (completely)

As 1., but MTSP6

5. Claims: 1-54, 130, 132, 137-139, 142 (all partially)

As 1., but corin

6. Claims: 1-54, 130, 132, 137-139, 142 (all partially)

As 1., but enterpeptidase

7. Claims: 1-54, 130, 132, 137-139, 142 (all partially)

As 1., but human airway trypsin-like protease

8. Claims: 1-54, 130, 132, 137-139, 142 (all partially)

As 1., but TMPRSS2

9. Claims: 1-54, 130, 132, 137-139, 142 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

As 1., but TMRSS4

10. Claims: 42-54, 137-139

Methods of identifying compounds that modulate the activity of MTSP or bind to single chain protease domain of MTSP, the method comprising the step of using the single chain protease domain of MTSP.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 16-18, 36, 37, 69, 88, 103, 109, 110, 136

Present claims 16 to 18 relate to a product defined by reference to a desirable characteristic or property, namely being a mutein of the polypeptide of claim 1 having a particular catalytic activity. The claims cover all products having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such products. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the product by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the muteins having a Cys in the catalytic domain replaced with another amino acid.

Present claims 36, 37, 69, 88, 103, 109, 110, 136 relate to a product/compound (or the use thereof) defined by reference to a desirable characteristic or property, namely being an inhibitor or modulator of the catalytic activity of type-II membrane-type serine proteases. The same reasoning as outlined above for claims 16 to 18 apply. The search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to antibodies or antisense oligonucleotides.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/03471

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9942120 A	26-08-1999	US 5972616 A AU 2869299 A EP 1056466 A	26-10-1999 06-09-1999 06-12-2000
WO 9936550 A	22-07-1999	US 6203979 B AU 2113599 A EP 1045913 A	20-03-2001 02-08-1999 25-10-2000
WO 9946281 A	16-09-1999	AU 3072199 A AU 3075099 A EP 1064382 A WO 9947677 A AU 735081 B AU 1532499 A EP 1032667 A WO 9927098 A AU 3757099 A EP 1071773 A WO 9954467 A AU 1070399 A EP 1025227 A WO 9920756 A	27-09-1999 11-10-1999 03-01-2001 23-09-1999 28-06-2001 15-06-1999 06-09-2000 03-06-1999 08-11-1999 31-01-2001 28-10-1999 10-05-1999 09-08-2000 29-04-1999
WO 0104141 A	18-01-2001	NONE	
WO 0052044 A	08-09-2000	AU 3720900 A	21-09-2000